#### THE USE OF IONIC LIQUIDS AS A PHASE TRANSFER CATALYST FOR THE DIALKYLATION OF AN ACTIVE METHYLENE COMPOUND

#### NUR LIYANA BINTI SHAMSUDEEN

BACHELOR OF SCIENCE (Hons.) CHEMISTRY FACULTY OF APPLIED SCIENCES UNIVERSITI TEKNOLOGI MARA

**APRIL 2009** 

This Final Year Project Report entitled "The Use of Ionic Liquids as a Phase Transfer Catalyst for the Dialkylation of an Active Methylene Compound" was submitted by Nur Liyana binti Shamsudeen, in partial fulfilment of the requirements for the Degree of Bachelor of Science (Hons.) Chemistry, in the Faculty of Applied Sciences, and was approved by

Prof. Dr.Ahmad Sazali Hamzah Supervisor B.Sc. (Hons.) Chemistry Faculty of Applied Sciences Universiti Teknologi MARA 40450 Shah Alam Selangor

Dr. Famiza Abdul Latiff Project Coordinator B.Sc. (Hons.) Chemistry Faculty of Applied Sciences Universiti Teknologi MARA 40450 Shah Alam Selangor

Prof. Madya Dr. Faizah Md Salleh Head of Programme B.Sc. (Hons.) Chemistry Faculty of Applied Sciences Universiti Teknologi MARA 40450 Shah Alam Selangor

7 MAY 2009 Date:

#### **TABLE OF CONTENTS**

iii
iv
vi
vii
ix
Х
xi

#### **CHAPTER 1 INTRODUCTION**

1.1	Background of Phase Transfer Catalyst	1
1.2	Introduction of ionic liquids	2
1.3	Introduction of methylacetoacetate	4
1.4	Chromatography techniques in organic synthesis	5
1.5	Problem statement of study	8
1.6	Significance of study	8
1.7	Objectives of study	8

#### **CHAPTER 2 LITERATURE REVIEW**

2.1	Previous work on the dialkylation of ethylacetoacetate using ionic liquid	9
2.2	Previous work on alkylation of diethyl malonate using ionic liquid	12

## **CHAPTER 3 METHODOLOGY**

3.1	Genera	l Experimental Procedures	14
	3.1.1	Purification of Reagents, Compounds and Solvents	14
	3.1.2	Preparation of glassware	15
	3.1.3	Instruments for data analysis	15
3.2	Experimental		17
	3.2.1	Dialkylation of methylacetoacetate with methyl iodide in the presence of $[bmim]BF_4$	17
	3.2.2	Dialkylation of methylacetoacetate with methyl iodide in the presence of $[bmim]PF_6^-$	18
	3.2.3	Dialkylation of methylacetoacetate with propyl iodide in the presence of $[bmim]BF_4$	19

### **CHAPTER 4 RESULTS AND DISCUSSION**

4.1	Synthe	sis of alkylated compounds via dialkylation reaction	21
	4.1.1	Dialkylation of methylacetoacetate with methyl iodide (CH <sub>3</sub> I)	21
		and propyl iodide ( $C_3H_7I$ ) by using phase transfer catalysts	
	4.1.2	Results for the dialkylation of methylacetoacetate in different	26
		ionic liquids	
4.2	Spectro	oscopic characterization for the dialkylation of methylacetoacetate	27
4.3	NMR (	characterization for the dialkylation of methylacetoacetate	28
	4.3.1	Dialkylation with methyl iodide in the presence of [bmim]BF <sub>4</sub>	28
	4.3.2	Dialkylation with methyl iodide in the presence of $[bmim]PF_6$	34
	4.3.3	Dialkylation with propyl iodide in the presence of $[bmim]BF_4$	40
СНА	PTER 5	5 CONCLUSION AND RECOMMENDATIONS	46

CITED REFERENCES	48
APPENDICES	49
CURRICULUM VITAE	50

#### ABSTRACT

# THE USE OF IONIC LIQUIDS AS A PHASE TRANSFER CATALYST FOR THE DIALKYLATION OF AN ACTIVE METHYLENE COMPOUND.

A study on the use of ionic liquids as a phase transfer catalysts on the dialkylation of methylacetoacetate has given a new finding for the development of environmentally friendly catalysts. The structures of these dialkylated compounds were elucidated using modern spectroscopic techniques such as IR and NMR. Studies on the dialkylation of methylacetoatate with different elecrophiles had been carried out in the presence of different ionic liquids. The reaction mixture was left stirring overnight at room temperature under nitrogen gas, N<sub>2</sub> until all compounds dissolved. The result showed that ionic liquid which was 1-butyl-3-methylimidazolium tetrafluoroborate, [bmim]BF<sub>4</sub><sup>-</sup> as the most successful phase transfer catalyst for the dialkylation of methylacetoacetate as compared to 1-butyl-3-methylimidazolium hexafluorophosphate, [bmim]PF<sub>6</sub><sup>-</sup>. With higher percentage yield obtained from ionic liquid of [bmim]BF<sub>4</sub><sup>-</sup>, it certainly showed that this ionic liquid has potential for further investigations in other chemical transformations.