


VOLUME 8 NO. 2
DEC 2011

ISSN 1675-7009

SCIENTIFIC RESEARCH JOURNAL



RESEARCH MANAGEMENT INSTITUTE



SCIENTIFIC RESEARCH JOURNAL

Chief Editor

Zaiki Awang
Universiti Teknologi MARA, Malaysia

Managing Editor

Razidah Ismail
Universiti Teknologi MARA, Malaysia

Editorial Board

Abu Bakar Abdul Majeed, Universiti Teknologi MARA, Malaysia
David Shallcross, University of Melbourne, Australia
Halila Jasmani, Universiti Teknologi MARA, Malaysia
Hamidah Mohd. Saman, Universiti Teknologi MARA, Malaysia
Huang Siew Lai, Universiti Teknologi MARA, Malaysia
Ichsan Setya Putra, Bandung Institute of Technology, Indonesia
Ideris Zakaria, Universiti Malaysia Pahang, Malaysia
Jamil Salleh, Universiti Teknologi MARA, Malaysia
K. Ito, Chiba University, Japan
Kartini Kamaruddin, Universiti Teknologi MARA, Malaysia
Luciano Boggione, University of Massachusetts Lowell, USA
Mohd Hanapiah Abidin, Universiti Teknologi MARA, Malaysia
Mohd Jamaludin Md Noor, Universiti Teknologi MARA, Malaysia
Mohd Rozi Ahmad, Universiti Teknologi MARA, Malaysia
Mohd. Nasir Taib, Universiti Teknologi MARA, Malaysia
Mohd Zamin Jumaat, University of Malaya, Malaysia
Muhammad Azmi Ayub, Universiti Teknologi MARA, Malaysia
Norashikin Saim, Universiti Teknologi MARA, Malaysia
Noriham Abdullah, Universiti Teknologi MARA, Malaysia
Nordin Abu Bakar, Universiti Teknologi MARA, Malaysia
Robert Michael Savory, Universiti Teknologi MARA, Malaysia
Saadiah Yahya, Universiti Teknologi MARA, Malaysia
Salmiah Kasolang, Universiti Teknologi MARA, Malaysia
Shah Rizam Mohd. Shah Baki, Universiti Teknologi MARA, Malaysia
Suhaimi Abd. Talib, Universiti Teknologi MARA, Malaysia
Titik Khawa Abd. Rahman, Universiti Teknologi MARA, Malaysia
Vasudeo Zambare, South Dakota School of Mines and Technology, USA
Wahyu Kuntjoro, Universiti Teknologi MARA, Malaysia
Zahrah Ahmad, University of Malaya, Malaysia

Journal Administrator

Puteri Murni Bt Salleh Hudin
Universiti Teknologi MARA, Malaysia

© UiTM Press, UiTM 2011

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means; electronics, mechanical, photocopying, recording or otherwise; without prior permission in writing from the Director of UiTM Press, Universiti Teknologi MARA, 40450 Shah Alam, Selangor Darul Ehsan, Malaysia. e-mail: penerbit@salam.uitm.edu.my

Scientific Research Journal is jointly published by Research Management Institute (RMI) and UiTM Press, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

The views and opinion expressed therein are those of the individual authors and the publication of these statements in the Scientific Research Journal do not imply endorsement by the publisher or the editorial staff. Copyright is vested in Universiti Teknologi MARA. Written permission is required to reproduce any part of this publication.

SCIENTIFIC RESEARCH JOURNAL

Vol. 8 No. 2

December 2011

ISSN 1675-7009

1. **Electrical Resistivity and Flexural Strength of Plastic Composite Reinforced with Pineapple Leaf Particles** 1
Shahril Anuar Bahari
Mohd Khairi Yahya
Masitah Abu Kassim
Khairul Safuan Muhammad
Rahimi Baharom

2. **Removal of Cd²⁺ from Aqueous Solution Using MgAlCO₃-Layered Double Hydroxide** 13
Nurul Izza Taib
Norzuyanti Mohd Aris

3. **Neurotoxicity and Neuroprotective Effects of Polyimides (PI) and Polyphenylenevinylene (PPV) against Hydrogen Peroxide (H₂O₂)** 33
Norfaezah Mazalan
Mazatulikhma Mat Zain
Nor Saliyana Jumali
Norhanim Mohalid
Zurina Shameri
Ahmad Sazali Hamzah

4. Palm Olein Corrosion Inhibitor for Aluminium in HCl 49
Solution

Junaidah Jai

Wan Shabuddin Wan Ali

Neurotoxicity and Neuroprotective Effects of Polyimides (PI) and Polyphenylenevinylene (PPV) against Hydrogen Peroxide (H_2O_2)

Norfaezah Mazalan¹, Mazatulikhma Mat Zain, Nor Saliyana Jumali, Norhanim Mohalid, Zurina Shameri and Ahmad Sazali Hamzah

Institute of Science,
Faculty of Applied Sciences, Universiti Teknologi MARA,
45000 Shah Alam, Selangor, Malaysia.
¹Email: faezahmazalan@yahoo.com

ABSTRACT

Recently, research and development in the field of drug delivery systems (DDS) facilitating site-specific therapy has reached significant progression. DDS based on polymer micelles, coated micro- and nanoparticles, and various prodrug systems including water-soluble polymer have been prepared and extensively studied as novel drugs designed for cancer chemotherapy and brain delivery. Since polymers are going to be used in human, this study has the interest of testing two types of polymer, polyimides (PI) and polyphenylenevinylene (PPV) on neuronal cells. The objective of this study was to determine the possible neurotoxicity and potential neuroprotective effects of PI and PPV towards SH-SY5Y neuronal cells challenged by hydrogen peroxide (H_2O_2) as an oxidant. Cells were pretreated with either PI or PPV for 1 hour followed by incubation for 24 hour with 100 μM of H_2O_2 . MTS assay was used to assess cell viability. Results show that PI and PPV are not harmful within the concentration up to 10 μM and 100 μM , respectively. However, PI and PPV do not protect neuronal cells against toxicity induced by H_2O_2 or further up the cell death.

Keywords: Neurotoxicity, Neuroprotection, Polyimide (PI), Polyphenylenevinylene (PPV), and Hydrogen Peroxide

Introduction

Over the last decade, there have been major advances in understanding the mechanisms involved in the development of central nervous system (CNS) disorders. Simultaneous with these events, huge progress has been made by researchers to identify the function of peptides and proteins as key molecules in the regulation of most physiological processes, both cellular and the molecular level. Peptides and proteins have turned out to be the main targets in neuropharmaceutical drug design for the treatment of a broad range of CNS disorders for example ischemia, inflammatory CNS diseases, neurodegenerative diseases, and acute and chronic pain syndromes [1 – 6]. Nevertheless, the inability of many existing peptide and protein drugs (P/P drugs) to effectively deliver and sustain within the brain have turned them to be ineffective for the treatment of these clinical problems. The main obstacle to targeting the brain with therapeutics in general (P/P drugs amongst them) is the presence of various barriers, in the first line the blood-brain barrier (BBB), which controls the concentration and entry of solutes into the CNS.

Currently, there is no effective treatment for the majority of brain diseases and it is a great challenge in designing potential P/P drugs with improved potency, selectivity and therapeutic efficacy for treating CNS disorders. The failure of treatment in some cases is due to the inability of P/P drugs to cross the BBB, which represents a limiting factor for the translation of drug research into clinical reality and commercialization. Other undetermined importance factors include the low stability and half-life of the P/P drugs in the blood, their non-selective accumulation, short residence time in the CNS and unspecific distribution of P/P drugs to target sites inside the brain [7].

Nowadays, research and development in the field of drug delivery systems (DDS) facilitating site-specific therapy has achieved significant progression. Safe and nontoxic formulations of a cytotoxic drug based on a polymer, its site-specific delivery and specific activation of the cytotoxic biologically active compound at its target (tumor tissue or tumor cell lines), have become the major goal of such research. DDS based on polymer micelles [8], coated micro- and nanoparticles [9], and various prodrug systems [10] including water-soluble polymer have been prepared and extensively studied as novel drugs designed for cancer chemotherapy and brain delivery [11, 12]. Since polymers are going to be used in human,

this research explored the effects of two types of polymers, polyimides (PI) and polyphenylenevinylene (PPV) on neuroblastoma cell line, SH-SY5Y. Generally, the toxicity of these polymers is identified before proceeding to the neuroprotective test because PI and PPV are considered safe to be used if they are not toxic to the cells. This study used hydrogen peroxide (H_2O_2), as an oxidative stress that induced cell death.




Figure 1: Chemical Structure of (a) Polyimides (PI). (b) Poly(1,4-bis(bromomethyl)-2-butoxy-5-methoxybenzene) (MBu-PPV). (c) Poly(2-Methoxy-5-(2'-Ethyl-Hexyloxy)-p-Phenylenevinylene) (MEH-PPV)

Materials and methods

Materials

Polyimides (PI) and polyphenylenevinylene (PPV) and its derivatives; poly(1,4-bis(bromomethyl)-2-butoxy-5-methoxybenzene) (MBu-PPV) and poly(2-Methoxy-5-(2'-Ethyl-Hexyloxy)-*p*-Phenylenevinylene) (MEH-PPV) were synthesized by Organic Synthesis and Biology laboratory, Institute of Science (IOS), Universiti Teknologi MARA (UiTM) Shah Alam, Selangor.

Cell Culture and Treatments

Human neuroblastoma SH-SY5Y cell lines were purchased from American Type Culture Collection (ATCC). The cells were adapted to grow in Dulbecco's modified Eagle's medium (DMEM) pH 7.4. 2×10^4 cells per ml were seeded in 96 well plates. The cells were differentiated to neural phenotype by adding retinoic acid (RA). After 24 hour, RA was added at a final concentration of 10 μ M in MEM-F12 supplemented with 1 % glutamine, 1 % non-essential amino acids, 1 % gentamicin and 10 % heat activated fetal bovine serum in a humidified atmosphere containing CO₂ at 37 °C. The media was changed after 3 days and cells were used at day 6.

Analysis of H₂O₂ Toxicity (IC₅₀ Determination)

Dilutions of H₂O₂ in DMEM were made fresh just prior to each experiment. Small volumes of H₂O₂ were added to each well to make final dilutions of 1 nM, 10 nM, 100 nM, 1 μ M, 10 μ M, 100 μ M, 1 mM, 10 mM, and 100 mM. The wells were then agitated lightly. Cultures were incubated in a humidified incubator at 37 °C, 5 % CO₂ with H₂O₂ for 24 hour. Cell viability was assessed by MTS assay. The concentration that caused 50 % reduction of cell viability was determined.

Analysis of Neurotoxicity of PI and PPV

Prior to each experiment, PI and PPV dilutions were made fresh in MEM-F12. Small volumes of PI were added to each well to make final dilutions of 1 ng/ml, 10 ng/ml, 100 ng/ml, 1 μ g/ml, 10 μ g/ml, 100 μ g/ml, 1 mg/ml and 10 mg/ml, while PPV was diluted to make the concentrations 1 ng/ml, 10 ng/ml, 100 ng/ml, 1 μ g/ml, 10 μ g/ml, and 100 μ g/ml. The wells

were agitated lightly. Cultures were incubated in a humidified incubator at 37 °C, 5 % CO₂ for 24 hour, and with PI or PPV for 1 hour and cell viability was assessed by MTS reduction assay.

Analysis of Neuroprotective Effects of PI and PPV Against H₂O₂-induced Cell Death

Differentiated SH-SY5Y cells were incubated with PI or PPV for 1 hour prior incubation with 100 μM H₂O₂ that cause 50 % cell viability (IC₅₀ determination earlier) for 24 h. Cell viability was assessed by MTS assay.

Analysis of Cell Viability

Cell viability was assessed with Cell Titer 96* Aqueous Non-Radioactive Cell Proliferation Assay (MTS, Promega, USA). The MTS assay is a colorimetric assay based on the reduction of 3-(4,5-dimethylthiazol-2-yl)-5-(3 carboxymethoxyphenyl)-2-(4 sulfophenyl)-2H-tetrazolium (MTS) by dehydrogenase enzymes found in metabolically active cells (Chemicon, USA). The amount of colored formazan product is proportional to the number of viable cells. Briefly, 20 μl MTS solution was added to each well and incubated in humidified incubator at 37 °C in 5 % CO₂ for 2 - 4 hours in the dark. The quantity of formazan product present was determined by measuring the absorbance at 490 nm with a micro titer plate reader (GloMax Integrated Systems by Promega USA). Values were expressed as the percentage of optical density of control cells in the absence of H₂O₂ and/or PI or PPV.

Statistical Analysis

Each experiment was carried out in triplicates with at least 3 independent cultures with comparable results. Data was reported as mean ±SD of at least three experiments. Comparison between control and treatment of PI and PPV were made by Student's t-test. *P* < 0.05 was considered statistically significant.

Results and Discussion

H₂O₂ Toxicity

The human SH-SY5Y cell line, derived from child neuroblastoma, is commonly used as an *in vitro* model to study the related neurodegenerative diseases such as Parkinson's and Alzheimer's disease [13, 14]. This is due to the fact that the cells express several markers for those diseases, such as dopaminergic markers, including tyrosine hydroxylase (TH) and the dopamine transporter (DAT) [15, 16]. However, in undifferentiated state, the cells are immature neuroblasts that maintain the characteristics of stem cell and aggressively proliferate for a long period of time [15, 16]. Because of this reason, the cells are differentiated with retinoic acid (RA), either alone, or in combination with other agents, such as growth factors and phorbol ester, to stop mitosis and enhance the neuronal and dopaminergic characteristics of the cells [15, 17]. In this study, the SH-SY5Y cell line was differentiated to neuron-like cells with RA alone before proceeding with the treatment at day 6.

Hydrogen peroxide (H₂O₂), a byproduct of oxidative stress derives from the enzymatic or spontaneous dismutation of superoxide anions, which are the byproducts of a wide and ubiquitous variety of oxidases [18]. H₂O₂ has been known for a long time to be toxic for numerous cell types [19, 20], [21]. Because of its high membrane permeability [22, 23] intracellularly formed H₂O₂ could induce its toxic effects not only within its cell of origin, but also in neighboring cells. Several findings indicate that neurons are particularly vulnerable to this mild oxidizing agent [24 – 27]. H₂O₂ directly stimulate endonuclease activity in renal tubular epithelial cells, which lead to DNA fragmentation and cell death [28]. In this study, we explored the effect of H₂O₂ on SH-SY5Y, the possible neurotoxicity and potential neuroprotective effects of PI and PPV by determining cell viability as measured by the MTS reduction assay.

MTS reduction assay has been used to reflect the number of viable cell [29]. Figure 2 shows the percentage of viable neuron-like cells after exposure to H₂O₂ as measured by the MTS assay. The result showed that at concentrations higher than 1 μM, H₂O₂ was toxic as it decreased the number of cell viability. At 100 μM, 50 % cell viability was observed. It was the IC₅₀ for the SH-SY5Y cell line and the concentration was used in determining the neuroprotective effects of PI and PPV. Previous study by

Musalma et al. [30] showed that H_2O_2 was cytotoxic at concentrations more than $100 \mu M$ as it decreased the cell viability as well as it increased the number of cell loss.




Figure 2: Effects of H_2O_2 Towards Viability of Neuron-like Cells as Assessed by the MTS. The Cells Line were First Differentiated and Cultured for Approximately 6 Days. The Cells were then Exposed to Increasing Concentrations of H_2O_2 for 24 Hour at $37^\circ C$. No Significant Difference in Viable Cell Number between Control and H_2O_2 was Observed when the Concentration of H_2O_2 was Less than $1 \mu M$. Cell Viability Decreased by 50% when Exposed to $100 \mu M$ of H_2O_2 . The IC_{50} of H_2O_2 for SH-SY5Y Cell Line was Found to be $100 \mu M$. Data is Presented as Means \pm SD (n = 3).

PI and PPV Toxicity

Figure 3 shows the neurotoxicity of PI on SH-SY5Y, as assessed by the MTS assays. Treatment with PI for 24 hour did not result in significant cell loss for a concentration range from 1 ng/ml to 10 mg/ml . The result shows that no significant difference in cell viability observed compare to control. A study by Kontush [31] showed that, Amyloid- β peptide, an antioxidant, has been reported to become a pro-oxidant at high concentration. The present data showed that high concentration ($100 \mu g/ml$) of PPV (MBu and MEH), but not PI, is toxic to the cells. The result (Figure 4) shows no significant difference in cell viability observed compared to control for

the concentration range from 1 ng/ml to 1 µg/ml for either MBu-PPV or MEH-PPV. There were more than 50 % reductions in viable cell number at concentration 100 µg/ml of both MBu-PPV and MEH-PPV. On the other hand, incubation with PI did not result in significant reduction in the number of viable cells up to 10 mg/ml concentration. However, at 10 µg/ml of MBu-PPV and MEH-PPV, the cell viability was more than 100 %. Further studies need to be done in order to understand why there is a huge reduction from concentration of 10 µg/ml to 100 µg/ml of MBu-PPV and MEH-PPV.




Figure 3: Neurotoxicity of PI on SH-SY5Y. SH-SY5Y Cell Line was Incubated with Increasing Concentrations of PI for 24 Hour at 37 °C. Prior PI Addition, Slight Difference in Cell Viability was Observed for a Concentration Range from 1 ng/ml to 10 mg/ml. The Results Show that no Significant Difference in Cell Viability Observed Compared to Control. *Denotes $P < 0.05$ Compared to Controls. Data are Presented as Mean \pm SD, from 3 Independent Experiments of Triplicate Wells (n = 3)




Figure 4: Neurotoxicity of PPV on SH-SY5Y. SH-SY5Y Cell Line were Incubated with Increasing Concentrations of PPV for 24 h at 37 °C.

The Result Shows that no Significant Difference in Cell Viability Observed Compared to Control for the Concentration Range from 1 ng/ml to 1 µg/ml for either MBu-PPV or MEH-PPV. A Slight Increase in Cell Viability was Shown at the Concentration of 10 µM for both MBu-PPV and MEH-PPV. However, there was a Huge Reduction in Cell Viability at the Highest Concentration, 100 µg/ml. *Denotes $P < 0.05$ Compared to Controls. Data are Presented as mean \pm SD, from 3 Independent Experiments of Triplicate Wells ($n = 3$)

Neuroprotective Effects of PI and PPV on H_2O_2 -Induced Cytotoxicity

Figure 5 and 6 show the neuroprotective effects of PI and PPV against H_2O_2 -induced cell loss in SH-SY5Y cell line, respectively. Cell viability was determined using the MTS reduction assay. The results show that both compounds, PI and PPV, were incapable to protect against cell loss and did not further up the cell death at concentration range from 1 ng/ml to 10 mg/ml and 100 µg/ml, respectively. Previous study reported that a compound can be considered in having a potential neuroprotective effects if the results lay in between positive and negative results [32]. As seen on the data, all concentrations did not have a potential neuroprotective effect towards neuron-like cells. However, both PI and PPV also do not cause further cell death after H_2O_2 treatment. This indicated that PI and PPV had

no ability to protect neuronal cells against H_2O_2 -induced cell loss or either further up the cell death.

Earlier study by Kang *et al.*, [33] used SH-SY5Y human neuroblastoma cells to test the neuroprotective effects of four structurally related flavones: wogonin, chrysin, apigenin, and luteolin against oxidative stress-induced cell death. Luteolin and apigenin significantly inhibited cell death induced by hydrogen peroxide, resulting in an increase in the cell. Chrysin at a concentration of 50 μM was able to inhibit cell death but by far lesser extent than that of luteolin and apigenin. Wogonin, however, had no effect on hydrogen peroxide-induced cell death.

As reported earlier, the high biodegradability by gastrointestinal, plasma and tissue peptidases are among the factors of low bioavailability of peptides. Various chemical strategies have been developed to try and overcome the limitations of peptides to increase their in vivo plasma residence time [34]. With further studies, PI and PPV could be the candidates to carry peptides to overcome the limitation of peptides to increase their in vivo plasma residence time or as prodrugs for improved CNS delivery.




Figure 5: Neuroprotective Effects of PI against H_2O_2 -induced Cell Loss in SH-SY5Y Cell Line. Cells were Pretreated with Different Concentrations of PI for 1 Hour before Exposure to 100 μM H_2O_2 for 24 hour at 37 °C. Cell Viability was Determined using the MTS Reduction Assay. From the Result, it shows that this Compound was Incapable to Protect against Cell Lost or Further Killing the Cells at a Concentration Range from 1 ng/ml to 10 mg/ml. *Denotes $P < 0.05$ Compared to Control. Data is Presented as means \pm SD, $n = 3$




Figure 6: Neuroprotective Effects of PPV against H₂O₂-induced Cell Loss in SH-SY5Y Cell Line. Cells were Pretreated with Different Concentrations of PPV (either MBu or MEH) for 1 Hour before Exposure to 100 µM H₂O₂ for 24 hour at 37 °C. Cell Viability was Determined using the MTS Reduction Assay.

The Result shows that MBu-PPV and MEH-PPV were Incapable to Protect against Cell Lost or Further Up the Cell Death at Concentration Range from 1 ng/ml to 100 µg/ml. *Denotes P < 0.05 Compared to Control.

Data is Presented as means ±SD, n = 3

Conclusion

The present findings showed that both PI and PPV are not toxic to the SH-SY5Y cells, and could not protect the cells against H₂O₂-induced cell injury as well as did not further up the cell death after H₂O₂ treatment. In addition, PI and PPV are safe to be used, as they are not toxic to the cells up to concentration of 10 mg/ml and 10 µg/ml, respectively. More studies need to be done such as apoptosis and histochemistry in order to understand the possible neurotoxicity of PI and PPV to be the candidates in the drug delivery system.

Acknowledgement

This research was funded by FRGS Phase 1/2007-Top-Down and Universiti Teknologi MARA (UiTM), MALAYSIA.

References

- [1] Balasubramaniam, A., 2002. 'Clinical potentials of neuropeptide Y family of hormones', *Am. J. Surg.*, 183, 430 – 434.
- [2] Binaschi, A., Bregola, G., Simonato, M., 2003. 'On the role of somatostatin in seizure control: clues from the hippocampus'. *Rev. Neurosci.*, 14, 285 – 301.
- [3] Strohle, A., and Holsboer, F., 2003. 'Stress responsive neurohormones in depression and anxiety', *Pharmacopsychiatry*, 36, 207 – 214.
- [4] Claes, S.J., 2004. 'Corticotropin-releasing hormone (CRH) in psychiatry: from stress to psychopathology', *Ann. Med.*, 36, 50 – 61.
- [5] Datar, P., Srivastava, S., Coutinho, E., Govil, G., 2004. 'Substance P: structure, function, and therapeutics', *Curr. Top. Med. Chem.* 4, 75 – 103.
- [6] Gentilucci, L., 2004. 'New trends in the development of opioid peptide analogues as advanced remedies for pain relief', *Curr. Top. Med. Chem.*, 4, 19 – 38.
- [7] Brasnjevic, I., Steinbusch, H. W. M., Schmitz, C. and Martinez, P. M., 2009. 'Delivery of peptide and protein drugs over the blood–brain barrier', *Progress in Neurobiology* 87, 212 – 251.
- [8] Han, X., Liu, J., Liu, M., Xie, C., Zhan, C., Gu, B., Liu, Y., Feng, L. and Lu, W., 2009. '9-NC-loaded folate-conjugated polymer micelles as tumor targeted drug delivery system: Preparation and evaluation in vitro', *Pharmaceutics* 372, 125 – 131.

- [9] Fontana, G., Licciardi, M., Mansueto, S., Schillaci, D. and Giammona, G., 2001. 'Amoxicillin-loaded polyethylcyanoacrylate nanoparticles: Influence of PEG coating on the particle size, drug release rate and phagocytic uptake', *Biomaterials* 22, 2857 – 2865.
- [10] Duncan, R., Gac-Breton, S., Keane, R., Musila, R., Sat, Y.N., Satchi R. and Searle, F., 2001. 'GacPolymer–drug conjugates, PDEPT and PELT: basic principles for design and transfer from the laboratory to clinic', *Controlled Release*, 74, 135 – 146.
- [11] Yoshikawa, T., Sakaeda, T., Sugawara, T., Hirano, K. and Stella, V. J., 1999. *Advanced Drugs Delivery Reviews*, 36, 255 – 275.
- [12] Kreuter, J., 2001. 'Nanoparticulate systems for brain delivery of drugs', *Advanced Drugs Delivery Reviews*, 47, 65 – 81.
- [13] Cheung, Y, Lau, W. K, Yu, M., Lai, C. S., Yeung, S., So, K. and Chang, R. C., 2009. Effects of all-trans-retinoic acid on human SH-SY5Y neuroblastoma as *in vitro* model in neurotoxicity research. *Neurotoxicology*, 30, 127 – 35.
- [14] Jämsä, A., Hasslund, K., Cowburn, R. F., Bäckström, A. and Vasänge, M., 2004. 'The retinoic acid and brain-derived neurotrophic factor differentiated SH-SY5Y cell line as a model for Alzheimer's disease-like tau phosphorylation', *Biochemical and Biophysical Research Communications*, 31, 993 – 1000.
- [15] Constantinescu, R., Constantinescu, A. T, Reichmann, H. and Janetzky, B., 2007. 'Neuronal differentiation and long-term culture of the human neuroblastoma line SH-SY5Y', *J Neural Trans Suppl*, 72, 17 – 28.
- [16] Presgraves, S. P, Ahmed, T., Borwege, S. and Joyce, J. N., 2004. 'Terminally differentiated SH-SY5Y cells provide a model system for studying neuroprotective effects of dopamine agonists', *Neurotox Res*, 5, 579 – 98.

- [17] Pählman, S., Hoehner, J. C., Nånberg, E., Hedborg, F., Fagerström, S. and Gestblom, C., 1995. 'Differentiation and survival influences of growth factors in human neuroblastoma', *Eur J Cancer*, 31, 453 – 8.
- [18] Cross, A. R. and Jones, O. T. G., 1991. 'Enzymic mechanisms of superoxide production', *Bioenergetics*, 1057, 281 – 298.
- [19] Byler, R. M., Sherman, N. A., Wallner, J. S. and Horwitz, L. D., (1994). 'Hydrogen peroxide cytotoxicity in cultured cardiac myocytes is iron dependent', *Am. J. Physiol.* 266, 121 – 127.
- [20] Cantoni, O., Guidarelli, A., Sestili, P., Mannello, F., Gazzanelli, G. and Cattabeni, F., 1994. 'Hydrogen peroxide cytotoxicity under conditions of normal or reduced catalase activity in IandO, -sensitive and -resistant Chinese hamster ovarv (CHO) cell variants', *Toxicol Lett*, 73, 193 – 199.
- [21] Clapp, P. A., Davies, M. J., French, M. S. and Gilbert, B.C., 1994. 'The bactericidal action of peroxides; an E.P.R. spin-trapping study', *Free Radical Research*, 21, 147 – 167.
- [22] Halliwell, B., 1992. 'Reactive oxygen species and the central nervous svstern. *J Neurochem*, 59,1609 – 1623.
- [23] Makino, N., Mochizuki, Y., Bannai, S. and Sugita, Y., 1994. Kinetic Studies on the Removal of Extracellular Hydrogen Peroxide by Cultured Fibroblasts', *Biological Chemistry*, 269, 1020 – 1026.
- [24] Théry, C., Chamak, B. and Mallat, M., 1991. 'Cytotoxic effect of brain macrophages on developing neurons', *Eur. J. Neurosci*, 3, 1155 – 1164.
- [25] Buckman, T. D., Sutphin, M. S. and Mitrovic, B., 1993. 'Oxidative stress in a clonal cell line of neuronal origin: effects of antioxidant enzyme modulation', *J. Neurochem*, 60, 2046 – 2058.
- [26] Behl, C., Davis, J. B., Lesley, R. and Schubert, D., 1994. 'Hydrogen peroxide mediates amyloid β protein toxicity', *Cell*, 77, 817 – 827.

- [27] Whittemore, E. R., Loo, D. T. and Cotman, C. W., 1994. 'Exposure to hydrogen peroxide induces cell death via apoptosis in cultured rat cortical neurons', *NeuroReport*, 5, 1485 – 1488.
- [28] Ueda, N. and Shah, S. Y., 1992. 'Endonuclease-induced DNA damage and cell death in oxidant injury to renal tubular epithelial cells', *J. Clin. Invest.*, 90, 2594 – 2597.
- [29] Wong, J. K., Kennedy, P. R. and Belcher, S. M., 2001. 'Simplified serum- and steroid-free culture conditions for high-throughput viability analysis of primary cultures of cerebellar granule neurons', *J. Neurosci. Methods*, 110, 45 – 55.
- [30] Musalmah, M., Mian, T.S., Gapor, M.T. and Wan, Z.W.N., 2006. 'Comparative effects of α -tocopherol and γ -tocotrienol against hydrogen peroxide induced apoptosis on primary-cultured astrocytes', *J. Neurosci. Sci.*, 243, 5 – 12.
- [31] Kontush, A., 2001. 'Amyloid- β : an antioxidant that becomes a pro-oxidant and critically contributes to Alzheimer's disease', *Free Radical Biology and Medicine*, 31, 1120 – 1131.
- [32] Mishima, K., Tanaka, T., Pua, F., Egashira, N., Iwasakia, K., Hidaka, R., Matsunaga, K., Takata, J., Karube, Y. and Fujiwara, M., 2003. 'Vitamin E isoforms α -tocotrienol and γ -tocopherol prevent cerebral infarction in mice', *Neuroscience Letters* 337, 56 – 60.
- [33] Kang, S. S., Lee, J. Y., Choi, Y. K., Kim, G. S. and Han, B. H., 2004. 'Neuroprotective effects of flavones on hydrogen peroxide-induced apoptosis in SH-SY5Y neuroblastoma cells', *Bio-organic and Medicinal Chemistry Letters*, 14, 2261 – 2264.
- [34] Vlieghe, P., Lisowski, V., Martinez, J. and Khrestchatisky, M., 2010. 'Synthetic therapeutic peptides: science and market', *Drug Discovery Today*, 15, 40 – 56.