MORPHOLOGICAL CHARACTERISTIC OF LUDAI(Sapium baccatum)

NUR FARAHIN BINTI SALLEH

This Final Year Project Report Submitted in Partial Fulfillment of the Requirements for the Degree of Bachelor of Science (Hons.) Furniture Technology in the Faculty of Applied Sciences
Universiti Teknologi MARA
Pahang

JULY 2016

CANDIDATE'S DECLARATION

I declare that the work in this thesis was carries out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the result of my own work, unless otherwise indicated or acknowledge as referenced work. This topic has not been submitter to any academic institution or non-academic institution for any other degree of qualification.

In the event that my thesis be found to violate the conditions mentioned above, I voluntary waive the right of conferment of my degree and agree to be subject to the disciplinary rules and regulation of Universiti Teknologi MARA.

Signature of Candidate:

Name of Candidate:

Nur Farahin Binti Salleh

Cabdidate's ID No.:

2014200092

Programme:

Furniture Technology

Faculty:

Faculty Applied Sciences

Thesis Title:

Morphological Characteristic of

Ludai

(Sapium

baccatum)

Date:

2016

ABSTRACT

MORPHOLOGICAL CHARACTERISTIC OF LUDAI (Sapium baccatum)

The scientific name of Ludai is *Sapium baccatum*. This species has the potential to be used for wood composite product in the furniture industry and wood plastic composite product. However, information about these *Sapium baccatum* is still inadequate especially those related to the morphological characteristic. In this study, morphological characteristic are carried out because of its importance to determine its fiber length, fiber diameter, lumen diameter and cell wall thickness towards different height portion and distance (from near bark to near pith). Runkel ratio and felting power are also calculated to determine the suitability of fiber in terms of producing a quality paper. The highest average recorded for fiber length, fiber diameter, lumen diameter and cell thicknesses are 1.217mm, 44.88µm, 31.33µm and 13.56µm respectively. While the highest runkel ratio and felting power recorded for *Sapium baccatum* are more than 1 and less than 60 percent. The result shows that this species is suitable for paper making.

TABLE OF CONTENTS

		Pag
ACKNOWL TABLE OF LIST OF TA LIST OF PI LIST OF AI	ONS E'S DECLARATION EDGEMENT CONTENTS ABLES GURES LATES BREVIATIONS PPENDICES	ii iii iv v vi viii ix x x xi xii xiii
CHAPTER		
1	INTRODUCTION	
	1.1 Background of Study1.2 Problems Statement1.3 Scope and Limitation of Study1.4 Objectives of Study	1 2 2 2
2	LITERATURE REVIEWS	
	 2.1 Hardwood 2.2 Ludai 2.2.1 Properties 2.2.2 Uses 2.3 Anatomical Properties 2.3.1 Heartwood and Sapwood 2.3.2 Wood Fiber 2.3.3 Fiber Morphology 2.3.4 Runkel Ratio 2.3.5 Felting Power 2.4 Effect of Fiber Density on Wood 	3 4 4 4 5 5 8 8 9 10 11
3	MATERIALS AND METHODS	
	3.1 Material Preparations3.2 Experimental of Design3.3 Methods3.3.1 Wood Density	12 14 15 15

	3.3.2 Fiber Maceration	15
	3.3.3 Fiber Morphology	19
4	RESULTS AND DISCUSSIONS	
	4.1 Introduction	22
	4.2 Density	23
	4.3 Fiber Morphology	24
	4.3.1 Statistical Analysis of Variance	25
	4.3.2 Effect of Portion on Morphological Characteristic	26
	4.3.3 Effect of Distance on Morphological Characteristic	28
	4.3.4 Runkel Ratio and Felting Power	30
5	CONCLUSIONS AND RECOMMENDATIONS	
	5.1 Conclusions	31
	5.2 Recommendations	32
REFERE	NCES	33
APPEND	ICES	35
CURRICI	JLUM VITAE	41
	TION OF THE PROJECT REPORT UNDERTAKING	45
	SION FOR REFERENCES AND PHOTOCOPYING	46