

**ABUNDANCE OF ANTS ( Hymenoptera : Formicidae ) IN UITM  
KUALA PILAH, NEGERI SEMBILAN**

**NURSHUADA BINTI YUNUS**

**Final Year Project Submitted in  
Partial Fulfillment of the Requirement for the  
Degree of Bachelor of Science (Hons.) Biology  
In The Faculty of Applied Sciences  
Universiti Teknologi MARA**

**JULY 2018**

This Final Year Project Report entitled “**Abundance of Ants (Hymenoptera : Formicidae) in UiTM Kuala Pilah, Negeri Sembilan**” was submitted by NurShuada Binti Yunus, in partial fulfillment of the requirements for the Degree of Bachelor of Science (Hons.) Biology, in the Faculty of Applied Sciences, and was approved by

---

Nursyazni Binti Abdul Rahim  
Supervisor  
Faculty of Applied Sciences  
Universiti Teknologi MARA (UiTM)  
Negeri Sembilan, Kampus Kuala Pilah,  
Pekan Parit Tinggi, 72000 Kuala Pilah  
Negeri Sembilan

---

Ahmad Zaimi Bin Mohd Zawawi  
Co-Supervisor  
Faculty of Applied Sciences  
Universiti Teknologi MARA (UiTM)  
Negeri Sembilan, Kampus Kuala Pilah,  
Pekan Parit Tinggi, 72000 Kuala Pilah  
Negeri Sembilan

---

Lili Syahani Binti Rusli  
Coordinator FSG661 AS201  
Faculty of Applied Sciences  
Universiti Teknologi MARA (UiTM)  
Negeri Sembilan, Kampus Kuala Pilah,  
Pekan Parit Tinggi, 72000 Kuala Pilah  
Negeri Sembilan

---

Dr. Aslizah Binti Mohd Aris  
Head of Biology School  
Faculty of Applied Sciences  
Universiti Teknologi MARA (UiTM)  
Negeri Sembilan, Kampus Kuala Pilah,  
Pekan Parit Tinggi, 72000 Kuala Pilah  
Negeri Sembilan

Date: \_\_\_\_\_

## TABLE OF CONTENTS

	<b>PAGE</b>
<b>ACKNOWLEDGEMENTS</b>	<b>iii</b>
<b>TABLE OF CONTENTS</b>	<b>iv</b>
<b>LIST OF TABLES</b>	<b>vi</b>
<b>LIST OF FIGURES</b>	<b>vii</b>
<b>LIST OF ABBREVIATIONS</b>	<b>viii</b>
<b>ABSTRACT</b>	<b>ix</b>
<b>ABSTRAK</b>	<b>x</b>
<b>CHAPTER 1: INTRODUCTION</b>	
1.1 Background of Study	1
1.2 Problem Statement	2
1.3 Significance of Study	3
1.4 Objectives of Study	4
<b>CHAPTER 2: LITERATURE REVIEW</b>	
2.1 Ant Subfamilies	5
2.1.1 Dolichoderinae	5
2.1.2 Dorylinae	6
2.1.3 Formicinae	7
2.1.4 Ectatomminae	7
2.1.5 Leptanillinae	8
2.1.6 Myrmicinae	9
2.1.7 Proceratiinae	9
2.1.8 Pseudomyrmecinae	10
2.2 Morphology of Ants	11
2.3 Habitat of Ants	12
2.4 Roles of Ants in Ecology	13
2.5 Ants as Social Insects	14
2.6 List of Ants that have Venom in the World	14
2.7 Damage the Venom may cause	15
<b>CHAPTER 3: METHODOLY</b>	
3.1 Materials	17
3.1.1 Raw Materials	17
3.1.2 Chemicals	17
3.1.3 Apparatus	17

3.2	Methods	17
3.2.1	Ant Sampling Method	18
3.2.2	Ant Cleaning and Sorting	18
3.2.3	Subfamily Identification	18
3.2.4	Wet Preservation	19
3.2.5	Dry Preservation	19
3.2.6	Labelling	19
3.3	Data Analysis	20
3.3.1	Shannon-Wiener Diversity Index	20
3.3.2	Shannon-Wiener Evenness Index	20
3.3.3	Margalef's Richness Index	21
<b>CHAPTER 4: RESULTS AND DISCUSSION</b>		
4.1	Frequency of Formicidae	22
4.1.1	Study Site A	23
4.1.2	Study Site B	24
4.1.3	Study Site c	25
4.2	Diversity of Ants at Three Different Study Sites	27
4.3	Evenness of Ants at Three Different Study Sites	28
4.4	Richness of Ants at Three Different Study Sites	29
<b>CHAPTER 5: CONCLUSIONS AND RECOMMENDATIONS</b>		<b>30</b>
<b>CITED REFERENCES</b>		<b>32</b>
<b>APPENDICES</b>		<b>36</b>
<b>CURRICULUM VITAE</b>		<b>37</b>

## **ABSTRACT**

### **ABUNDANCE OF ANTS (Hymenoptera: Formicidae) IN UiTM KUALA PILAH, NEGERI SEMBILAN**

Ants are dominant group of insect that can be found anywhere except for polar region. There are few factors that contribute toward habitation of ants at an area such as soil, temperature and resources. Ants have 22 subfamilies and each of them have their own habitat preferences and feeding behavior. This study aims to determine the abundance of ants in UiTM Kuala Pilah. Method used in this study is bait-fall trap. It is recommended compared to manually captured ants because it is accurate. This is because ants have different timing of foraging. By using bait-fall trap, we are able to captured ants with different foraging behavior because the trap is left for 24 hours. From the result, there are three subfamilies exist in UiTM Kuala Pilah. Each area has different resources availability. This contributes to the abundance of ants at an area. Ants choose habitat by considering the amount of disturbance occur in an area. One of the areas has low abundance due to high human interaction in the area. The ants migrate its nesting to other area that offers resources and have less disturbance in order to have high chances of survival.