

**ECOLOGICAL ANALYSIS OF BEETLES POPULATION IN  
UNIVERSITI TEKNOLOGI MARA NEGERI SEMBILAN**

**ADDIBA NUR SYAZWANNIE BTE ABD WAHID (2011296038)**

**AIN SYAFIQAH BT SA'AT (2011821254)**

**ALEEZA HANEE BT ALYKALLAM (2011435614)**

**AMY ZAWIAH BINTI ABDULLA (2011806754)**

**UNGKU AMIRUL ARIF BIN UNGKU ABDULLAH (2011284352)**

**Mini Project Report Submitted in  
Partial Fulfillment of the Requirements for the  
Diploma in Science  
in the Faculty of Applied Sciences  
Universiti Teknologi MARA**

**SEPTEMBER 2013**

## TABLE OF CONTENT

	<b>Page</b>
<b>DECLARATION</b>	iii
<b>APPROVAL SHEET</b>	iv
<b>ACKNOWLEDGEMENTS</b>	v
<b>TABLE OF CONTENTS</b>	vi
<b>LIST OF TABLES</b>	viii
<b>LIST OF FIGURES</b>	ix
<b>LIST OF ABBREVIATIONS</b>	x
<b>ABSTRACT</b>	xi
<b>ABSTRAK</b>	xii
<b>CHAPTER 1 INTRODUCTION</b>	
1.1 Background	1
Significance of study	
Objective of study	
<b>CHAPTER 2 LITERATURE REVIEW</b>	
2.1 Importance of Beetles	4
2.2 Habitat of Beetles	6
2.3 Diversity of Beetles	7
<b>CHAPTER 3 MATERIALS AND METHODS</b>	
3.1 Capturing of Beetles	9

3.2 Killing of Beetles	10
3.3 Pinning of Beetles	11
3.4 Preserving the Beetles	12
3.5 Identification of Beetles	13
List of Equipment	14
List of Chemicals/Media	14
<b>CHAPTER 4 RESULTS AND DISCUSSION</b>	
4.1 Taxonomy Composition	15
4.2 Diversity Indices	19
<b>CHAPTER 5 CONCLUSION</b>	22
<b>REFERENCES</b>	23
<b>APPENDICES</b>	26
<b><i>CURRICULUM VITAE</i></b>	27

## LIST OF FIGURES

<b>Figure 3.1</b> Capturing beetles using plastic cups	9
<b>Figure 3.2</b> Beetles captured in cups	9
<b>Figure 3.3</b> Dropping ethyl ethanoate onto the beetles	10
<b>Figure 3.4</b> Pinning Process	11
<b>Figure 3.5</b> Drying oven (40°C to 45°C)	12
<b>Figure 3.6</b> Beetles captured to be identified	13
<b>Figure 4.1</b> Beetles population in UiTM Negeri Sembilan	26

## ABSTRACT

### ECOLOGICAL ANALYSIS OF BEETLES POPULATION IN UNIVERSITI TEKNOLOGI MARA NEGERI SEMBILAN

Beetles are under the order of Coleopteran. It can be found whether on land or in the water. In Malaysia, there are variety species of beetles can be found and it is believed by scientists that some of the species are yet to be discovered, especially in the unexplored and virgin tropical rain forest in Sarawak and Sabah. Beetles can also easily be found in UiTM Negeri Sembilan especially at night. The main objective of this study is to determine the taxonomy composition of beetles in UiTM Negeri Sembilan. The study is conducted around the area of UiTM Negeri Sembilan. The beetles found within this area are collected and is kept to be preserve. In UiTM Negeri Sembilan. It found that *Chrysina restlendens* species has the most number of beetles captured. Otherwise, the least species captured are *Halyomorpha halys* and *Heterocerus marginatus*. There are three different of diversity indices uses in this study which is Shannon-Weiner index, Richness index and Evenness index. Shannon-Weiner index shows a low diversity of 1.83, Richness index shows a low richness of 2.1 and the Evenness index shows of high evenness which is 0.83.