# ANATOMICAL PROPERTIES OF MENGKIRAI (TREMA ORIENTALIS)

FENAINAH AMAT MUKSIN

DIPLOMA IN WOOD INDUSTRIES UNIVERSITI TEKNOLOGI MARA

2000

#### ACKNOWLEDGEMENT

I would like to state my grateful appreciation to Dr. Suhaimi Muhammed as my final project advisor for their leadership, encouragement and critical comment.

I also wish to express great thanks to Prof. Madya Abdul Jalil Ahmad as Course Tutor, Dr. Jamaluddin Kasim as Lecturer of WTE 375 and Mr.Rudaini Mohd. Nawawi as laboratory assistant for their valuable help during the preparation for this final project.

Finally, special thanks to all of my beloved friends, classmates also to those who are involved either directly or indirectly for their help and spiritual during the whole period of this project.

## **TABLES OF CONTENTS**

.

## Page

DEDICATION	iii
ACKNOWLEDGEMENTS	iv
LISTS OF FIGURES	vii
LISTS OF PLATES	viii
LISTS OF ABBREVIATIONS	ix
ABSTRACT	x
ABSTRAK	xi

## CHAPTER

1.	INTRODUCTION	1
	1.1 Overview	1
	1.2 Objectives	5
2.	LITERATURE REVIEW	6
	2.1 Anatomical features of hardwoods	6
	2.1.1 Vessels	8
	2.1.2 Wood Parenchyma	9
	2.1.3 Rays	10
	2.1.4 Fibers	11
	2.1.5 Tyloses	11
	2.1.6 Included phloem	12
	2.2 Others features	12
	2.3 Heartwood and Sapwood	13
	2.4 Source of Raw Materials	13
3.	MATERIALS AND METHODS	14
	3.1 Source of Raw Material	14
	3.2 Preparation of Materials	14
	3.3 Sectioning	18

	3.4 Staining and Mounting	21
	3.5 Scanning Microscope of Wood Specimens	22
4.	RESULTS AND DISCUSSIONS	24
	4.1 Transverse Section	24
	4.2 Tangential Section	29
	4.3 Radial section	31
5.	CONCLUSION	39
		41
REFEI	RENCES	
VITA.		42

÷,

#### ABSTRACT

## ANATOMICAL PROPERTIES OF MENGKIRAI (Trema orientalis)

By

### JUNAINAH AMAT MUKSIN

April 2000

Anatomical properties of Mengkirai Trema Orientalis were ascertained based on parts or tree such as base, middle and top which base was describe as diameter of breast height, middle was 50% from bark and top was upper 50%. The sources of Mengkirai woods were from Reserve Forest of UiTM Pahang, which named as medium tree and growth at edge the forest. Anatomical properties that had in this wood were it has diffuse porous and three types vessel likes solitary, pairs, triple and a few tail's vessels. In solitary vessels had a tylosis. Wood rays were uniseriate types and homocellular. It was a place of food reserve for this tree. This wood has apotrachea and paratrachea Paratrachea parenchyma was incomplete vecisentric. parenchyma. Beside that apotrachea parenchyma was not arranged. Cell walls were thin effected from wide lumen cells. From this information, the conclusion is Mengkirai wood is easy to preserve because it has wide lumen cell and thin cell walls. Both of these properties can make absorption of preservation materials in the wood. Thus Mengkirai wood is still not commercial yet.