COMPARISION OF EXTRACTIVE CONTENT BETWEEN SAPWOOD AND HEARTWOOD FOR MAHANG GAJAH (Macaranga gigantae)

by

Che Azahar bin Adzemin

Final Project Paper Submitted in Partial Fulfillment for the Diploma in Wood Industry,

Faculty of Applied Sciences,

Universiti Teknologi MARA Pahang Branch

November 2005

ACKNOWLEDGEMENT

Alhamdullilah.....

Firstly, I would like to thank Al- Mighty Allah for granting me strength and courage to complete this project.

I wish to express my gratitude to Prof. Dr. Suhaimi bin Muhammed, my project advisor, who helped and advised me to complete this final project. I also wish to acknowledge my sincere thanks to our lectures directly and indirectly involved during this project in progress. Last but not least, I express my thanks also to our Wood Industry Workshop and laboratory staffs En. Sardey, En. Anuar and En. Rudaini for their efforts and technical help.

Acknowledgements are also due to my friends of my own course, my own part Jen, Hakim, Azhan, Rizduan, Fadhil, Nizam, Naim and other who are give their support and cooperation for me to finish my final project.

"Thanks a lots"

V

TABLE OF CONTENTS

Pages

PROJECT TITLE	ii
APPROVAL SHEET	iii
DEDICATION	iv
ACKNOWLEDGEMENT	v

LIST OF ABBREVATIONS	ix
ABSTRACT	X
ABSTRAK	xi

CHAPTER I

1.0	INTRODUCTION AND OBJECTIVES			
	2			

CHAPTER II

2.0	LITERATURE REVIEW		3
	2.1	Field characteristic of Mahang Gajah	3
	2.2	Definition of Sapwood and Heartwood	4
	2.3 Extractives in wood2.4 Terpenoids and steroids		7
			8
		2.4.1 Occurrence	8

	2.4.2	Chemica	l composition	9
		2.4.2.1	Monoterpenoids	10
		2.4.2.2	Sesquiterpenoids	10
		2.4.2.3	Diterpenoids	11
		2.4.2.4	Triterpenoids and steroids	11
		2.4.2.5	Polyterpenoids	12
2.5	Fats a	and Waxes	S	12
2.6	Phene	olic Const	ituents	13

CHAPTER III

3.0	MATERIALS AND METHODS			
	3.1	Source of material	14	
	3.2	Chemical Analysis	15	
		3.2.1 Moisture Content	15	
		3.2.2 Hot Water Soluble in Sapwood and Heartwood	16	
		3.2.3 Alcohol-Benzene Soluble in Sapwood and Heartwood	17	
		3.2.4 1% NaOH Soluble in Sapwood and Heartwood	18	

ABSTRACT

COMPARISON OF EXTRACTIVE CONTENT BETWEEN SAPWOOD AND HEARTWOOD FOR MAHANG GAJAH (Macaranga gigantae)

By

Che Azahar bin Adzemin

November 2005

The amount of extractives may vary greatly between species, between trees and within a tree. The wood just inside the inner bark of a tree (sapwood) will normally contain few or no extractives while the extractive content of the inner part of a tree (heartwood) can be very high. For this final project, the Mahang Gajah species (*Macaranga gigantae*) is choice to find out the amount of the extractives content in sapwood and heartwood portion with find the values of moisture content, hot water soluble, Alcohol-benzene soluble and 1% NaOH soluble.

Wood extractives such as hot-water solubles, 1% NaOH solubles and alcohol-benzene solubles was studied for the variation according to sapwood and heartwood. The finding indicated that Mahang gajah process extractives of higher percentage at the heartwood as compared to the sapwood region.