

UNIVERSITI TEKNOLOGI MARA

**IN-VITRO ANTI-INFLAMMATORY ACTIVITY OF
STANDARDIZED AQUEOUS EXTRACT OF *ERYTHROXYLUM
CUNEATUM* IN LIPOPOLYSACCHARIDE-INDUCED RAW 264.7
CELLS**

SITI KHUZAIMAH BINTI MAAROF

2010859614

BACHELOR OF PHARMACY

2013

ACKNOWLEDGEMENTS

In the name of Allah, the Compassionate and the Merciful. Alhamdulillah, with His will and guidance, this research project is finally completed. The success and final outcome of this project required a lot of guidance and assistance from many people and I am extremely fortunate to have them along completion of my project work.

My sincere gratitude, deepest appreciation and respect to my research supervisor Dr. Mizaton Hazizul Hasan for providing continuous support and guidance throughout the period of this research. I am very grateful that she had spared me her time for me to discuss about my progress, shared her knowledge and giving me advice for improvement.

I owe my profound gratitude to our project guide Siti Rohani Saleh, who took keen interest on our project work and guided us all along by providing all the needful information and had spared some of her time for us until the completion of our project. In addition, thanks to all people from the Pharmacotoxicology Laboratory for invaluable assistant and encouragement.

I am also heartily thankful to all my family members for their continuous support. Their prayer, motivational comfort and assistance in which has given me strength to overcome all the difficulties throughout the project work. Last but not least, thanks to all my fellow members especially to my housemate for their support and guidance and of course to my team work, NurHafizah bt Abd Mutalib.

TABLE OF CONTENTS

TITLE PAGE	
APPROVAL FORM	
ACKNOWLEDGEMENT	ii
TABLE OF CONTENTS	iii
LIST OF FIGURES	v
LIST OF ABBREVIATIONS	vi
ABSTRACT	viii
CHAPTER ONE: INTRODUCTION	
1.1 Background Of Study	1
1.2 Objective	6
1.2.1 Specific Objectives	6
1.3 Hypothesis	6
CHAPTER TWO: LITERATURE REVIEW	
2.1 Inflammation	7
2.2 Inflammation Process	9
2.2.1 Phase 1	9
2.2.2 Phase 2	10
2.3 Others Mediators of Inflammation	12
2.4 Chronic Inflammation	16
2.5 Anti Inflammatory Drugs	18
2.6 <i>Erythroxylum Cuneatum</i>	18

ABSTRACT

Erythroxylum cuneatum has long been used by the folks as a traditional herbal medicine to treat inflammatory disease such as rheumatoid arthritis by chewing the dried leaves. This plant was chosen because only few studies were conducted on *Erythroxylum cuneatum*. The aim of the study is to determine the in-vitro anti-inflammatory activity of standardized aqueous extract of *Erythroxylum cuneatum* in lipopolysaccharide-induced RAW 264.7 cells. We evaluate the protective and toxicity effect of *Erythroxylum cuneatum* by measuring the percent cell death in MTT assay. The pro-inflammatory cytokines such as Tumor Necrosis Factor alpha (TNF- α) and Interleukin-1 β (IL-1 β) production were evaluated after the cells were induced by the lipopolysaccharide (LPS) of gram negative bacteria with presence of *Erythroxylum cuneatum*. The pro-inflammatory cytokines were evaluated by ELISA. In this research, upon stimulation of *Erythroxylum cuneatum* in lipopolysaccharide-induced RAW 264.7 cell showed that there are significantly reduced of the amount of pro-inflammatory cytokines production. These results suggest that *Erythroxylum cuneatum* may have anti-inflammatory activity by inhibiting TNF- α and IL-1 β in RAW 264.7 cell.

CHAPTER ONE

INTRODUCTION

1.1 Background of study

Erythroxylum cuneatum is a plant of interest in this research. This plant is from the Erythroxylaceae family. The common names of this plant are *Chinta mula*, *inai-inai*, *kayu urang*, wild cocaine and *Gu Ko Yi* (Ody, 2000). Leaves from the Erythroxylaceae family such as *Erythroxylum coca* are used as an anti-inflammatory agent by the old folks. The leaves contain several alkaloid, the most important alkaloid are cocaine. The used of this leaves is as stimulant, carminative and also has a potent local anesthetic effect when applied to mucosal membranes (Jarald & Jarald, 2006). So, it is believed that *Erythroxylum cuneatum* will have the same indication for the treatment of inflammation.

The inflammatory response or inflammation is when tissues are damaged by inflammagen (Rainsford, 2004). Inflammagen is an irritant that elicits both edema and the cellular response of inflammation. Inflammagens can be bacteria, trauma, toxins, heat or any other cause. Inflammation produces five major symptoms that are redness, heat, swelling, pain and disturbance of function (Stoelting & Hillier, 2012).