By

MOHD HARIS ISMAIL

Final Project Paper Submitted in Partial Fulfillment for the

Diploma in Wood Industry, Faculty of Applied Sciences,

Universiti Teknologi MARA

March 2005

ACKNOWLEDGEMENT

Firstly I would like to take opportunity to shows my thankfulness towards ALLAH S.W.T for rendering upon me such tremendous with determination and courage the period of time leading to complete this project.

Besides that, I pleased to express my great appreciation to my project advisor Prof. Dr. Suhaimi Bin Muhammed, due to the courage on the advices, comments and suggestion which had expanded the valuable during the preparation of this project.

I would also like to express my appreciation to all lecturers and staff I known in this Wood Technology Industry, for their helping during my study in this UiTM Pahang.

Finally, special thank to all my friend also to those who are involved either directly or indirectly for their help and moral support during the whole period of this project.

Thank you very much.....

TABLE OF CONTENTS

	Page
APPROVAL SHEET	. i
DEDICATION	. ii
ACKNOWLEDGEMENTS	. iii
LIST OF FIGURE	. vi
LIST OF PLATE	. vii
LIST OF TABLE AND GRAPH.	viii
ABSTRACT	ix
ABSTRAK	x
CHAPTER	
1.0 INTRODUCTION	
1.1 General	. 1
1.2 Justification	. 1
1.3 Problem Statement	3
1.4 Guidelines for the average gardener	4
1.4.1 A viable Alternative	6
1.4.2 Economic Potential	7
1.4.3 Environmental Performance	8
1.5 Objective	9

2.0 LITERATURE REVIEW

2.1 Characteristic: Seeds, Leaves, Stalks, Flowers and seed pods	10
2.2 Uses for Kenaf	11
2.3 Anatomical Properties.	12
3.0 MATERIAL AND METHODS	
3.1 Preparation of Materials	13
3.2 Sectioning.	15
3.3 Staining and Mounting	16
3.4 Scanning Microscope of Frond Specimens	17
4.0 RESULTS AND DISCUSSION	
4.1 Compare Stem and Branch	18
5.0 CONCLUSION.	21
REFERENCES	22
VITA	23

DETERMINATION OF PORE DENSITY IN KENAF (Hibiscus cannabinus)

By

MOHD HARIS ISMAIL

MARCH 2005

ABSTRACT

Determination of pore density in Kenaf's tree to find number of pore based on part of cross section. The raw materials were from Jengka 8, Pahang. The cell had in these tree were it has a same cell structure but different in number of pores. This information can help for provide in study for the future because no data available. It can be concluded, there are differences at number of pores which in cross sectional. Stem portion of Kenaf has more pores than the branch portion of Kenaf.