

UNIVERSITI TEKNOLOGI MARA

**MODELING SEASONAL WIND
PATTERN FROM IMAGING AND
NON-IMAGING DATA**

SITI NORATIQA BINTI MOHAMAD DEROS

Thesis submitted in fulfillment
of the requirements for the degree of
Doctor of Philosophy

Faculty of Applied Science

December 2016

ABSTRACT

Wind speed in Malaysia is categorized as low with annual mean of 3-5 m/s and influenced primarily by four monsoon season; Northeast and Southwest monsoon with two transition period; April and October inter monsoon. Monsoon seasons were classified based on the origin of wind blows that brings unique character of wind speed and direction. Current wind study utilized limited wind data to projects wind behavior in wind pattern model generation. Short-term wind data insufficient to explain wind criteria by neglects the seasonality behavior of wind caused by different monsoon season. Wind observation using ground measurement devices such as anemometer and wind vane produced limited spatial resolution data that initiated the used of imagery data with 100-150 km² swath width. Wind direction extraction using wavelet transform (WT) technique is an example of wind study using imagery data. However, the study focused on high wind speed area due to the limitations of wavelet coefficient derivation that requires minimum 7 m/s wind speed and underestimates streak produced by lower speed. This study attempts to develop wind pattern model by forecasting the seasonal wind speed, determine the seasonal wind speed distribution model and extract wind direction from SAR images acquired at low wind speed area. The study site represents the seasonal and low wind speed condition in Pulau Langkawi, Malaysia. The timely wind speed data used for year 2000-2010. The imagery data used is Standard-2 (100 km²) and Wide-3 (150 km²) RADARSAT-1 SAR images to picture high spatial resolution wind direction in Langkawi. Autoregressive and Moving Average (ARMA) model was used to combine the seasonal and non-seasonal component of wind speed that will be used in wind speed forecast. The seasonal wind speed distribution model was determined among Lognormal, Weibull and Gamma distribution model; evaluated using goodness-of-fit test with the lowest error value is the most fit distribution model. Introduction of new WT technique to extract low wind speed wind direction was performed on Standard-2 and Wide-3 RADARSAT-1 SAR images. Enhancement involved the derivation of wavelet coefficient of low wind-induced streak on SAR data that then was transformed by using Fast Fourier Transform (FFT), Short Time Fourier Transform (STFT) and Mexican-Hat wavelet transform technique. Finally, the wind pattern model used is the Multi-Layer Conveyor (MLC) model simulated from wind speed, direction and the day in monsoon as the time unit, and the wind speed distribution as wind capacity measure. SARIMA (1,1,1)×(1,1,3) model is the ARMA model that best represents the seasonal wind speed with high r-squared value (>0.9) in each monsoon season. This showed that the coefficient of determination between sample data and forecasted data is relatively high. Lognormal distribution model is the best model used to describe the seasonal distribution of wind with goodness-of-fit test showed lowest error (0.07-0.14) between the model performance and data. The introduction of new wavelet transform technique using FFT domain spectrum is the most suitable technique to extract wind direction at low wind speed area with r-squared value of 0.71 for Standard-2 and 0.99 for Wide-3 image. Wind pattern model developed showed high r² (0.94) and correlation (0.97) with actual wind data. As a conclusion, the seasonal wind pattern model developed using MLC model by using derived parameters successfully described the wind behavior of distinct monsoon season and able to projects the long-term scenario of low wind.

ACKNOWLEDGEMENTS

All praises and thanks to Allah s.w.t., the supreme ruler of the universe for the spiritual and gift bestowed upon me in the performance of my routines. No words can describe how much I should be thankful to each of everyone who involved whether directly or indirectly during my PhD completion. My hearty gratitude expression goes to Associate Professor Dr. Arnis Asmat (PhD.), the Senior Lecturer in Faculty of Applied Science, Universiti Teknologi MARA, Shah Alam, Malaysia. This wonderful lady had been guide me closely, helped me endlessly and always understand me with her kind cooperation, friendly supervision in this work. I never forget to thank my co-supervisor, Prof. Hj. Dr. Shattri bin Mansor, Faculty of Engineering, Universiti Putra Malaysia for his valuable suggestions, guidance and encouragements.

I extend this appreciation to my family and my parents especially. Their support is everything for me during this thesis completion. I also appoint my appreciation to my research team, my colleagues whom I exchange the sweet and hard moments, ideas and knowledge. My journey will be so hard without them.

Special thanks to myBrain15 program in Scholarship Department of Malaysian Minister of Science, Technology and Innovation (MOSTI), for the doctoral sponsorship.

My appreciation goes to the following organizations for the data provision and guidance.

Malaysian Remote Sensing Agency (MRSA)
Malaysian Meteorological Department (MMD)
Universiti Teknologi MARA (UiTM)

TABLE OF CONTENTS

	Page
CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR’S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENTS	v
TABLE OF CONTENTS	vi
LIST OF TABLES	ixi
LIST OF FIGURES	xii
LIST OF PLATES	xviii
LIST OF SYMBOLS	xix
LIST OF ABBREVIATIONS	xixi
CHAPTER ONE: INTRODUCTION	
1.1 Research Context	1
1.1.2 Importance of Wind Observation	3
1.1.3 Wind Observation in Malaysia and Wind-related Studies	6
1.2 Research Problem and Justification	7
1.3 Research Aims	8
1.4 Significant of Study	11
1.5 Novelty	12
1.6 Thesis Structure	14
CHAPTER TWO: LITERATURE REVIEW	15
2.1 Winds in Malaysia	15
2.2 Statistical Wind Study and Observation	17
2.2.1 Wind Frequency Distribution Study	18
2.2.2 Wind Seasonality Study	21
2.2.3 Limitations of Statistical Wind Study	23
2.3 Wind Observation Using Radar System	24
2.3.1 Wind Study Using Remote Sensing Radar System	27
2.3.2 Radar Contribution in Wind Analysis	28

CHAPTER ONE

INTRODUCTION

1.1 RESEARCH CONTEXT

Wind is one of important climatic factor that may affect the daily routine especially in coastal area. Wind was created due to the rise of temperature that increases the pressure in one regional area. The high pressure moved to another area that has less pressure and this movement creates wind. Monsoonal wind is the phenomena where the various character of wind was created periodically and creates unique pattern.

Monsoon season is a climate that has obvious seasonal shift of prevailing winds between winter and summer. This climate involved certain region includes Asia, Australia, Africa & Indian Ocean that experience clear fluctuation between dry winter and rainy summer. The term ‘monsoon’ derives by the word ‘mausim’ that can be defined as season. This monsoon season is mainly due to the seasonal shift of prevailing wind direction that occurs due to the difference in temperature of land and sea across the Tropics. Over half of the world’s population, includes most developing countries, they live under the climates that was dominated by monsoon season (Chabangborn, 2014). Meteorologically, monsoon season refers to the directional shift of wind from one season to another season. Monsoon occur when humid wind blows from an ocean towards the land in summer and cooler wind blows towards an ocean off the land in winter. This cause 180° changes in wind direction (Weller and Cai, 2014).

Monsoon seasonality is due to the rotation of the Earth, while the monsoon weather systems are a result of the land-sea temperature differences due to the effect of solar radiation. When the Earth rotates and revolves around the sun, different seasons occur due to the different land-masses of the northern and southern hemispheres. This phenomena is mainly due to the land surface at northern hemisphere is larger than the southern hemisphere. Therefore, northern hemisphere temperature rise higher and results in the opposing seasons between the northern and southern hemisphere. This seasonality is directly influence the wind behavior. During