

UNIVERSITI TEKNOLOGI MARA

**TLC-BIOAUTOGRAPHIC METHOD FOR
DETECTION OF α -AMYLASE INHIBITION IN
MARINE ALGAE**

NURUL NILLIE AMANIN BINTI WAN ZIN

**Dissertation submitted in partial fulfillment of the requirements for
the degree of Bachelor of Pharmacy (Hons.)**

Faculty of Pharmacy

JUNE 2017

ABSTRACT

Bioautographic method using TLC plays an important role in search of plant bioactive compound. In this study, TLC-bioautographic method was conducted to investigate antidiabetic activity in marine algae. Marine algae are abundant and known to have bioactive compound with health benefit including as antioxidant and antidiabetic effects. The objective of this study was to investigate the activity of α -amylase inhibition and antioxidant in marine algae using TLC-bioautographic method. Marine algae samples were extracted in two different solvents, ethanol and ethyl acetate. After migration of the samples, the first plate was dipped in α -amylase solution and incubated. Then it was dipped in starch solution and incubated again. The reaction was visualized by washing the plate with Gram's iodine. Another developed plate was dipped in DPPH• solution. The ethyl acetate extracts of marine algae showed higher α -amylase inhibition and antioxidant activity. Sample 8 showed the highest inhibition with moderate antioxidant activity. Antioxidant activity may contribute to α -amylase inhibition activity. Comparison of results may show relationship but must be further investigated. Marine algae showed the potential candidate for management of type 2 diabetes.

ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious, the Most Merciful.

Alhamdulillah, praise be to Allah for giving me the strength and health for me to be able to complete this research project in given time.

I would like to express my deepest appreciation to my supervisor, Prof Snezana Aganotovic-Kustrin for all her guidance, support and kind supervision.

I would like to forward my gratitude to my teammates, Syahira, Alia, and Lutfi for being very helpful and supportive friends. In addition, I would like to thank to the technicians from the chemistry lab UiTM Puncak Alam , especially Mrs Farhanah and Mr Rahimi for always helping us in providing all that we need during laboratory session.

To my family members, thank you for all the love and motivation. I apologize for not being able to spend more time with all of you during the completion of this research. Last but not least, I want to thank all my friends and everyone that had indirectly helped me.

Thank you.

TABLE OF CONTENTS

ABSTRACT	ii
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATIONS	viii
 CHAPTER ONE	 1
INTRODUCTION	1
1.1 Introduction	1
1.2 Problem Statement	3
1.3 Objectives	4
1.4 Significant of Study	4
1.5 Hypothesis	4
 CHAPTER TWO	 5
LITERATURE REVIEW	5
2.1 Marine Algae	5
2.2 Antioxidant	7
2.3 Diabetes Mellitus	9
2.4 TLC-Bioautography	10

CHAPTER ONE

INTRODUCTION

1.1 Introduction

In the recent years, researchers have gained more interest on natural resources for the source of antioxidant in view of toxic and carcinogenic effect of synthetic antioxidants. Many phytochemicals from terrestrial and marine plants have been evaluated for the past two decades. Marine algae have gained many interests as many studies have proven to contain a bioactive compound of the phenolic compound with antioxidant. The phenolic-rich extract of marine algae also found to have anti-diabetic properties through the inhibition of α -amylase and α -glucosidase (Nwosu et al., 2011).

Marine algae can be classified according to its pigmentation; red (Rhodophyta), brown (Phaeophyta), and green (Chlorophyta). Previous studies have proven that algae rich in bioactive particularly due to antioxidant that have wide potential in the industry especially in food, pharmaceuticals and biomedical industry (S. H. Lee et al., 2010). Therefore, the number of studies on algae increases in the recent years due to this interest on the antioxidant compound.