

Abnormal Behaviors in Captive Wildlife: To Keep or Not to Keep?

Nurfarah Ain Limin¹, Siti Nurfatiha Najihah Fakhrol Hatta¹, Nurul Adilla Zaki¹
and Nur Nadiyah Md Yusof^{1,2}

¹*School of Biology, Faculty of Applied Science, Universiti Teknologi MARA, Selangor, Malaysia*

²*Institute of Biodiversity and Sustainable Development, Universiti Teknologi MARA, Shah Alam, Selangor, Malaysia*

ARTICLE INFO**Article history:**

Received 22 November 2024

Revised 25 December 2024

Accepted 30 December 2024

Published 30 January 2025

Keywords:

Captive

Captive wildlife

Stereotypic behavior

Stress behavior

Wildlife conservation

DOI: 10.24191/scl.v19i1.6356

ABSTRACT

Wildlife in captivity serves several purposes, including research, conservation, agriculture, and tourism. Many zoos and wildlife parks are driven towards conservation roles as they help prevent endangered wildlife from becoming extinct. However, a captive environment is often not the best manifestation of the animal's natural habitat, thus hindering them from performing natural behavior as they would in the wild. This imposes stress on the captive animals, leading to the display of abnormal behaviors, such as stereotypic behaviors, which are repetitive, invariant, and functionless behaviors. Stereotypic behaviors have been observed in many captive animals, such as pacing in tigers and bears, swaying and bobbing in elephants, over-grooming, self-mutilating, coprophilia, and coprophagia among captive primates, as well as fur and/or feather plucking in primates and birds. This article explores the abnormal behaviors of captive animals in response to their environment and highlights the critical importance of enrichment and naturalistic habitat design. Creating environments encouraging species-specific behaviors can significantly improve animal welfare, enhance conservation outcomes, and educate the public about wildlife conservation. Improved welfare practices not only support animal well-being but also strengthen public engagement and advocacy for conservation initiatives, ultimately aiding in protecting endangered species.

* Corresponding author. *E-mail address:* nurnadiyah@uitm.edu.my

INTRODUCTION

The suboptimal welfare of captive animals poses serious concerns that warrant effective interventions to improve their well-being. Beyond ethical considerations, this issue is pivotal in understanding the evolutionary development of animal behaviors [1] and its implications for conservation

efforts. Observing unnatural behaviors exhibited by animals in zoos and wildlife parks often evoke a sense of unease, as signs of lethargy, inactivity, and stereotypic behaviors, such as pacing, swaying, or object-licking, are indicative of compromised welfare [2-5]. These behaviors are more than symptoms of psychological distress; they also reflect an animal's inability to adapt to specific environmental stimuli [6], posing significant challenges to conservation outcomes and raising important questions about captive wildlife's welfare and psychological well-being. Zoos and conservation centers are primarily established to conserve endangered species by captive breeding [7] and enhance research on animal behavior and ecology while also educating the public about the critical roles of wildlife in sustaining ecosystems [8-10], ultimately contributing to biodiversity preservation efforts [11].

However, when animals experience chronic stress in captivity [1], it can lead to a cascade of adverse effects, including impaired reproductive success, increased susceptibility to diseases, display of stereotypic behaviors [12], increased cortisol levels [10], all of which threaten long-term species survival [13]. A list of evidence points out the impaired well-being of captive wild animals, which indicates their needs are not being satisfied [14]. For example, herbivores, such as giraffes, are short-lived in zoos compared to when they are conserved in their natural habitat due to the lack of nutritional value and low energy in their food supply, indicating insufficiency in their diets [15]. Moreover, many giraffes in captivity displayed stereotypic behaviors such as object-licking [16, 17]. Similarly, stereotypic behaviors are widespread among other herbivores, including captive elephants [18-20] and carnivores such as tigers and leopards [10, 21, 22]. Addressing these welfare concerns of captive wildlife is not only critical for improving the quality of life for individual captive animals. However, it is also instrumental in achieving the overarching goals of conservation programs. Understanding the underlying factors and consequences of abnormal behaviors can inform the development of evidence-based strategies to mitigate stress, enhance reproductive outcomes, and support species' survival. Furthermore, captive wildlife displaying maladaptive behaviors may face challenges upon reintroduction to the wild. This article explores the types of abnormal behaviors displayed by captive wildlife, their relation to environmental stressors in captivity, and the measures required to minimize the occurrence of these behaviors and improve conservation outcomes.

Animal behaviours in captive wildlife

Natural behavior is displayed in environments that are pleasant for the animals and support their biological functioning [23]. Examples of natural behaviors common in wildlife are foraging and exploring habitat, as observed in many species, including sun bears [24, 25]. Wild orangutans also invest almost half of their time foraging and feeding [26] and prefer to stay in high spots throughout the day and night [27]. Conversely, Macaques are often observed to perform locomotor, investigative, and social activities in the wild [28]. Another example is free-ranging tigers that typically live in solitary and often patrol their wide home ranges or territories [29-33]. They are the apex predators in the wild and are great hunters. They normally hunt at night and occasionally during the day [30].

In captive environments, however, natural behaviors are commonly substituted with abnormal behaviors, including stereotypic behaviors such as pacing [34]. Abnormal behaviors can be defined as atypical responses that are not typically observed in the wild, displayed by animals exposed to specific stimuli and conditions they are unable to cope with [14], which are frequently interpreted as a form of coping mechanism that organisms utilize in response to stressors [35]. Captivity has been shown to induce anomalies in behavioral patterns across various species [36], including wildlife [37, 38]. This scenario poses significant concerns, particularly for threatened species, as it could adversely affect reproductive success, physiological health, and overall life span [39-41]. Behavior is a practical and widely utilized stress indicator in zoological research, as it can be systematically measured. This allows for an accurate assessment of animals' responses to environmental alterations, thereby providing insights into their well-being about environmental changes [42]. The inability of animals in zoos to exhibit normal behavior is often a result of environmental factors such as unfavorable weather conditions, unpleasant substrates,

unnatural lighting, exposure to loud noises, and disturbing odors [40]. Barren settings, spatial limitations, restricted locomotion, lack of retreat area, inescapable proximity with humans, restricted contact with other species, lack of feeding variety and chances, social housing, and other limitations in behaviors are some of the stressors present in captivity [4, 43]. The effects of stressors depend significantly on how an individual perceives and responds to the stress agents [44].

Warwick et al. (2013) [45] stated that animals living in a confined setting or artificial ecosystem encounter numerous stressors that deviate from those typically experienced in their natural habitats. In most cases, animals housed in confinement with limited space have a high probability of developing stereotypic behaviors [4, 46], as the limited availability of space hinders the potential for environmental enrichment [47]. The inadequate space offered by the enclosures restricts animals from eliciting their normal species-specific behaviors, such as foraging and exploring their environment [4, 24, 25]. In addition, smaller enclosures are also associated with aggressiveness [48] and stereotypical pacing in tigers [10, 32, 49, 50]. In addition to limited space within enclosures, a lack of environmental stimulation can develop stereotypic behaviors [47]. In barren enclosures, animals typically respond to the lack of stimulation by either (1) decreasing their exploratory behaviors or (2) seeking to fulfill their exploratory needs through stereotypic behaviors [4], as can be seen in bears [24, 40].

Furthermore, enclosure type, high visitor density, and humidity also contribute to the development of abnormal behaviors. A study conducted by Arumugam et al. (2018) [51] highlighted the significant influence of enclosure type, humidity, and visitor density on the behavior of Malayan tapirs (*Tapirus indicus*) in captivity. Their findings indicate that tapirs housed in semi-natural environments exhibit a higher feeding frequency than those housed in artificial environments.

Introduction to stereotypic behaviors

Numerous studies across various animal species indicate that reproductive success and survival rates are higher in captivity than in their wild counterparts [52, 53]. However, when wildlife is placed in environments that significantly deviate from their natural habitats, they become increasingly vulnerable to developing stereotypic behaviors [54]. For instance, inadequate housing conditions, such as the absence of perches and hiding places for captive birds, primarily contribute to the manifestation of stereotypic behaviors [55]. Stereotypic behaviors are conclusively described as behaviors that are repetitious, invariant, and without clear functions, such as pacing, head bobbing, fur/feather plucking, and over-grooming [54, 56, 57], and are often indicative of psychological distress [58]. These behaviors are typically associated with stress experienced by animals in captivity [59] and are thought to arise from various stresses caused by boredom, physical constraint, fright, agitation, and frustration [50, 60-62]. Such frustration typically stems from the animals' inability to resolve persistent problems in captivity or perform natural species-specific behaviors [54]. For example, captive European starlings often exhibit stereotypic somersaulting [63], attributed to many repetitive, failed attempts to escape [64].

In this context, stereotypic behaviors are considered psychological behaviors synonymous with human psychological conditions such as anxiety, depression, and obsessive-compulsive disorder (OCD) [65]. In the early stages, such behaviors may serve as coping mechanisms to help animals adapt to environmental challenges. However, lacking opportunities to engage in diverse natural behaviors can lead to significant neurobiological changes, potentially impairing brain function and altering neural connections, physiology, and anatomy. Importantly, these neuroanatomical are often considered as a consequence rather than the root cause of the behaviors [66]. Furthermore, the manifestation of stereotypic behaviors has been extensively investigated concerning several factors, including personality traits [67], individual variability [68], sex-specific susceptibility [69], and genetic dispositions [70].

Stereotypic behaviors are widespread across taxa in captivity [3, 37, 71]. Primates were observed to display behaviors that are equivalent to certain depressive patterns observed in humans [72, 73]. Birkett and Newton-Fisher (2011) [36] reported that all 40 chimpanzees living in six accredited zoos displayed stereotypic behaviors, suggesting mental dysfunction akin to mental illness in humans. In addition to prosimians, giant pandas [37, 71, 74] and red pandas [75] in captivity were also observed to exhibit stereotypic behaviors that illustrate stressful conditions or poor well-being. Research also highlights differences in behaviors based on housing conditions. For example, fully captive omnivorous sun bears displayed significantly higher levels of stereotypic behaviors than those housed in semi-captivity (i.e., they were allowed to the forest during the day but housed in enclosures at night), likely due to multiple factors such as enclosure size and environmental complexity [76]. Captive wildlife exhibits diverse stereotypic behaviors, reflecting their responses to environmental stressors. The following sections explore common forms of stereotypies observed in captivity, including pacing, head tossing/bobbing/body swaying, excessive grooming, coprophagia, fur and/or feather plucking, self-mutilating, object gnawing and/or licking, genital stimulation, urophagia, investigative behaviors, and fearful behaviors, passive behaviors and aggression.

Types of stereotypic behaviors

Pacing

Pacing or circling is one of the most common stereotypic behaviors in captive wildlife. It is characterized by repetitive walking along the same path with no obvious objective [3, 32, 77] and is often an indicator of stress [78]. For example, horses often exhibit stereotypic circling within their stalls, even during feeding, showing a lack of interest in their surroundings [4]. In tigers, pacing is typically identified as walking in a figure-eight pattern with more than two rotations [56]. Among chimpanzees, the pacing is described as a quadrupedal locomotion along a repeated route in a conformed manner with no obvious goals [36]. The prevalence of pacing behavior has been widely documented across taxa. Carnivorous animals in captivity, such as tigers, spend much time pacing back and forth within their enclosure [10, 21, 22, 28, 56, 79]. Similarly, stereotypic pacing has been reported in various wildlife species, including leopards, bears, giraffes, and even sloths. Environmental stimuli, such as enclosure design, feeding routines, or human activities, often trigger such behavior. For instance, Mallapur and Chellam (2002) [80] reported that stereotypic behaviors in captive leopards increased in correspondence to the zookeepers' activities and during longer feeding periods. It was observed that the leopards paced significantly more when the zookeepers were cleaning the off-exhibit enclosures. Similar behaviors were observed during feeding time, in which they were seen to be agitated and restless. The leopards were also found to significantly increase their stereotypical pacing during high numbers of visitors, especially during festive seasons. Visitors can be stimulating for the animals, but constant exposure to the high density of visitors can be traumatizing [54].

Environmental stressors, such as limited control over the environment [81], restricted space, social isolation [16], unmet intrinsic needs [14], and inability to perform species-specific behaviors [82], exacerbate stereotypic pacing. For territorial species such as tigers, who naturally patrol their extensive wild home ranges [50, 83, 84], spatial constraints in captivity prevent them from engaging in such territorial patrolling [32, 50]. Furthermore, while wild tigers are skilled hunters, the fixed feeding schedules typical of captive settings limit opportunities for foraging behaviors, such as hunting prey. This restriction, coupled with anticipatory behavior around feeding times, often contributes to the development of pacing behavior [50]. Similar patterns have been observed in other carnivores, such as leopards [10, 85] and lions [86]. In addition, herbivores, such as giraffes, exhibit stereotypic behaviors in the form of repetitive pacing [16, 17], potentially due to spatial constraints. Other herbivores, including captive sloths, are also not excluded from displaying pacing [87].

Tailored strategies, including enrichment and enclosure modifications, have shown promise in reducing stereotypic behaviors. Pitsko [47] demonstrated that tigers kept in enclosures designed to mimic their natural habitat with the provision of natural substrates, vegetation, and shaded areas exhibited reduced pacing behavior and increased exploration. Interestingly, there was no incidence of stereotypic behaviors by two tigers that were kept in bigger enclosures throughout the observation. Natural substrates such as grass or hay beddings, leaves and grass, and wood chips are recommended to encourage typical species-specific behaviors in animals in captivity [88]. The captive tigers invested 90% of their time in well-shaded areas, indicating the necessity of having large, shaded areas, particularly on sunny days. For captive felids, artificial substrates that do not reflect naturally occurring substrates in the environment, such as concrete, not only increase stereotypic pacing [89] but can also cause physical injuries to the legs and throbbing footpads [88]. Such findings extend to other species. For example, a study comparing fully captive sun bears with semi-captive sun bears found that the fully captive bears spent significantly more time pacing [76]. The stereotypical pacing exhibited by the captive bears aligns with findings that associate pacing with animal stress, particularly in bears [38, 90]. The bears also paced significantly in the evening in anticipation of food, as it was routinely provided to them during that period [76]. Recognizing such taxa-specific differences is critical for developing strategies to improve welfare, such as enrichment that meets species-specific needs [91].

Head tossing, bobbing, and body swaying

Tossing or bobbing of the head is a circular or up-and-down motion performed without clear functions in a repetitive manner, classifying it as a type of stereotypic behavior [28, 92, 93]. This behavior is particularly prevalent in primates. For instance, zoo chimpanzees have reportedly exhibited repetitive head bobbing [36], as have captive-born macaques, frequently repeatedly moving their heads up and down [28]. Beyond primates, head bobbing has also been documented in elephants [35]. On the other hand, Swaying involves repetitive side-to-side movements of the head and body, typically performed at least three consecutive times [5, 46, 94] or by transferring weight rhythmically from front to rear and back again [46, 95]. Swaying is common among elephants [5, 19, 20] and has also been observed in cynomolgus monkeys [28]. Weaving, a related behavior characterized by head-swinging combined with shifting weight from one foot to the other without forward movement, is frequently observed in horses [4]. Bettinger et al. (1997) [96] suggested that head-swaying in the largest mammals, such as elephants, is a coping mechanism to soothe themselves. Maternal deprivation has been identified as a potential contributing factor to the development of these stereotypic behaviors. For example, captive elephant calves separated from their mothers after weaning are more likely to exhibit head bobbing or swaying, highlighting the role of early social and environmental conditions in the emergence of these behaviors [35]. This highlights the critical role of appropriate maternal care and enriched environments in mitigating the development of stereotypes in captivity.

Excessive grooming

Grooming or preening in some taxa, such as birds, fulfills both physiological and social functions in various species and is instrumental for survival and adaptation [97, 98] and ectoparasite removal [99]. However, in the presence of environmental stressors, grooming behavior is often displaced, resulting in the development of pathological excessive grooming [100-102]. Over-grooming or excessive grooming is characterized by repetitive grooming focused on specific body parts, apparently without any clear objective or intention. This behavior may also involve using objects such as sticks or other tools to lightly drag across body surfaces in a non-focused manner [36]. Over-grooming is considered a stress-coping behavior [103] and has been documented across taxa, particularly primates. For instance, captive chimpanzees often exhibit stereotypic over-grooming [36], while captive macaques lick their tails repetitively and various body parts, suggesting analogous behavioral patterns in these primates [28]. These behaviors may arise from the inability to engage in species-typical social grooming, such as allo-grooming among conspecifics [104].

Carnivores, including captive jaguars, have also been observed to display over-grooming behavior [105]. Potential underlying causes of over-grooming include chronic itching, leading the animals to over-groom, or even self-mutilating to alleviate discomfort [106]. Over-grooming may also result in hair loss, impairing thermoregulation and further exacerbating the animals' physiological stress [107].

Coprophagy

Coprophagy, the deliberate act of eating feces, involves animals ingesting their feces or that of another animal [108]. This behavior is often attributed to stress, boredom, and compromised welfare [109]. It has been observed across various taxa, including horses, pigs, koalas, and birds [110]. Among primates, chimpanzees have been documented engaging in coprophagy, often picking through their excretions and consuming the remnants of food and seeds found in their feces [111]. A study examining chimpanzees across six zoos identified coprophagy as one of the most prevalent stereotypic behaviors observed [36], a finding supported by additional research [108, 112]. Interestingly, infant chimpanzees were found to mimic this behavior after observing their mothers, suggesting that coprophagy can be a socially learned behavior despite its classification as a stereotype [108, 109, 112]. A similar pattern has been observed in koalas, where their infants consume maternal feces to acquire microbial strains necessary for digesting eucalyptus leaves [113]. Coprophagy is also reported in avian species, particularly among precocial birds such as quails, turkeys, and ptarmigans [110, 114].

Fur or feather plucking

Fur or feather plucking, depending on the species' concern, is a stereotypical grooming behavior in which an animal self-inflicts itself by pulling out its fur, hair, or feathers for no obvious reason. Fur-plucking is a behavioral abnormality observed in various species, including felids [115] and primates such as the great apes [116]. Captive chimpanzees, for example, have been reported to pluck not only their body hair but also the hair of their conspecifics [36]. Among carnivores, fur-plucking has been documented in captive tigers, cougars, and lions, often resulting in fur loss and skin irritation [117, 118]. Feather-plucking is particularly prevalent among captive birds, such as parrots [100] and cockatoos, and is frequently associated with limited opportunities for naturalistic behaviors, including foraging, social interactions, and physical exercise [55].

Hand-rearing has also been implicated in developing stereotypic behaviors such as feather-plucking, as observed in hand-reared grey parrots [55]. When directed toward conspecifics, this behavior is called feather-pecking, which leads to loss of feathers, tissue injuries, and pain, signaling serious welfare concerns [119]. The resulting bald patches can attract further tissue pecking, escalating cannibalistic behavior that can be fatal [120]. Feather loss also compromises the birds' physical condition, resulting in prolonged discomfort [119]. Controlling feather-pecking is particularly challenging in large groups, as social learning promotes the spread of these behaviors, including cannibalism, throughout the flocks [121]. Feather damage caused by feather-pecking is often linked to fear and the pain of being pecked, as observed in confined hens [55].

Self-mutilating

Self-mutilating or self-harming behaviors, including self-hitting and self-biting, occur when animals intentionally inflict injury upon themselves. These actions may involve striking their bodies against surfaces using hands, paws, or limbs to strike themselves or biting their skin or appendages [36, 92]. Various factors, such as improper diet, social frustration, aggressive tendencies, and boredom, have been suggested as potential contributors to the development of these behaviors [122]. Self-mutilating has been documented across a wide range of taxa. For example, a captive king vulture exhibited severe self-inflicted injuries that became contaminated, exposing underlying bone [123]. Similarly, self-harming behaviors such as self-biting and head-banging are also prevalent in primates [124]. Japanese macaques and pig-tailed monkeys have been observed repetitively rushing into wired enclosures, leading to self-harm [122]. In a study of zoo-living chimpanzees, Birkett and Newton-Fisher (2011) [36] reported self-harming behavior, such as hitting themselves with their hands. Apart from that, captive sun bears have also demonstrated self-mutilating behaviors, including self-biting, hitting, and pinching repetitively without any purpose [76]. Tail-biting is another form of self-injury commonly observed in pigs [125] and is often linked to improper diet, gut discomfort, compromised health, and genetics [4]. Similarly, self-mutilation is prevalent among felids, manifesting as behaviors such as excessing, licking, biting, or scratching [126]. In equines, self-injurious behaviors include self-biting, kicking, and pushing against stall structures, often reflecting frustration or discomfort in captivity [4]. Collectively, these examples highlight that self-mutilation is not confined to a specific group of animals but is a widespread issue in captive settings, often indicative of underlying welfare concerns.

Objects gnawing or licking.

Oral stereotypic behaviors, such as bar biting, wall licking, and substrate gnawing, are commonly observed in captive animals. These repetitive actions involve using teeth to lick, gnaw, or chew on objects within their enclosures. Such behaviors have been documented across various species. For instance, in a study of 40 male macaques on a breeding farm, individuals housed in solitary conditions exhibited repetitive oral stereotypies, specifically bar gnawing and licking [28]. Similarly, captive giraffes frequently display pronounced substrate-licking behavior, including wall-licking [16, 17]. In pigs, oral stereotypies such as bar-biting and sham chewing are prevalent [127]. Bar-biting is often interpreted as an attempt to escape, whereas sham chewing is a displacement behavior that substitutes actual feeding [128]. A survey revealed that nearly 80% of giraffes and okapis displayed stereotypic behaviors, with substrate licking accounting for 72% of these observed behaviors, as stated by Lewis et al. [129]. In horses, oral stereotypies such as crib-biting, wood chewing, and wind sucking are particularly common. Crib biting involves biting into their cribs, fences, or other stall structures [4], and these stereotypies are typically associated with increased stomach acidity [4]. These behaviors emphasize the need for tailored interventions to address underlying causes, such as environmental stressors or physiological imbalances.

Genitals non-sexual stimulation

Genital stimulation refers to the repetitive act of an animal stimulating its genitalia through actions such as touching, patting, clinging, fondling, and/or rubbing it continuously, typically in a non-sexual context. These behaviors have been observed in multiple captive species. For instance, Camus et al. [28] reported that macaques kept in a breeding farm repeatedly clung to their genitalia and atypically stimulated neighboring conspecific's genitals. Similarly, captive chimpanzees have been documented to touch, pat, stroke, and fondle their genitalia repetitively in non-mating conditions [36].

Urophagia

Urophagia or urine consumption is a form of stereotypic behavior in which an animal ingests its urine or that of others, typically by licking. This abnormal behavior is prevalent in primates such as macaques, chimpanzees [130], and other species, including birds. For example, urophagia has been documented in chimpanzees across six different zoos [36].

Investigative behaviours

Investigative or attentive behaviors, including sniffing and observing surroundings, are often performed when an animal is curious about environmental stimuli. This stimuli-seeking or exploratory behavior is typical of animals to survive, especially in the wild. However, repetitively displaying it due to stressors present in captivity is considered an abnormality. For example, Arumugam et al. [51] in his study also reported that investigative behaviors displayed by the Malayan tapirs, such as sniffing, general alertness, or observing surroundings while standing in stationary, sitting, or lying down, were higher in artificial enclosures than the semi-natural ones. This is due to the possibility of the tapirs being kept in an artificial environment exposed to blaring sounds from activities such as renovation works, traffic, and zoo visitors.

Fearful behaviours

Fearful behaviors such as being vigilant, constantly alert, quivering, fleeing, or hiding from non-harmful variables often result from stressful, traumatic experiences. For instance, captive tapirs in Zoo Melaka were found to be vigilant and always watchful of their surroundings. They displayed fearful behaviors such as tensing, alert, and fleeing to hide when hearing the noises from visitors, which frequently led to them quivering [51]. These behaviors were also observed in a tiger kept at the National Wildlife Rescue Centre (NWRC). According to the zookeepers, the said tiger possessed fearful behaviors such as panting, tensing, alertness, and fleeing to hide under platforms every time it heard growls of other tigers as well as any noise from surrounding due to its experience of being contained next to sun bears, not near its conspecifics. Consequently, the poor tiger had difficulty coping with the presence of other tigers. This aligns with the findings of Mason and Latham [131] and Wickins-Drazilova [132], who suggested that stereotypes arising from specific past experiences may diverge from their original causes and can persist in later stages of an animal's life, even in circumstances where such stereotypic behaviors are less likely to manifest.

Passive behaviours

Being passive includes standing still (stationary), lying down, resting, and sleeping. For example, sun bears in fully captive environments were observed to be significantly more passive than those in semi-captive environments [76], as they spent more time resting and sleeping, mostly in the afternoon. Kamaruzaman et al. [133] also reported that giant pandas invest most of their time inactive, which is different from their counterparts in the wild due to the constant routine provision of food in zoos. This gives them plenty of time to rest and be inactive, whereas the wild giant pandas forage for food to survive. Research showed that male giant pandas preferably spend plenty of time feeding and moving while female giant pandas tend to rest more [134]. Giant pandas in Zoo Negara, Malaysia, preferred to invest their time being inactive in rocky and cool areas as it mirrors the natural habitat of giant pandas, which is rocky, lush terrain in China [133]. With many visitors, fewer locomotor behaviors were also observed in Malayan tapirs [51]. They swam, investigated less often, and became more passive as they rested more when the number of visitors peaked. They also spent much time submerged inside the pool to avoid visitors' disturbances, such as poking and pouring water onto them [51].

Aggression or agonistic behavior

Aggressive or agonistic behavior is a type of negative social interaction [135] that can trigger stress reactions [136]. Nelson [137] stated that aggressive behavior is elicited to inflict harm or distress on another animal. Limitations in mating partners, food resources, and territory contribute to the appearance of aggressive behaviors [136]. Aggressive interactions can be observed in mammals, such as in sun bears. Restriction in exhibiting their typical behaviors in captivity contributes to stressful conditions and discouragement, which cause them to become hostile, violent, and harmful to each other and, therefore, could be destructive to their welfare [76].

Another example is captive Dorcas gazelles. They demonstrated aggressive behaviors such as butting, hitting, thrusting, striking, and pushing each other with their forehead, horns, and other body parts with force, chasing, and fighting [136]. Aggressive behaviors such as pecking, standing offs in which individuals vocalize and swing their heads in the display, and chasing [138] were also observed in captive flamingos [139].

Ways to reduce abnormal behaviors and nurture natural behaviors

The ethical debate over keeping wildlife in captivity hinges on our ability to minimize the adverse implications of confinement, particularly the prevalence of stereotypic behaviors. Animal rights advocates contend that captivity inherently infringes upon animals' intrinsic values and freedom [140]. On the other hand, utilitarian conservationists justify captivity as a necessary means to safeguard species from extinction, foster public education, and facilitate research [141]. Reconciling these perspectives is essential to ensure ethical practices align with conservation goals while prioritizing animal welfare. Central to this discussion is the design of captive environments that prioritize the provision of naturalistic habitats, which encourage captive animals to behave naturally while mitigating abnormal behaviors. Environmental enrichment is a significant tool for promoting wildlife welfare by enhancing reproductive and behavioral health [142, 143]. Through the use of diverse strategies [144], such as the inclusion of natural substrates, foliage, water bodies (i.e., pools), hiding spots, climbing structures such as logs, platforms, ropes, and other environmental items, make captive conditions more favorable and conducive to promote active, exploratory and natural behaviors [145] for various species including felids and primates. For instance, Gomes et al. [146] demonstrated that increasing environmental enrichment was associated with reducing stereotypical behaviors in captive tigers, reaffirming the potential of well-managed captivity to nurture natural behaviors. One widely used approach is foraging enrichment, which involves introducing novel food items to captive animals [147] or increasing the complexity of accessing food [148]. In cheetahs, for instance, exposure to such enrichment led to a marked increase in exploratory behaviors and a notable reduction in pacing behaviors [149]. Olfactory enrichment has similarly been shown to reduce stereotypic behaviors [150-152] and to foster natural and territorial behaviors [153, 154], as evident in black-footed cats [155].

In the wild, free-ranging animals can choose their habitat based on the necessities they require, such as food sources, shelter, and protection, adapting their behaviors to suit their environments [156]. In contrast, captive animals are subjected to rigid conditions, including routine activities and fixed diets, limiting their natural responses and changing their behavioral patterns [157], leaving them with little to no control over their environment [158]. To address these limitations, zoos and institutions must optimize enclosure designs, incorporating larger, more complex spaces with sufficient features and proper management to nurture natural, species-specific behaviors such as scent-marking [159], locomotor repertoires [32, 160] and minimize the prevalence of stereotypies [32, 161]. Alternatives to traditional captivity, sanctuaries, and semi-captive environments provide more naturalistic habitats and prioritize animal welfare. Sanctuaries typically offer animals greater autonomy and access to environments that closely mimic their natural habitats, leading to reduced stress and a decline in abnormal behaviors [162]. Similarly, semi-captive environments integrate aspects of wild habitats with managed care, fostering

improved welfare outcomes and encouraging the expression of natural behaviors while maintaining opportunities for conservation and education [163]. Evaluating these approaches offers insights into more ethical and effective captive wildlife management. Enriched environments encourage captive animals to utilize more naturalistic areas [164], and studies indicate that spatial constraints contribute to displaying stereotypic behaviors. For example, Breton and Barrot [50] identified an inverse relationship between enclosure size and pacing, aligning with the findings of Clubb and Mason [60]. Animals housed in bigger, more complex enclosures exhibited reduced stereotypic behaviors compared to those housed in smaller, barren spaces [158]. Tigers with access to larger, enriched environments displayed more inherent behaviors and experienced a reduction in stereotypies [32].

In addition to larger spaces, providing manipulable and consumable items has significantly elevated natural behaviors in captive animals, particularly primates such as orangutans, where such enrichment is associated with a marked reduction in passive and stereotypic behaviors [44]. Gippoliti [27] reported that wild orangutans invested their day and night by being in aerial environments, and this offers a great window for zoological institutions to install ropes and lofty platforms to nurture the development of arboreal activities, which are synonyms with primates like orangutans [165]. Social enrichment has also proven beneficial, particularly in felids, where housing animals with conspecifics or near them has been shown to reduce stress-related behaviors [166 - 168]. Furthermore, auditory enrichment, such as the playback of recorded lion roars, has been shown to stimulate natural behaviors, including increased live roaring and playful interactions among lions [169]. Similar effects have been observed in African birds, where auditory stimuli such as natural sounds enhanced activity levels [170], further emphasizing the potential of auditory enrichment to encourage naturalistic behaviors in captive animals.

Emerging digital tools, such as computers and tablets, offer innovative cognitive enrichment, providing challenges and diverse problem-solving opportunities for captive animals, particularly primates [171-174]. Advances in digital technology have enabled researchers to develop virtual environments where animals can interact with and exert control over their environment [174], offering sustained cognitive stimulation and mitigating the risk of rapid habituation often associated with traditional enrichment methods [171, 172, 175]. These tools can be tailored to accommodate various animals based on age, sex, social rank, and cognitive abilities [174]. Environmental enrichment devices (EEDs) have similarly demonstrated significant positive effects on animal behaviors and physiological and psychological advantages, as mentioned in Donald et al. [91]. For example, EEDs have been shown to reduce stereotypic pattern swimming and increase exploratory and foraging behaviors in ill harbor seal pups [176]. Tailored EEDs have also effectively reduced stereotypical behaviors in California sea lions and northern elephant seals in rehabilitation centers [91]. While genetic selection is a potential method for reducing stereotypies, it is rarely utilized in zoos despite its common application in agricultural and breeding settings [177]. This method, which avoids breeding individuals exhibiting stereotypic behaviors [38], is less favored in captivity due to the underlying environmental factors contributing to such behaviors. Genetic engineering, although potentially effective in addressing issues such as feather pecking [119], is not widely considered, as stereotypies typically stem from environmental deprivation and frustration.

To optimize enrichment, captive management and caretakers must consider species-specific needs and individual differences, including personality traits such as shyness or boldness, which influence how animals respond to enrichment [178]. These traits may manifest as variations in the intensity or expression of specific behaviors or as distinct behavioral patterns within similar contexts, ages, or timeframes [179]. Such behavioral differences often arise from the interaction of endocrine, neural, and immune processes, collectively called “psycho-neuroimmunology,” which influence and are influenced by behavior. These processes significantly shape personality development within the boundaries of an individual’s genetic predispositions [180]. When introducing enrichment, individual preferences, background history, and needs must also be considered [181]. For instance, while some captive macaques benefit from enrichment, others exhibit no significant changes in stereotypic behaviors [181]. Social enrichment may have little to no

positive effects on species such as giraffes and okapi, which are solitary by nature [16]. Conversely, species such as horses have shown marked reductions in stereotypies when provided with more fibrous diets, special feeding devices, stalls, and social enrichment [4]. This finding underscores the importance of considering variation in species and individuals in responding to enrichment.

While enrichment strategies should address the species' general behavioral and physiological needs, tailoring these approaches to align with individual differences, including personality traits, history, and preferences, is crucial to optimize their effectiveness and mitigate stereotypic behaviors [91]. Enrichment strategies specific to carnivores such as tigers and lions, such as frozen blood cubes, food hung on climbing poles, feeding boxes, feeding poles, simulated prey, and buried meats, have been shown to foster feeding and hunting behaviors [77, 102, 117, 158, 182, 183]. These methods not only improve animal welfare but also enhance visitor experiences. Visitors prefer observing animals in naturalistic enclosures displaying various interactive and active behaviors [184], while stereotypic behaviors and a lack of natural behaviors can detract from the exhibit's appeal [185]. Transitioning enclosures to more complex and naturalistic designs benefits the well-being of animals [186, 187] and conservation efforts by providing a more enriching environment for the animals and a more engaging experience for the visitors [184]. Ultimately, addressing abnormal behaviors in captivity is not about negating the values of captivity itself but rather about improving it to align with animal welfare standards. With effective strategies, captivity can contribute positively to conservation, education, and research, ensuring the well-being of the animals while furthering broader conservation initiatives.

CONCLUSION

Wildlife in captivity plays an instrumental role in conservation, education, and public engagement. However, it inevitably raises welfare concerns, particularly regarding the development of abnormal behaviors, particularly stereotypies. These behaviors indicate poor welfare and challenge the ethical basis of keeping animals in captivity. The answer to “To keep or not to keep?” lies in the equilibrium between the benefits and challenges of captivity. Captivity can be justified if it significantly contributes to conservation, education, and research while upholding rigorous welfare standards. To ensure ethical captivity practices, zoos and institutions should adhere to clear guidelines, which include prioritizing enclosure designs that closely resemble wildlife natural habitats, implementing a variety of species-specific and individualized enrichment measures to minimize the occurrence of abnormal behaviors, and tailoring management practices to address animal needs and preferences. Captivity should not be viewed as a binary condition but as a continuum where consistent improvement in welfare practices and interventions can significantly impact. For example, bolstering conservation programs and fostering natural behaviors through innovative technologies that can elevate the quality of life for captive wildlife. Housing wildlife in captivity is not inherently detrimental when done responsibly and ethically. Institutions must actively address welfare concerns and continually advance conservation goals. By doing so, captivity can remain a viable solution for wildlife survival and public education and awareness, ensuring that captive wildlife serves as conservation ambassadors rather than confinement symbols. However, if these standards are neglected, the ethical justification for captivity diminishes. Looking ahead, further research is needed to evaluate the long-term effects of enrichment on animal welfare, particularly in reducing stereotypic behaviors and nurturing natural behaviors.

Additionally, exploring the role of genetics in susceptibility to stress could offer valuable insights into tailoring individualized management strategies for different species and individuals. These studies will refine existing practices and guide the development of more effective and ethical captivity models. A forward-thinking approach requires ongoing commitment and innovation, ensuring that captive wildlife thrives while inspiring global conservation efforts.

ACKNOWLEDGMENTS

We thank Universiti Teknologi MARA and the Ministry of Higher Education, Malaysia. This study was supported by MOHE Grant (FRGS/1/2022/WAB11/UITM/02/01), and the first author is supported by MyBrainSc Scholarship (KPT(BS).1/21-900919045460).

AUTHOR'S CONTRIBUTION

Nurfarah Ain Limin conceptualized, drafted, wrote, and revised the article. Nurfatiha Najihah Fakhrol Hatta drafted and provided technical input. Nur Adilla Zaki provided technical input and contributed to the article's revision. Nur Nadiah Md Yusof conceptualized, supervised, and reviewed the article.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

REFERENCES

- [1] Mason, G. J. (2010). Species differences in responses to captivity: stress, welfare and the comparative method. *Trends in Ecology & Evolution*, 25(12), 713–721. <https://doi.org/10.1016/j.tree.2010.08.011>
- [2] Fureix, C., & Meagher, R. K. (2015). What can inactivity (in its various forms) reveal about affective states in non-human animals? A review. *Applied Animal Behaviour Science*, 171, 8–24. <https://doi.org/10.1016/j.applanim.2015.08.036>
- [3] Poirier, C., & Bateson, M. (2017). Pacing stereotypies in laboratory rhesus macaques: Implications for animal welfare and the validity of neuroscientific findings. *Neuroscience & Biobehavioral Reviews*, 83, 508–515. <https://doi.org/10.1016/j.neubiorev.2017.09.010>
- [4] Radkowska, I., Godyn, D., and Kinga, F. (2020). Stereotypic behaviour in cattle, pigs and horses – a review. *Animal Science Papers and Reports*, 38(4), 303 – 319.
- [5] Fuktong, S., Yuttasaen, P., Punyapornwithaya, V., Brown, J. L., Thitaram, C., Luevitoonvechakij, N., & Bansiddhi, P. (2021). A survey of stereotypic behaviors in tourist camp elephants in Chiang Mai, Thailand. *Applied Animal Behaviour Science*, 243, 105456. <https://doi.org/10.1016/j.applanim.2021.105456>
- [6] Cooper, J.J., & Albentosa, M J. (2005). Behavioural adaptation in the domestic horse: potential role of apparently abnormal responses including stereotypic behaviour. *Livestock Production Science*, 92(2), 177–182. <https://doi.org/10.1016/j.livprodsci.2004.11.017>
- [7] Marinath, L., Vaz, J., Kumar, D., Thiyagesan, K., & Baskaran, N. (2019). Drivers of stereotypic behaviour and physiological stress among captive jungle cat (*Felis chaus* Schreber, 1777) in India. *Physiology & Behavior*, 210, 112651. <https://doi.org/10.1016/j.physbeh.2019.112651>
- [8] Marino, L., Bradshaw, G. And Malamud, R. (2009). The captivity Industry. *Animals and Society, Best Friends Magazine*. March/April. 25 – 27 pp.
- [9] Suárez, P., Recuerda, P., & Arias-De-Reyna, L. (2017). Behaviour and welfare: the visitor effect in captive felids. *Animal Welfare*, 26(1), 25–34. <https://doi.org/10.7120/09627286.26.1.025>
- [10] Vaz, J., Narayan, E. J., Kumar, R. D., Thenmozhi, K., Thiyagesan, K., & Baskaran, N. (2017). Prevalence and determinants of stereotypic behaviors and physiological stress among tigers and leopards in Indian zoos. *PLoS ONE*, 12(4), e0174711. <https://doi.org/10.1371/journal.pone.0174711>

- [11] Mohapatra, B.S., and Sethy, U. (2020). Behaviour of zoo-housed tigers: A case study. *Journal of Environmental Science, Toxicology and Food Technology*, 14(12), 41 – 45. <https://doi.org/10.9790/2402-1412014145>
- [12] Budzyńska, M. (2014). Stress reactivity and coping in horse adaptation to environment. *Journal of Equine Veterinary Science*, 34(8), 935–941. <https://doi.org/10.1016/j.jevs.2014.05.010>
- [13] Mason, G., Burn, C.C., Dallaire, J.A., Kroshko, J., Kinkaid, H.M., & Jeschke, J.M. (2013). Plastic animals in cages: behavioural flexibility and responses to captivity. *Animal Behaviour*, 85(5), 1113–1126. <https://doi.org/10.1016/j.anbehav.2013.02.002>
- [14] Brunberg, E. I., Rodenburg, T. B., Rydhmer, L., Kjaer, J. B., Jensen, P., & Keeling, L. J. (2016). Omnivores Going Astray: A review and new synthesis of abnormal behavior in pigs and laying hens. *Frontiers in Veterinary Science*, 3. <https://doi.org/10.3389/fvets.2016.00057>
- [15] Clauss, M., Franz-Odenaal, T., Brasch, J., Castell, J.C. and Kaiser, T. (2007). Tooth wear in captive giraffes (*Giraffa camelopardalis*): mesowear analysis classifies free-ranging specimens as browsers but captive ones as grazers. *Journal of Zoo and Wildlife Medicine*, 38(3), 433 – 445. <https://doi.org/10.1638/06-032.1>
- [16] Bashaw, M. J., Tarou, L. R., Maki, T. S., & Maple, T. L. (2001). A survey assessment of variables related to stereotypy in captive giraffe and okapi. *Applied Animal Behaviour Science*, 73(3), 235–247. [https://doi.org/10.1016/s0168-1591\(01\)00137-x](https://doi.org/10.1016/s0168-1591(01)00137-x)
- [17] Orban, D. A., Siegford, J. M., & Snider, R. J. (2016). Effects of guest feeding programs on captive giraffe behavior. *Zoo Biology*, 35(2), 157–166. <https://doi.org/10.1002/zoo.21275>
- [18] Mason, G. J., & Veasey, J. S. (2010). What do population-level welfare indices suggest about the well-being of zoo elephants? *Zoo Biology*, 29(2), 256–273. <https://doi.org/10.1002/zoo.20303>
- [19] Greco, B. J., Meehan, C. L., Hogan, J. N., Leighty, K. A., Mellen, J., Mason, G. J., & Mench, J. A. (2016). The Days and Nights of Zoo Elephants: Using Epidemiology to Better Understand Stereotypic Behavior of African Elephants (*Loxodonta africana*) and Asian Elephants (*Elephas maximus*) in North American Zoos. *PLoS ONE*, 11(7), e0144276. <https://doi.org/10.1371/journal.pone.0144276>
- [20] Greco, B. J., Meehan, C. L., Heinsius, J. L., & Mench, J. A. (2017). Why pace? The influence of social, housing, management, life history, and demographic characteristics on locomotor stereotypy in zoo elephants. *Applied Animal Behaviour Science*, 194, 104–111. <https://doi.org/10.1016/j.applanim.2017.05.003>
- [21] Miller, L. J. (2011). Visitor reaction to pacing behavior: influence on the perception of animal care and interest in supporting zoological institutions. *Zoo Biology*, 31(2), 242–248. <https://doi.org/10.1002/zoo.20411>
- [22] Sajjad, S., Farooq, U., Anwar, M., Khurshid, A., & Bukhari, S. (2011). Effect of captive environment on plasma cortisol level and behavioral pattern of Bengal tigers (*Panthera tigris tigris*). *Pakistan Veterinary Journal*, 31(3), 195–198. <https://agris.fao.org/agris-search/search.do?recordID=PK2011001213>
- [23] Bracke, M. B. M., & Hopster, H. (2006). Assessing the importance of natural behavior for animal welfare. *Journal of Agricultural and Environmental Ethics*, 19(1), 77–89. <https://doi.org/10.1007/s10806-005-4493-7>
- [24] Carlstead, K., & Shepherdson, D. (2000). Alleviating stress in zoo animals with environmental enrichment. In *CABI Publishing eBooks* (pp. 337–354). <https://doi.org/10.1079/9780851993591.0337>
- [25] McPhee, M. E., & Carlstead, K. (2010). The importance of maintaining natural behaviors in captive mammals. *Wild mammals in captivity: Principles and techniques for zoo management*, 2, 303-313.

- [26] Campbell-Smith, G., Campbell-Smith, M., Singleton, I., & Linkie, M. (2011). Raiders of the Lost Bark: Orangutan foraging strategies in a degraded landscape. *PLoS ONE*, 6(6), e20962. <https://doi.org/10.1371/journal.pone.0020962>
- [27] Gippoliti, S. (2000). Orangutans in zoos: Husbandry, welfare and management in a typical arboreal solitary mammal. *International Zoo News*, 47, 356 – 368.
- [28] Camus, S. M. J., Blois-Heulin, C., Li, Q., Hausberger, M., & Bezard, E. (2013). Behavioural profiles in Captive-Bred cynomolgus macaques: towards monkey models of mental disorders? *PLoS ONE*, 8(4), e62141. <https://doi.org/10.1371/journal.pone.0062141>
- [29] Zheng-sheng, L., Feng, L., Li-Wei, T. and Xiao-Yu, Z. (2002). Time budget of semifree-ranging Amur tigers (*Panthera tigris altaica*). *Zoological Research*, 5, 389 – 439.
- [30] Sunquist, M. (2010). What is a tiger? Ecology and behavior. In *Tigers of the world. The Science, Politics and Conservation of Panthera tigris*, 2nd edition, Tilson, R. and Nhyus, P.J. (editors). Elsevier, London. 19 – 33.
- [31] Galardi, E.G., Fabbrani, M., Rausa, F.A., Preziosi, R., Brereton, J.E., Pastorino, Q.G. (2021). An investigation into the behavior, sociality, and enclosure use of group-housed lions and tigers. *Journal of Veterinary Medicine and Animal Sciences*, 4(1), 1068.
- [32] Smith, K.D., Snider, R.J., Dembiec, D. P., Siegford, J.M., & Ali, A.B. (2022). Effects of a modern exhibit design on captive tiger welfare. *Zoo Biology*, 42(3), 371–382. <https://doi.org/10.1002/zoo.21746>
- [33] Holland, A., Galardi, E. G., Fabbroni, M., Hashmi, A., Catinaud, J., Preziosi, R., Brereton, J. E., & Pastorino, G. Q. (2023). Exploration of social proximity and behavior in captive malayan tigers and their cubs. *Animals*, 13(6), 1040. <https://doi.org/10.3390/ani13061040>
- [34] Jepsen, E. M., Scheun, J., Dehnhard, M., Kumar, V., Umopathy, G., & Ganswindt, A. (2021). Non-invasive monitoring of glucocorticoid metabolite concentrations in native Indian, as well as captive and re-wilded tigers in South Africa. *General and Comparative Endocrinology*, 308, 113783. <https://doi.org/10.1016/j.ygcen.2021.113783>
- [35] Rose, P. E., Nash, S. M., & Riley, L. M. (2017). To pace or not to pace? A review of what abnormal repetitive behavior tells us about zoo animal management. *Journal of Veterinary Behavior*, 20, 11–21. <https://doi.org/10.1016/j.jveb.2017.02.007>
- [36] Birkett, L. P., & Newton-Fisher, N. E. (2011). How Abnormal Is the Behaviour of Captive, Zoo-Living Chimpanzees? *PLoS ONE*, 6(6), e20101. <https://doi.org/10.1371/journal.pone.0020101>
- [37] Liu, J., Chen, Y., Guo, L., Gu, B., Liu, H., Hou, A., Liu, X., Sun, L., & Liu, D. (2006). Stereotypic behavior and fecal cortisol level in captive giant pandas in relation to environmental enrichment. *Zoo Biology*, 25(6), 445–459. <https://doi.org/10.1002/zoo.20106>
- [38] Mason, G., Clubb, R., Latham, N. and Vickery, S. (2007). Why and how should we use environmental enrichment to tackle stereotypic behaviour. *Applied Animal Behaviour Sciences*, 102 (3), 163 – 188. <https://doi.org/10.1016/j.applanim.2006.05.041>
- [39] Hosey, G. R. (2004). How does the zoo environment affect the behavior of captive primates? *Applied Animal Behaviour Science*, 90(2), 107–129. <https://doi.org/10.1016/j.applanim.2004.08.015>
- [40] Morgan, K.N. and Tromborg, C.T. (2007). Sources of stress in captivity. *Applied Animal Behaviour Sciences*, 102 (3), 262 – 302. <https://doi.org/10.1016/j.applanim.2006.05.032>
- [41] Cinková, I. and Bičík, V. (2013). Social and reproductive behaviour of critically endangered northern white rhinoceros in zoological garden. *Mammalian Biology*, 78 (1), 50 – 54. <https://doi.org/10.1016/j.mambio.2012.09.007>
- [42] Swaisgood, R.R. (2007). Current status and future directions of applied behavioral research for animal welfare and conservation. *Applied Animal Behaviour Science*, 102(3/4), 139 – 162. <https://doi.org/10.1016/j.applanim.2006.05.027>

- [43] McEwen, B.S. (2000). The neurobiology of stress: from serendipity to clinical relevance. *Brain Research*, 886(1-2), 172 – 189. [https://doi.org/10.1016/s0006-8993\(00\)02950-4](https://doi.org/10.1016/s0006-8993(00)02950-4)
- [44] Pizzutto, C.S., Nichi, M., Sgai, M.G.F.G., Correa, S.H.R., Viau, P., Beresca, A.M., de Oliveira, C.A., Barnabe, R.C. and Vaz Guimarães, M.A. (2008). Effect of environmental enrichment on behavioral and endocrine aspects of a captive orangutan (*Pongo pygmaeus*). *Laboratory Primate Newsletter*, 47(2), 10 – 14.
- [45] Warwick, C., Arena, P., Lindley, S., Jessop, M., & Steedman, C. (2013). Assessing reptile welfare using behavioural criteria. *In Practice*, 35(3), 123–131. <https://doi.org/10.1136/inp.f1197>
- [46] Varadharajan, V., Krishnamoorthy, T., & Nagarajan, B. (2016). Prevalence of stereotypies and its possible causes among captive Asian elephants (*Elephas maximus*) in Tamil Nadu, India. *Applied Animal Behaviour Science*, 174, 137–146. <https://doi.org/10.1016/j.applanim.2015.10.006>
- [47] Pitsko, L.E. (2003). *Wild tigers in captivity: A study of the effects of the captive environment on tiger behavior*. Master Thesis. Virginia Polytechnic Institute and State University, Blacksburg, VA, USA.
- [48] Li, C., Juang, Z., Tang, S. and Zeng, Y. (2007). Evidence of effects of human disturbance on alert response in Pere David’s deer (*Elaphurus davidianus*). *Zoo Biology*, 26, 461 – 470. <https://doi.org/10.1002/zoo.20132>
- [49] Brummer, S. P., Gese, E. M., & Shivik, J. A. (2010). The effect of enclosure type on the behavior and heart rate of captive coyotes. *Applied Animal Behaviour Science*, 125(3–4), 171–180. <https://doi.org/10.1016/j.applanim.2010.04.012>
- [50] Breton, G. and Barrot, S., (2014). Influence of enclosure size on the distances covered and paced by captive tigers (*Panthera tigris*). *Applied Animal Behaviour Science*, 154:66–75. <https://doi.org/10.1016/j.applanim.2014.02.007>
- [51] Arumugam, K.A., Luan, L.Q., Wan Ibrahim, W.N., Mohd. Toh @Tah., M., Buesching, C.D. and Annavi, G. (2018). Influence of enclosure conditions and visitors on the behavior of captive Malayan tapir (*Tapirus indicus*): Implications for ex-situ management and conservation. *International Journal of Scientific and Research Publications*, 8 (7), 22 – 33. <https://doi.org/10.29322/ijsrp.8.7.2018.p7906>
- [52] Robeck, T.R., Willis, K., Scarpuzzi, M.R. and O’Brien, J.K. (2015). Comparisons of life-history parameters between free-ranging and captive killer whale (*Orcinus orca*) populations for application toward species manahement. *Journal of Mammalogy*, 96(5), 1055 – 1070. <https://doi.org/10.1093/jmammal/gyv113>
- [53] Lahdenpera, M., Mark, K.U., Courtiol, A. and Lummaa, V. (2018). Differences in age-specific mortality between wild-caught and captive-born Asian elephants. *Nature Communications*, 9, 1 – 10. <https://doi.org/10.1038/s41467-018-05515-8>
- [54] Yasmeen, R., Aslam, I., Ahmad, M., & Ali Shah, M.H. (2022). Zoochosis: A short review on stereotypical behavior of captive animals. *Journal of Wildlife and Biodiversity*, 7(2), 8–20. <https://doi.org/10.5281/zenodo.7362442>
- [55] Peng, S.; Broom, D.M. (2021). The sustainability of keeping birds as pets: should any be kept? *Animals*, 11, 582. <https://doi.org/10.3390/ani11020582>
- [56] Mohapatra, R.K., Panda, S. And Acharya, U.R. (2014). Study on activity pattern and incidence of stereotypic behavior in captive tigers. *Journal of Veterinary Behavior*, 1 – 5. <https://doi.org/10.1016/j.jveb.2014.04.003>
- [57] Scherpenhuizen, J.M., 2022. Evaluation of the welfare and reproductive biology of captive tigers using non-invasive techniques: Behaviour and hormone monitoring. Thesis, Charles Sturt University, Australia.
- [58] Rollin, B. (2006). *Science and ethics*. Cambridge University Press, Cambridge, UK. 304.
- [59] Shih, H-Y., Yu, J-F. and Wang, L-C. (2016). Stereotypic behaviours in bears. *Taiwan Veterinary Journal*, 42(1), 11 – 17. <https://doi.org/10.1142/s168264851530004x>

- [60] Clubb, R., & Mason, G.J. (2007). Natural behavioural biology as a risk factor in carnivore welfare: How analysing species differences could help zoos improve enclosures. *Applied Animal Behaviour Science*, 102(3–4), 303–328. <https://doi.org/10.1016/j.applanim.2006.05.033>
- [61] Boccacino, D., Maia, C. M., Santos, E. F. D., & Santori, R. T. (2018). Effects of environmental enrichments on the behaviors of four captive jaguars: individuality matters. *Oecologia Australis*, 22(01), 63–73. <https://doi.org/10.4257/oeco.2018.2201.06>
- [62] Miller, L., Vicino, G., Sheftel, J., & Lauderdale, L. (2020). Behavioral diversity as a potential indicator of positive animal welfare. *Animals*, 10(7), 1211. <https://doi.org/10.3390/ani10071211>
- [63] Greenwood, V. J., Smith, E. L., Goldsmith, A. R., Cuthill, I. C., Crisp, L. H., Walter-Swan, M. B., & Bennett, A. T. (2004). Does the flicker frequency of fluorescent lighting affect the welfare of captive European starlings? *Applied Animal Behaviour Science*, 86(1–2), 145–159. <https://doi.org/10.1016/j.applanim.2003.11.008>
- [64] Brilot, B. O., Asher, L., Feenders, G., & Bateson, M. (2009). Quantification of abnormal repetitive behaviour in captive European starlings (*Sturnus vulgaris*). *Behavioural Processes*, 82(3), 256–264. <https://doi.org/10.1016/j.beproc.2009.07.003>
- [65] Kumar, V. (2021). Clinical management of maggot wounds in Asiatic Black Bear (*Ursus thibetanus*). *Scientific Reports in Life Sciences*, 2(1), 7–12. <https://doi.org/10.22034/srsls.2020.520564.1006>
- [66] Tatemoto, P., Broom, D. M., & Zanella, A. J. (2022). Changes in Stereotypies: Effects over Time and over Generations. *Animals*, 12(19), 2504. <https://doi.org/10.3390/ani12192504>
- [67] Joshi, S., & Pillay, N. (2016). Association between personality and stereotypic behaviours in the African striped mouse *Rhabdomys dilectus*. *Applied Animal Behaviour Science*, 174, 154–161. <https://doi.org/10.1016/j.applanim.2015.11.021>
- [68] Joshi, S., & Pillay, N. (2018). Is wheel running a re-directed stereotypic behaviour in striped mice *Rhabdomys dilectus*? *Applied Animal Behaviour Science*, 204, 113–121. <https://doi.org/10.1016/j.applanim.2018.04.011>
- [69] Hogan, L. and Tribe, A. (2007). Prevalence and cause of stereotypic behaviour in common wombats (*Vombatus ursinus*) residing in Australian zoos. *Applied Animal Behaviour Science* 105 (1-3) 180-191. <https://doi.org/10.1016/j.applanim.2006.06.006>
- [70] Jeppesen, L., Heller, K., & Bildsøe, M. (2004). Stereotypies in female farm mink (*Mustela vison*) may be genetically transmitted and associated with higher fertility due to effects on body weight. *Applied Animal Behaviour Science*, 86(1–2), 137–143. <https://doi.org/10.1016/j.applanim.2003.11.011>
- [71] Liu, H., Duan, H., & Wang, C. (2017). Effects of ambient environmental factors on the stereotypic behaviors of giant pandas (*Ailuropoda melanoleuca*). *PLoS ONE*, 12(1), e0170167. <https://doi.org/10.1371/journal.pone.0170167>
- [72] Qin, D., Rizak, J., Chu, X., Li, Z., Yang, S., Lü, L., Yang, L., Yang, Q., Yang, B., Pan, L., Yin, Y., Chen, L., Feng, X., & Hu, X. (2015). A spontaneous depressive pattern in adult female rhesus macaques. *Scientific Reports*, 5(1). <https://doi.org/10.1038/srep11267>
- [73] Xu, F., Wu, Q., Xie, L., Gong, W., Zhang, J., Zheng, P., Zhou, Q., Ji, Y., Wang, T., Li, X., Fang, L., Li, Q., Yang, D., Li, J., Melgiri, N. D., Shively, C., & Xie, P. (2015). Macaques exhibit a Naturally-Occurring depression similar to humans. *Scientific Reports*, 5(1). <https://doi.org/10.1038/srep09220>
- [74] Martin, M. S., Owen, M., Wintle, N. J. P., Zhang, G., Zhang, H., & Swaisgood, R. R. (2020). Stereotypic behavior predicts reproductive performance and litter sex ratio in giant pandas. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-63763-5>
- [75] Khan, A. S., Lea, S. E. G., Chand, P., Rai, U., & Baskaran, N. (2022). Predictors of psychological stress and behavioral diversity among captive red panda in Indian zoos and

- their implications for global captive management. *Scientific Reports*, 12(1). <https://doi.org/10.1038/s41598-022-17872-y>
- [76] Abdul Mawah, S.S., Chor-Wai, L. and Jasnie, F. (2021). Comparative study on daily activity budget of sun bear (*Helarctus malayanus*) in captive and semi-captivity. *Malaysian Applied Biology*, 50(1), 115 – 124. <https://doi.org/10.55230/mabjournal.v50i1.18>
- [77] Johnson, B., and Langton, J. (2021). Behaviour change in Amur tigers *Panthera tigris altaica* after an enclosure move. *Journal of Zoo and Aquarium Research*, 9(3). <https://doi.org/10.19227/jzar.v9i3.520>
- [78] Poirier, C., Oliver, C. J., Bueno, J. C., Flecknell, P., & Bateson, M. (2019). Pacing behaviour in laboratory macaques is an unreliable indicator of acute stress. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-43695-5>
- [79] Bashaw, M.J., Kelling, A.S., Bloomsmith, M.A. and Maple, T.L. (2007). Environmental effects on the behavior of zoo-housed lions and tigers, with a case study of the effects of a visual barrier on pacing. *Journal of Applied Animal Welfare Science*, 10(2), 95 – 109. <https://doi.org/10.1080/10888700701313116>
- [80] Mallapur, A. and Chellam, R. (2002). Environmental influences on stereotypy and the activity budget of Indian leopards (*Panthera pardus*) in four zoos in southern India. *Zoo Biology*, 21(6), 585 – 595. <https://doi.org/10.1002/zoo.10063>
- [81] Broom, D.M. (2022). Broom and Fraser's domestic animal behaviour and welfare, 6th ed., CABI, Wallingford, UK, p. 545. <http://dx.doi.org/10.1079/9/9781789249835.0001>
- [82] Polverino, G., Manciocco, A., Vitale, A., & Alleva, E. (2015). Stereotypic behaviours in *Melopsittacus undulatus*: Behavioural consequences of social and spatial limitations. *Applied Animal Behaviour Science*, 165, 143–155. <https://doi.org/10.1016/j.applanim.2015.02.009>
- [83] Simcharoen, A., Savini, T., Gale, G.A., Simcharoen, S., Duangchantrasiri, S., Pakpien, S., & Smith, J. L. (2014). Female tiger *Panthera tigris* home range size and prey abundance: important metrics for management. *Oryx*, 48(3), 370–377. <https://doi.org/10.1017/s0030605312001408>
- [84] Chundawat, R. S., Sharma, K., Gogate, N., Malik, P. K., & Vanak, A. T. (2016). Size matters: Scale mismatch between space use patterns of tigers and protected area size in a Tropical Dry Forest. *Biological Conservation*, 197, 146–153. <https://doi.org/10.1016/j.biocon.2016.03.004>
- [85] Maulana, R., Gawi, J.M., & Utomo, S.W. (2020). Architectural design assessment of Javan leopard rehabilitation facility regarding the occurrence of stereotypical pacing. *IOP Conference Series Earth and Environmental Science*, 426(1), 012075. <https://doi.org/10.1088/1755-1315/426/1/012075>
- [86] Williams, I.J., Finch, K., Agnew, R., & Holmes, L. (2021). Effects of nearby construction work on the behavior of Asiatic lions (*Panthera leo persica*). *Journal of Zoological and Botanical Gardens*, 2, 66 – 74. <http://doi.org/10.3390/jzbg2010005>
- [87] Bauer, E., Babitz, M., Boedeker, N., & Hellmuth, H. (2013). Approaches to understanding and managing pacing in sloth bears in a zoological setting. *International Journal of Comparative Psychology*, 26(1). <https://doi.org/10.46867/ijcp.2013.26.01.04>
- [88] Mittal, S.K., Rao, R.J., Shakya, S., and Tripathi, S.M. (2019). Modern Naturalistic Enclosures: Comparatively an enhanced management practice of captive felids in the zoological park. *Journal of Animal Research*, 9(1). <https://doi.org/10.30954/2277-940x.01.2019.14>
- [89] Mohapatra, R.K., Mishra, A.K., Parida, S.P., and Mishra, S. (2010). Behavioural responses to environmental enrichment in captive tigers (*Panthera tigris*) at Nandankanan Zoological Park, Orissa. *e-planet*, 8(2), 44 – 48.

- [90] Montaudouin, S., and Le Pape, G. (2004). Comparison of the behaviour of European brown bears (*Ursus arctos arctos*) in six different parks, with particular attention to stereotypes. *Behavioural Processes*, 67(2), 235 – 244. <https://doi.org/10.1016/j.beproc.2004.02.008>
- [91] Donald, K., Benedetti, A., Goulart, V.D. L. R., Deming, A., Nollens, H., Stafford, G., & Brando, S. (2023). Environmental enrichment devices are safe and effective at reducing undesirable behaviors in California sea lions and northern elephant seals during rehabilitation. *Animals*, 13(7), 1222. <https://doi.org/10.3390/ani13071222>
- [92] Veeraselvam, M., Sridhar, R., Jayathangaraj, M.G. and Perumal, P. (2013). Behavioural study of captive sloth bears using environmental enrichment tools. *International Journal of Zoology*, 526905. <https://doi.org/10.1155/2013/526905>
- [93] Stokes, J. (2014). Observation in behavioural patterns of Bornean sun bears *Helarctos malayanus euryspilus* in Rehabilitation Center Tropical Rainforest Reserve in Sabah. Thesis, Oregon State University, USA.
- [94] Gruber, T.M., Friend, T.H., Gardner, J.M., Packard, J.M., Beaver, B. and Bushong, D. (2000). Variation in stereotypic behavior related to restraint in circus elephants. *Zoo Biology*, 19, 209 – 221. [https://doi.org/10.1002/1098-2361\(2000\)19:3](https://doi.org/10.1002/1098-2361(2000)19:3)
- [95] Haspesslagh, M., Stevens, J.M.G., De Groot, E., Dewulf, J., Kalmar, I.D. and Moons, C.P.H. (2013). A survey of foot problems, stereotypic behaviour and floor type in Asian elephants (*Elephas maximus*) in European zoos. *Animal Welfare*, 22, 437 – 443. <https://doi.org/10.7120/09627286.22.4.437>
- [96] Bettinger, T., Larry, M., Goldstein, M. and Laudenslager, M. (1997). Plasma cortisol concentrations and behavioral traits of two female Asian elephants. *American Zoological and Aquarium Association Annual Conference Proceedings*, 13 – 17 September 1997. Albuquerque Biological Park, Albuquerque, NM, USA. 88 -90.
- [97] Arakawa, H. (2021). Implication of the social function of excessive self-grooming behavior in BTBR T+*lpr*3^{tf}/J mice as an idiopathic model of autism. *Physiology and Behavior*, 237, 113432. <https://doi.org/10.1016/j.physbeh.2021.113432>.
- [98] Mooring, M. (2024). Programmed grooming after 30 years of study: A review of evidence and future prospects. *Animals*, 14(9), 1266. <https://doi.org/10.3390/ani14091266>
- [99] Blank, D. A. (2022). Grooming behavior in goitered gazelles: the programmed versus stimulus-driven hypothesis. *Ethology Ecology & Evolution*, 35(1), 62–82. <https://doi.org/10.1080/03949370.2021.2015449>
- [100] Van Zeeland, Y.R.A., Spruit, B.M., Rodenburg, T.B., Riedstra, B., Van Hierden, Y.M., Buitenhuis, B., Korte, S.M. and Lumeij, J.T. (2009). Feather damaging behavior in parrots: A review with consideration of comparative aspects. *Applied Animal Behaviour Science*, 121(2), 75 – 95. <https://doi.org/10.1016/j.applanim.2009.09.006>
- [101] Estanislau, C., Veloso, A.W., Filgueiras, G.B., Maio, T.P., Dal-Cól, M.L., Cunha, D.C., Klein, R., Carmona, L.F., Fernández-Teruel, A., (2019). Rat self-grooming and its relationships with anxiety, dearousal and perseveration: Evidence for a self-grooming trait. *Physiology and Behavior*, 209, 112585. <https://doi.org/10.1016/j.physbeh.2019.112585>.
- [102] Clayton, M., and Shrock, T., (2020). Making a tiger’s day: free-operant assessment and environmental enrichment to improve the daily lives of captive Bengal tigers (*Panthera tigris tigris*). *Behavior Analysis in Practice*, 13(4), 883–893. <https://doi.org/10.1007/s40617-020-00478-z>.
- [103] Klenowski, P.M., Zhao-Shea, R., Freels, T.G., Molas, S., Zinter, M., M’Angale, P., Xiao, C., Martinez-Núñez, L., Thomson, T., & Tapper, A.R. (2023). A neuronal coping mechanism linking stress-induced anxiety to motivation for reward. *Science Advances*, 9(49). <https://doi.org/10.1126/sciadv.adh9620>
- [104] Pomerantz, O., Meiri, S., & Terkel, J. (2013). Socio-ecological factors correlate with levels of stereotypic behavior in zoo-housed primates. *Behavioural Processes*, 98, 85–91. <https://doi.org/10.1016/j.beproc.2013.05.005>

- [105] Morris, Megan Colleen, "Treatment Analysis of a Captive Male Jaguar (*Panthera onca*)" (2018). UNF Graduate Theses and Dissertations. 799. <https://digitalcommons.unf.edu/etd/799>
- [106] Chitty, J. (2003). Feather plucking in psittacine birds 2. Social, environmental and behavioural considerations. *Practice*, 25(9), 550–555. <https://doi.org/10.1136/inpract.25.9.550>
- [107] Bellows, J., Center, S., Daristotle, L., Estrada, A. H., Flickinger, E. A., Horwitz, D. F., Lascelles, B. D. X., Lepine, A., Perea, S., Scherk, M., & Shoveller, A. K. (2016). Aging in cats. *Journal of Feline Medicine and Surgery*, 18(7), 533–550. <https://doi.org/10.1177/1098612x16649523>
- [108] Hopper, L. M., Freeman, H. D., & Ross, S. R. (2016). Reconsidering coprophagy as an indicator of negative welfare for captive chimpanzees. *Applied Animal Behaviour Science*, 176, 112–119. <https://doi.org/10.1016/j.applanim.2016.01.002>
- [109] Goldsborough, Z., Sterck, E., De Waal, F., & Webb, C. (2022). Individual variation in chimpanzee (*Pan troglodytes*) repertoires of abnormal behavior. *Animal Welfare*, 31(1), 125–135. <https://doi.org/10.7120/09627286.31.1.011>
- [110] Videvall, E., Bensch, H. M., Engelbrecht, A., Cloete, S., & Cornwallis, C. K. (2023). Coprophagy rapidly matures juvenile gut microbiota in a precocial bird. *Evolution Letters*, 7(4), 240–251. <https://doi.org/10.1093/evlett/grad021>
- [111] Krief, S., Jamart, A., & Hladik, C. (2004). On the possible adaptive value of coprophagy in free-ranging chimpanzees. *Primates*, 45(2), 141–145. <https://doi.org/10.1007/s10329-003-0074-4>
- [112] Jacobson, S. L., Ross, S. R., & Bloomsmith, M. A. (2016). Characterizing abnormal behavior in a large population of zoo-housed chimpanzees: prevalence and potential influencing factors. *PeerJ*, 4, e2225. <https://doi.org/10.7717/peerj.2225>
- [113] Blyton, M. D. J., Soo, R. M., Hugenholtz, P., & Moore, B. D. (2022). Maternal inheritance of the koala gut microbiome and its compositional and functional maturation during juvenile development. *Environmental Microbiology*, 24(1), 475–493. <https://doi.org/10.1111/1462-2920.15858>
- [114] Kobayashi, A., Tsuchida, S., Ueda, A., Yamada, T., Murata, K., Nakamura, H., & Ushida, K. (2019). Role of coprophagy in the cecal microbiome development of an herbivorous bird Japanese rock tarmigan. *Journal of Veterinary Medical Science*, 81(9), 1389–1399. <https://doi.org/10.1292/jvms.19-0014>
- [115] Phillips, C. J., Tribe, A., Lisle, A., Galloway, T. K., & Hansen, K. (2017). Keepers' rating of emotions in captive big cats, and their use in determining responses to different types of enrichment. *Journal of Veterinary Behavior*, 20, 22–30. <https://doi.org/10.1016/j.jveb.2017.03.006>
- [116] Brand, C. M., Boose, K. J., Squires, E. C., Marchant, L. F., White, F. J., Meinelt, A., & Snodgrass, J. J. (2016). Hair plucking, stress, and urinary cortisol among captive bonobos (*Pan paniscus*). *Zoo Biology*, 35(5), 415–422. <https://doi.org/10.1002/zoo.21320>
- [117] Goldsborough, Z. (2017). The effect of visitor density on the behavior of two Siberian tigers (*Panthera tigris altaica*) housed in a zoo enclosure. *Reinvention: an International Journal of Undergraduate Research*, BLASTER 2017, Special Issue.
- [118] Babb, M.H., Goforth, K.M., Lohmann, K.J., & Lohmann, C.M.F. (2020). Behavioural comparison of cougars (*Puma concolor*) and lions (*Panthera leo*) between zoo and sanctuary. Thesis, University of North Carolina. <https://doi.org/10.17615/ksqn-qw63>
- [119] Fijn, L.B., Van Der Staay, F.J., Goerlich-Jansson, V.C., & Arndt, S.S. (2020). Importance of basic research on the causes of feather pecking in relation to welfare. *Animals*, 10(2), 213. <https://doi.org/10.3390/ani10020213>
- [120] Rodenburg, T., Van Krimpen, M., De Jong, I., De Haas, E., Kops, Riedstra, B., Nordquist, R., Wagenaar, J., Bestman, M., & Nicol, C. (2013). The prevention and control of feather

- pecking in laying hens: identifying the underlying principles. *World's Poultry Science Journal*, 69(2), 361–374. <https://doi.org/10.1017/s0043933913000354>
- [121] Rodenburg, T.B., De Reu, K., & Tuytens, F.A.M. (2012). Performance, welfare, health and hygiene of laying hens in non-cage systems in comparison with cage systems. In CABI eBooks, pp. 210–224. <https://doi.org/10.1079/9781845938246.0210>
- [122] Giljam, P. (2021). Self mutilation as problem behaviour. *Zoospensefull*. Retrieved date: 11 December 2024. <https://zoospensefull.com/2021/08/02/self-mutilation-as-problem-behaviour/>
- [123] Santos, I.C., Rodrigues, M.C., Silva, F.A.N. & Campos, M.A.S. (2022). Automutilation in real vulture (*Sarcorampus papa*) maintained in captivity. *CABI Databases*, 25(2), e8154. <https://doi.org/10.25110/arqvet.v25i2conv.8154>
- [124] Dorey, N. R., Rosales-Ruiz, J., Smith, R., Lovelace, B., & Roane, H. (2009). Functional Analysis And Treatment Of Self-Injury In A Captive Olive Baboon. *Journal of Applied Behavior Analysis*, 42(4), 785–794. <https://doi.org/10.1901/jaba.2009.42-785>
- [125] Cronin, G. M., & Glatz, P. C. (2020). Causes of feather pecking and subsequent welfare issues for the laying hen: a review. *Animal Production Science*, 61(10), 990–1005. <https://doi.org/10.1071/an19628>
- [126] Stanton, L. A., Sullivan, M. S., & Fazio, J. M. (2015). A standardized ethogram for the felidae: A tool for behavioral researchers. *Applied Animal Behaviour Science*, 173, 3–16. <https://doi.org/10.1016/j.applanim.2015.04.001>
- [127] Williams, J., & Randle, H. (2017). Is the expression of stereotypic behavior a performance-limiting factor in animals? *Journal of Veterinary Behavior*, 20, 1–10. <https://doi.org/10.1016/j.jveb.2017.02.006>
- [128] Broom, D. M., & Johnson, K. G. (2019). Stress and animal welfare. In *Animal welfare*. <https://doi.org/10.1007/978-3-030-32153-6>
- [129] Lewis, K., Parker, M. O., Proops, L., & McBride, S. D. (2022). Risk factors for stereotypic behavior in captive ungulates. *Proceedings of the Royal Society B Biological Sciences*, 289(1983). <https://doi.org/10.1098/rspb.2022.1311>
- [130] Olsen, A., Kristensen, H.G., Iversen, K.W., Pedersen, N.H., Pertoldi, C., Alstrup, A.K.O., Jensen, T.H., & Pagh, S. (2020). Assessment of abnormal behavior and the effect of enrichment on captive chimpanzees in Aalborg Zoo. *Genetics and Biodiversity Journal*, 4(2), 73–91. <https://doi.org/10.46325/gabj.v4i2.99>
- [131] Mason, G. and Latham, N.R. (2004). Can't stop, won't stop: Is stereotypy a reliable animal welfare indicator? *Animal Welfare*, 13, S57 – S69.
- [132] Wickins-Drazilova, D. (2006). Zoo animal welfare. *Journal of Agricultural and Environmental Ethics*, 19, 27 – 36. <http://dx.doi.org/10.1007/s10806-005-4380-2>.
- [133] Kamaruzaman, A.S., Samsaimon, N.S., Hamzah, M.L., Amir, A., Hambali, K., Ramli, M.N. and Caiwu, L. (2017). Daily activity budget, feeding preferences and habitat choices of giant panda (*Ailuropoda melanoleuca*) at Zoo Negara Malaysia. *Malayan Nature Journal*, 69(3), 183 – 192.
- [134] Liu, D., Wang, Z., Tian, H., Yu, C., Zhang, G., Wei, R. and Zhang, H. (2003). Behavior of giant pandas (*Ailuropoda melanoleuca*) in captive conditions: gender differences and enclosure effects. *Zoo Biology*, 22, 77 – 82.
- [135] Rault, J.L. (2012). Friends with benefits: social support and its relevance for farm animal welfare. *Applied Animal Behaviour Science*, 136, 1 – 14.
- [136] Salas, M., Temple, D., Abáigar, T., Cuadrado, M., Delclaux, M., Enseñat, C., Almagro, V., Martínez-Nevado, E., Quevedo, M.A., Carbajal, A., Tallo-Parra, O., Sabés-Alsina, M., Amat, M., Lopez-Bejar, M., Fernández-Bellon, H. and Manteca, X. (2016). Aggressive behavior and hair cortisol levels in captive Dorcas Gazelles (*Gazella Dorcas*) as animal-based welfare indicators. *Zoo Biology*, 9999, 1 – 7.

- [137] Nelson, R.J. (1995). Aggression and social behaviour. In *An Introduction to behavioural endocrinology*, Nelson, R.J. (editor). Sinauer Associates Inc. Publisher, Sunderland. 443 – 484.
- [138] Hinton, M.G., Bendelow, A., Lantz, S. et al. (2013). Patterns of aggression among captive American flamingos (*Phoenicopterus ruber*). *Zoo Biology*, 32, 445 – 453.
- [139] Frumkin, N.B., Wey, T.W., Exnicios, M., Benham, C., Hinton, M.G., Lantz, S., Atherton, C., Forde, D. and Karubian, J. (2016). Inter-annual patterns of aggression and pair bonding in captive American Flamingos (*Phoenicopterus ruber*). *Zoo Biology*, 35, 111 – 119.
- [140] Gruen, L. (Ed.). (2014). *The Ethics of Captivity*. Oxford University Press.
- [141] Minter, B.A., & Collins, J.P. (2013). Ecological ethics in captivity: Balancing values and responsibilities in zoo and aquarium research under rapid global change. *Institute for Laboratory Animal Research Journal*, 54(1), 41–51.
- [142] Barber, J. C. (2009). Programmatic approaches to assessing and improving animal welfare in zoos and aquariums. *Zoo Biology*, 28(6), 519–530. <https://doi.org/10.1002/zoo.20260>
- [143] Gardiánová, I., and Bolechová, P. (2014). Tigers stereotypic pacing and enrichment. *Annual Research and Review in Biology*, 4(10):1544 – 1550. <https://doi.org/10.9734/arrb/2014/6487>
- [144] Maple, T. L. (2007). Toward a science of welfare for animals in the zoo. *Journal of Applied Animal Welfare Science*, 10(1), 63–70. <https://doi.org/10.1080/10888700701277659>
- [145] Sureshmarimuthu, N. S., Babu, S., Kumara, N. H. N., & Rajeshkumar, N. N. (2021). Factors influencing the flush response and flight initiation distance of three owl species in the Andaman Islands. *Journal of Threatened Taxa*, 13(11), 19500–19508. <https://doi.org/10.11609/jott.7339.13.11.19500-19508>
- [146] Gomes, D., McSweeney, L., and Santos, M. (2019). Effects of environmental enrichment techniques on stereotypical behaviors of captive Sumatran tigers: A preliminary case study. *Journal of Animal Behaviour and Biometeorology*, 7, 144 – 148. <https://doi.org/10.31893/2318-1265jabb.v7n4p144-148>
- [147] McPhee, M. E. (2002). Intact carcasses as enrichment for large felids: Effects on on- and off-exhibit behaviors. *Zoo Biology*, 21(1), 37–47. <https://doi.org/10.1002/zoo.10033>
- [148] Burgener, N., Gusset, M., & Schmid, H. (2008). Frustrated Appetitive Foraging Behavior, Stereotypic Pacing, and Fecal Glucocorticoid Levels in Snow Leopards (*Uncia uncia*) in the Zurich Zoo. *Journal of Applied Animal Welfare Science*, 11(1), 74–83. <https://doi.org/10.1080/10888700701729254>
- [149] Quirke, T., & Riordan, R. M. O. (2011). The effect of different types of enrichment on the behaviour of cheetahs (*Acinonyx jubatus*) in captivity. *Applied Animal Behaviour Science*, 133(1–2), 87–94. <https://doi.org/10.1016/j.applanim.2011.05.004>
- [150] De Souza Resende L, Gomes KCP, Andriolo A, Genaro G, Remy GL, De Almeida Ramos V (2011) Influence of cinnamon and catnip on the stereotypical pacing of onyx cats (*Leopardus tigrinus*) in captivity. *Journal of Applied Animal Welfare Science*, 14(3):247–254. <https://doi.org/10.1080/10888705.2011.576981>
- [151] Vidal L, Guilherme FR, Silva VF, Faccio MCSR, Martins M, Briani DC (2016) The effect of visitor number and spice provisioning in pacing expression by jaguars evaluated through a case study. *Brazilian Journal of Biology*, 76(2):506–510. <https://doi.org/10.1590/1519-6984.22814>
- [152] Antonenko TV, Panchuk KA, Medvedeva YuE (2017) The influence of olfactory enrichment on the welfare of large felines in captivity. *Ukrainian Journal of Ecology*, 7(4):134–138.
- [153] Tarou, L.R., and Bashaw, M.J. (2007). Maximizing the effectiveness of environmental enrichment: Suggestions from the experimental analysis of behavior. *Applied Animal Behaviour Science*, 102(3–4):189–204. <https://doi.org/10.1016/j.applanim.2006.05.026>

- [154] Wells, D. L. (2009). Sensory stimulation as environmental enrichment for captive animals: A review. *Applied Animal Behaviour Science*, 118(1–2), 1–11. <https://doi.org/10.1016/j.applanim.2009.01.002>
- [155] Wells, D. L. & Egli, J. (2004). The influence of olfactory enrichment on the behavior of captive black-footed cats, *Felis nigripes*. *Applied Animal Behaviour Science*, 85(1–2):107–119. <https://doi.org/10.1016/j.applanim.2003.08.013>
- [156] Mak, B., Francis, R. A., & Chadwick, M. A. (2021). Living in the concrete jungle: a review and socio-ecological perspective of urban raptor habitat quality in Europe. *Urban Ecosystems*, 24(6), 1179–1199. <https://doi.org/10.1007/s11252-021-01106-6>
- [157] Mishra, A. K., Charan Guru, B., & Patnaik, A. K. (2013). Effect of feeding Enrichment on behaviour of captive tigers. *Indian Zoo Yearbook*, 7, 124 – 133.
- [158] Vashisth, S., Singh, R., Singh, D.N., and Sethi, N. (2023). Evaluation of factors affecting the behavior of Bengal tigers (*Panthera tigris*) in captivity. *Journal of Wildlife and Biodiversity*, 7, 30 – 51. <https://doi.org/10.5281/zenodo.10019718>
- [159] White BC, Houser LA, Fuller JA, Taylor S, Elliott JL (2003) Activity-based exhibition of five mammalian species: Evaluation of behavioral changes. *Zoo Biology*, 22(3):269–285. <https://doi.org/10.1002/zoo.10085>
- [160] Ritzler CP, Lukas KE, Bernstein-Kurtycz LM, Koester DC (2021) The effects of choice-based design and management on the behavior and space use of zoo-housed Amur tigers (*Panthera tigris altaica*). *Journal of Applied Animal Welfare Science*, 26(2):256–269. <https://doi.org/10.1080/10888705.2021.1958684>
- [161] Biolatti, C., Modesto, P., Dezzutto, D., Pera, F., Tarantola, M., Gennero, M.S., Maurella, C., & Acutis, P.L. (2016). Behavioural analysis of captive tigers (*Panthera tigris*): A water pool makes the difference. *Applied Animal Behaviour Science*, 174, 173–180. <https://doi.org/10.1016/j.applanim.2015.11.017>
- [162] Doyle, R. (2016). Captive wildlife sanctuaries: Definition, ethical considerations, and public perception. *Animal Studies Journal*, 5(1), 5–28.
- [163] Grant, R. (2022). Captive animals: Perspectives, practices, challenges, and ethics. *Animals*, 12(10), 1300.
- [164] Bashir, M. (2020). Dietary ecology of Markhor (*Capra falconeri cashmiriensis*) in winter range of Kazinag National Park, Kashmir, J&K, India. *Indian Journal of Science and Technology*, 13(24), 2463–2474. <https://doi.org/10.17485/ijst/v13i24.432>
- [165] Hebert, P.L. and Bard, K. (2000). Orangutan use of vertical space in an innovative habitat. *Zoo Biology*, 19, 239 – 251.
- [166] De Rouck, M., Kitchener, A., Law, G., & Nelissen, M. (2005). A comparative study of the influence of social housing conditions on the behaviour of captive tigers (*Panthera tigris*). *Animal Welfare*, 14(3), 229–238. <https://doi.org/10.1017/s0962728600029390>
- [167] Miller, L., Bettinger, T., & Mellen, J. (2008). The reduction of stereotypic pacing in tigers (*Panthera tigris*) by obstructing the view of neighboring individuals. *Animal Welfare*, 17(3), 255–258. <https://doi.org/10.1017/s0962728600032176>
- [168] Macri, A. M., & Patterson-Kane, E. (2011). Behavioural analysis of solitary versus socially housed snow leopards (*Panthera uncia*), with the provision of simulated social contact. *Applied Animal Behaviour Science*, 130(3–4), 115–123. <https://doi.org/10.1016/j.applanim.2010.12.005>
- [169] Kelling, A. S., Allard, S. M., Kelling, N. J., Sandhaus, E. A., & Maple, T. L. (2012). Lion, ungulate, and visitor reactions to playbacks of lion roars at Zoo Atlanta. *Journal of Applied Animal Welfare Science*, 15(4), 313–328. <https://doi.org/10.1080/10888705.2012.709116>
- [170] Robbins, L., & Margulis, S. W. (2016). Music for the birds: effects of auditory enrichment on captive bird species. *Zoo Biology*, 35(1), 29–34. <https://doi.org/10.1002/zoo.21260>
- Perdue, B. M., Clay, A. W., Gaalema, D. E., Maple, T. L., & Stoinski, T. S. (2011). Technology at the zoo: The influence of a touchscreen

- computer on orangutans and zoo visitors. *Zoo Biology*, 31(1), 27–39. <https://doi.org/10.1002/zoo.20378>
- [171] Boostrom, H. (2013). Problem-Solving with orangutans (*Pongo pygmaeus* and *Pongo abelii*) and chimpanzees (*Pan troglodytes*): Using the IPAD to provide novel enrichment opportunities. <https://oaktrust.library.tamu.edu/items/6036ad48-9b23-49c1-a15e-b3c7a7b82bfa>
- [172] Webber, S., Carter, M., Smith, W., & Vetere, F. (2016). Interactive technology and human-animal encounters at the zoo. *International Journal of Human-Computer Studies*, 98, 150–168. <https://doi.org/10.1016/j.ijhcs.2016.05.003>
- [173] Kim-McCormack, N. N., Smith, C. L., & Behie, A. M. (2016). Is interactive technology a relevant and effective enrichment for captive great apes? *Applied Animal Behaviour Science*, 185, 1–8. <https://doi.org/10.1016/j.applanim.2016.09.012>
- [174] Tarou, L.R., Kuhar, C., Adcock, D., Bloomsmith, M., & Maple, T. (2004). Computer-assisted enrichment for zoo-housed orangutans (*Pongo pygmaeus*). *Animal Welfare*, 13(4), 445–453. <https://doi.org/10.1017/s0962728600028712>
- [175] Chudeau, K.R., Johnson, S.P., & Caine, N.G. (2019). Enrichment reduces stereotypical behaviors and improves foraging development in rehabilitating Eastern Pacific Harbor Seals (*Phoca vitulina richardii*). *Applied Animal Behaviour Science*, 219, 104830. <https://doi.org/10.1016/j.applanim.2019.07.001>
- [176] Pedonti, J. (2016). Coping with captivity: Stereotypies in zoo animals. Thesis, New College of Florida, US.
- [177] Watters, J. V., & Meehan, C. L. (2006). Different strokes: Can managing behavioral types increase post-release success? *Applied Animal Behaviour Science*, 102(3–4), 364–379. <https://doi.org/10.1016/j.applanim.2006.05.036>
- [178] Stamps, J., & Groothuis, T. G. G. (2009). The development of animal personality: relevance, concepts and perspectives. *Biological Reviews/Biological Reviews of the Cambridge Philosophical Society*, 85(2), 301–325. <https://doi.org/10.1111/j.1469-185x.2009.00103.x>
- [179] Trillmich, F., & Hudson, R. (2011). The emergence of personality in animals: The need for a developmental approach. *Developmental Psychobiology*, 53(6), 505–509. <https://doi.org/10.1002/dev.20573>
- [180] Cannon, T.H., Heistermann, M., Hankison, S.J., Hockings, K.J., & McLennan, M.R. (2016). Tailored enrichment strategies and stereotypic behavior in captive individually housed macaques (*Macaca* spp.). *Journal of Applied Animal Welfare Science*, 19(2), 171–182. <https://doi.org/10.1080/10888705.2015.1126786>
- [181] Szokalski, M.S., Litchfield, C.A., & Foster, W.K. (2012). Enrichment for captive tigers (*Panthera tigris*): Current knowledge and future directions. *Applied Animal Behaviour Science*, 139(1–2), 1–9. <https://doi.org/10.1016/j.applanim.2012.02.021>
- [182] Law, G., & Kitchener, A.C. (2019). Twenty years of the tiger feeding pole: review and recommendations. *International Zoo Yearbook*, 54(1), 174–190. <https://doi.org/10.1111/izy.12249>
- [183] Godinez, A.M., & Fernandez, E.J. (2019). What is the zoo experience? How zoos impact a visitor’s behaviors, perceptions, and conservation efforts. *Frontiers in Psychology*, 10. <https://doi.org/10.3389/fpsyg.2019.01746>
- [184] Lichtenwaller, C. (2020). Effect of the Prospect of Transport on Captive Tiger Behavior and Fecal Cortisol in Naïve and Experienced Tigers. Thesis. University of Arkansas, US.
- [185] Nakamichi, M. (2007). Assessing the effects of new primate exhibits on zoo visitors’ attitudes and perceptions by using three different assessment methods. *Anthrozoös*, 20(2), 155–165. <https://doi.org/10.2752/175303707x207945>
- [186] Fernandez, E. J., Tamborski, M. A., Pickens, S. R., & Timberlake, W. (2009). Animal–visitor interactions in the modern zoo: Conflicts and interventions. *Applied Animal Behaviour Science*, 120(1–2), 1–8. <https://doi.org/10.1016/j.applanim.2009.06.002>