

UNIVERSITI TEKNOLOGI MARA

**GEOMETRIC PROPERTIES OF NEW
SUBCLASSES OF UNIVALENT AND
BI-UNIVALENT FUNCTIONS
ASSOCIATED WITH
SUBORDINATION AND
GENERALISED DIFFERENTIAL
OPERATOR**

MUNIRAH BINTI ROSSDY

Thesis submitted in fulfillment
of the requirements for the degree of
**Doctor of Philosophy
(Mathematics)**

**College of Computing, Informatics and
Mathematics**

October 2024

ABSTRACT

Let A denote the class of normalised analytic functions f on the unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$, in the form $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$. Let S be the class of functions $f \in A$ that are univalent in D . A function f in A is univalent if it is a one-to-one mapping. Based on Duren (1983), the theorem of Koebe's one-quarter showed the images of D for every univalent function, $f \in S$ enclosing $1/4$ radius disc. Hence, every function $f \in S$ has an inverse f^{-1} , which is defined by $f^{-1}(f(z)) = z, z \in D$, and $f(f^{-1}(\omega)) = \omega$ ($|\omega| < r_0(f)$; $r_0(f) \geq \frac{1}{4}$), with the power series expansion $g(\omega) = f^{-1}(\omega) = \omega - a_2 \omega^2 + (2a_2^2 - a_3) \omega^3 - (5a_2^3 - 5a_2 a_3 + a_4) \omega^4 + \dots$. Meanwhile, a function f is said to be bi-univalent in D if both f and its inverse f^{-1} are univalent in D . Let Σ denote the class of bi-univalent functions in D . This thesis is concerned with the geometric properties of the class of analytic univalent and bi-univalent functions in the complex plane. Thus, there are three major objectives to be achieved and the results are presented in three different chapters. The first objective is to apply the subordination method on the class of analytic univalent and bi-univalent functions involving radius and coefficient problems. The radii of starlikeness for concave univalent functions related to certain rational functions, lune, cardioid, and the exponential equation are obtained. Then, the Third Hankel Determinants $|a_3(a_2 a_4 - a_3^2) + a_4(a_2 a_3 - a_4) + a_5(a_3 - a_2^2)|$ and Generalised Zalcman Conjecture $|a_n a_m - a_{n+m-1}|$ of bi-univalent functions are acquired. Next, continuation from the subordination method in the previous objective, the second objective investigated the coefficient problems for the class of bi-univalent functions by employing subordination and a generalised differential operator. The initial coefficients $|a_2|$ and $|a_3|$ of Taylor-Maclaurin series associated with Fibonacci numbers, Chebyshev polynomials and Chebyshev polynomials with respect to symmetric and symmetric conjugate points defined by the Al-Oboudi operator are attained. Hence, the results obtained were an improvement and more generalised from the recent studies. Finally, the last objective is to introduce a new differential operator. Hence, motivated by the generalised differential operator in the second objective, this study proposed a new generalised differential operator by utilising the binomial series. This new operator is applied in obtaining the inclusion properties for the class of analytic univalent functions. Then, the results on covering theorem, distortion theorem, rotation theorem, growth theorem, and the radius of convexity for functions of the class of bi-univalent functions were derived by utilising the new differential operator. The final chapter of this thesis discussed on the suggestion problems for future research. In conclusion, this thesis introduces new subclasses of univalent and bi-univalent functions and extends several classical results in geometric function theory. The findings contribute significantly to the understanding of these classes, offering new techniques and insights for future research.

ACKNOWLEDGEMENT

Firstly, I offer my heartfelt gratitude to Allah S.W.T for bestowing upon me the strength and opportunity to pursue and culminate my journey to a doctorate degree. I am deeply indebted to my esteemed supervisors and mentors, Dr Rashidah Omar and Associate Professor Dr Shaharuddin Cik Soh, for their unwavering guidance and support throughout this endeavour.

I extend my sincere appreciation to the Ministry of Higher Education of Malaysia and Universiti Teknologi MARA. Their generous financial support through the Skim Latihan Akademik IPTA (SLAI) scholarship and the provision of study leave with full salary have been pivotal in realizing my academic aspirations.

Lastly, I dedicate this work to the pillars of my life, my parents, who nurtured and moulded me into the individual I stand as today. My husband and son, who have been my beacon of joy and hope, and my siblings, family and friends, whose unwavering encouragement and support served as my bedrock. I am immeasurably grateful for your presence and constant faith in me throughout this journey. Alhamdulillah.

TABLE OF CONTENTS

	Page
CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR'S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENT	v
TABLE OF CONTENTS	vi
LIST OF SYMBOLS	ix
 CHAPTER ONE INTRODUCTION	 1
1.1 Introduction	1
1.2 Analytic Functions	3
1.3 Univalent Functions	4
1.4 Subclasses of Analytic Univalent functions	6
1.5 Analytic Bi-Univalent functions	9
1.6 Radius Problem	11
1.7 Initial Coefficients	14
1.8 Hankel Determinant	16
1.9 Zalcman Conjecture	19
1.10 Subordination	21
1.11 Operators	22
1.12 Problem Statement	24
1.12.1 Problem 1 : Radius of Starlikeness for Concave Univalent Functions and Coefficient Problems of Zalcman Conjecture and Hankel Determinants	24
1.12.2 Problem 2 : Coefficient Bounds for Bi-Univalent Functions associated with a Generalised Operator	25
1.12.3 Problem 3 : Characterisation Properties of Analytic Bi-Univalent Functions associated with a Generalised Operator	26
1.13 Research Questions	27
1.14 Research Objectives	27

CHAPTER ONE

INTRODUCTION

1.1 Introduction

Complex analysis and geometric function theory (GFT) play a crucial role in both mathematics and physics, serving as foundational concepts in these fields. Complex analysis, also known as the theory of functions with complex variables, was intensively studied in the nineteenth century. Complex analysis is a branch of mathematics focused on complex functions, derivatives, and the intricacies of the complex plane. A complex function is a function that receives a complex number as input and another complex number as output. These functions can be represented as a mapping from points in the complex plane to other points in the complex plane. The derivatives of these functions are also very interesting and are used to study the behaviour of complex functions near points in the complex plane. The complex plane is a two-dimensional coordinate system in which the x-axis represents the real part, and the y-axis represents the imaginary part of a complex number.

Complex analysis also studies the properties of complex functions such as analytic functions, complex integrals, and power series representations. If a complex function is differentiable at every point in its domain, it is called analytic. The derivatives of analytic functions have significant effects on the behaviour of the function. For example, the Cauchy-Riemann equations are a set of differential equations that complex analytic functions must satisfy. These equations have several important implications for the behaviour and characteristics of analytic functions, specifically demonstrating that these functions have the properties of local conformance and the ability to preserve angles between curves.

Complex integration is another important concept in complex analysis. Complex functions can be integrated along curves in the complex plane, just as real functions can be integrated along curves in the real plane. The contour integral is the most common form of the complex integral, where a function is integrated along a closed path in the complex plane. A key result in this area is Cauchy's theorem, which states that the value of a contour integral depends only on the homotopy class of the curve and not on the actual curve itself.