

JCHS

Journal of Clinical and Health Sciences

#2
2026

SUPPLEMENTARY ISSUE

Malaysian Obesity Society
MYOS



ND **2 NATIONAL OBESITY SCIENTIFIC CONFERENCE**

ABSTRACT BOOK

11th - 12th April 2026
**Connexion Conference &
Event Centre,**
Kuala Lumpur, Malaysia

Theme
**Obesity as a Treatable
Chronic Disease:
Science, Strategy,
and Success**

Issue **September 2026** | Volume **11** Issue **02 (Supplementary)**
eISSN 0127-984x
jchs-medicine.uitm.edu.my



**UNIVERSITI
TEKNOLOGI
MARA**

PD-48

Association Between Diabetes and the Prevalence of Diabetic Peripheral Neuropathy – Sub Analysis from Previous Conducted Study

Mohd Zin S.K.¹, Mohamed Shah F.Z.¹, Mohd Fauzi N.A.², Abdul Ghani R.¹, Eddy Warman N.A.

¹Department of Endocrine, Universiti Teknologi MARA, Hospital Al-Sultan Abdullah

²Department of Neurology, Universiti Teknologi MARA, Hospital Al-Sultan Abdullah

Corresponding Author email: kaamilahzin@gmail.com

Introduction:

Obesity and diabetes have reached epidemic proportions worldwide in recent years. The coexistence of these conditions, termed diabetes, may further elevate the risk of developing diabetic peripheral neuropathy (DPN), a prevalent complication of diabetes. This study aimed to evaluate the association between obesity and the prevalence of DPN, as well as other microvascular complications of diabetes, among patients with T2DM

Methods:

This study represents a sub analysis of a previously conducted cross-sectional study carried out at UiTM Specialist Centre Sungai Buloh and Hospital Al-Sultan Abdullah. It involved patients aged 18 to 60 years with type 2 diabetes mellitus (T2DM) of more than 5 years duration (n = 132). Obesity was assessed using anthropometric measurements and defined as a body mass index (BMI) >27.5 kg/m². The diagnosis of DPN was established based on positive nerve conduction study (NCS) findings and other microvascular complications were confirmed from hospital records.

Results:

The study population had a mean age of 60.16 ± 10.28 years and a mean duration of diabetes of 14.82 ± 6.66 years. The overall prevalence of diabetic peripheral neuropathy (DPN) was 51.5% (n = 68). More than half of the participants were classified as obese (53%, n = 70), and among these individuals, 64% (n = 45) had DPN. Obesity was significantly associated with DPN, with individuals with obesity having approximately threefold higher odds of developing DPN compared to non-obese individuals (OR = 3.052, 95% CI: 1.500–6.210, p = 0.002). Additionally, nephropathy was also significantly associated with obesity (OR = 2.157, 95% CI: 1.044–4.453, p = 0.038).

Conclusion:

More than half of participants with diabetes were found to have DPN. Furthermore, the presence of obesity was significantly associated with nephropathy, highlighting the importance of early screening for diabetic microvascular complications especially in individuals with diabetes.

Keywords: Diabetic peripheral neuropathy, obesity, diabetes, nephropathy, nerve conduction study