

### Zika on Board

Ariza Adnan, Nurhuda Ismail

Faculty of Medicine, Universiti Teknologi MARA (UiTM), Selangor, Malaysia

The Olympics and Paralympic games bring joy to millions of people worldwide. This year it was held in Rio de Janeiro, Brazil amidst the Zika virus epidemic. On 1 February 2016, six months leading to the Olympics, the excitement was dampened by the World Health Organization declaration of the unprecedented vector-borne Zika virus (ZIKV) infection as a public health emergency of international concern [1]. At that point in time, people in Malaysia also felt the heat but had the consolation that the epidemic occurred across the Atlantic on the other side of the world. On 27 August 2016, Singapore reported the first local confirmed case of ZIKV infection in the city-state [2]. It was only a matter of time that Malaysia reported its first confirmed case of ZIKV infection on 2 September 2016 [3]. A 58-year old woman from Klang, Selangor was diagnosed as the first ZIKV case, who had earlier visited her daughter in Singapore who was infected by the ZIKV. As of 22 September 2016, the number of confirmed ZIKV cases in Malaysia has mounted to six with both Polynesia and Micronesia strains [4]. The occurrence of the disease in our continent brings to light how rapidly globalization and free movement of population across geographical borders can accelerate the arbovirus threat across the globe.

The ZIKV is spread through the bite of infected female *Aedes* mosquitoes and evidence has shown it can also spread via sexual and blood transmission [5]. Most of the cases are asymptomatic or subclinical while the symptomatic cases are self-limiting. Other manifestations include neurological (Guillain-Barré syndrome and meningoencephalitis) and autoimmune (thrombocytopenic purpura and leukopenia) complications. More alarming is the association of the virus with infants born with microcephaly as a result of pregnant mothers infected

with ZIKV with the risk of microcephaly ranges between 1-13% [6, 7], though the direct causal relationship is still under investigation.

In line with the international response, Malaysia has intensified the surveillance and management activities to control its ZIKV epidemic. These include clinical surveillance, laboratory surveillance, microcephaly and Guillain-Barre Syndrome case detection as well as preparedness and response at all ports of entry [8]. The public health delivery system in Malaysia has improved tremendously over the years following lessons learned from the emerging and re-emerging infectious diseases that affected the country over the recent years. Apart from those commendable measures; in light of this ZIKV outbreak, there are looming questions even though may appear elementary are nevertheless pertinent which the healthcare fraternity should address. Where do we go from here? What more do we need to know and do to help us manage and control this outbreak more efficiently and effectively? These questions would certainly pose a challenge to our public health especially when this arbovirus shares the same transmission vector with dengue and chikungunya i.e., *Aedes* mosquito where the authorities in Malaysia are still struggling to achieve a satisfactory control level in the country.

What shall we do? We need a paradigm shift. We need to look at the re-emergence of ZIKV in a bigger perspective and manage it accordingly. Thus, instead of reacting merely to the notified ZIKV cases, we need to start strategizing how the transmission dynamics of the arbovirus family can be altered; this possibly means to look out of the box for the solutions. The conventional measures for prevention and control should also be complemented with two other elements

which are often neglected and underestimated in most outbreak response i.e., effective communication and social mobilization [9]. We must actively engage in communication with the public to hasten the containment by using all available forms of social medium. With tons of information about the virus in the media, the message should emphasize more on health education; empowering community on the case reports, transmission routes and infection complications rather than general statements on impact and world reaction towards ZIKV [10]. Only then social mobilization, which is commonly underutilized, may help mitigate the social and economic impact during an outbreak. An informed public understands the limitation and the need for the community. Only then we will appreciate the ripple effects - they will bring the community on board, educate the community to actively participate in the outbreak management, and share the responsibility as well as the outcome. Even when the community is faced with great anxiety, an informed public would be able to understand and support any move or decision made by the authorities concerned.

Of late, the controversy which surrounded ZIKV in Malaysia involved the ethical issue in dealing with termination of pregnancy in women with possible ZIKV-related fetal brain abnormalities. This was following a statement made by the Mufti of the Federal Territory, saying that Muslim women could abort their pregnancies if they were infected by the Zika virus to avoid the adverse effect on the lives of their families or the baby itself [11]. In Malaysia, the current law does not provide for abortion for pregnant mothers infected with Zika unless the pregnancy poses a threat to the mother's life [12]. The recent Centres for Disease Control guideline does not include pregnancy termination as an option in managing suspected or confirmed Zika infection. It advocates monitoring the pregnancy with serial fetal ultrasounds in suspected or inconclusive cases and retest for ZIKV when ultrasound suggests abnormalities consistent with Zika infection and fall short in mentioning the alternative path of termination of pregnancy [13]. On the other hand, World Health Organization guideline mentions subtly on the discontinuation of pregnancy as a possible next step in the management of pregnancies with the likelihood of foetal brain abnormalities and states that

women who wish to discontinue their pregnancy should receive accurate information about their options to the full extent of the law [14]. The failure to include guidelines on the option of safe, legal termination of pregnancy in Zika-response strategies is not only an issue of reproductive rights but also an issue of reproductive justice [15]. At the time of writing, it is learned that the Ministry of Health of Malaysia will hold a discussion on the matter with the National Fatwa Council regarding termination of pregnancy for women infected by the Zika virus in order to reach a consensus. Irrespective of the outcome of the *fatwa*, we are in the opinion that whether a woman who wishes to carry her pregnancy to term or discontinue the pregnancy should be offered appropriate counselling so that she, together with her partner, will be able to make a fully informed choice on the next step of action.

Despite being a re-emerging disease, there is still much evidence required to effectively manage and control the ZIKV outbreak. The disease behaviour remains dynamic, and a concerted effort by the health authorities and policy makers in implementing the appropriate dynamic alignment to meet the challenges is imperative. It also requires heightened public awareness of personal responsibility which is of paramount importance. The public health preventive strategies remain the cornerstone in the control of this mosquito-borne disease.

## REFERENCES

1. World Health Organization. Director-General summarizes the outcome of the emergency committee regarding clusters of microcephaly and Guillain-Barré syndrome. <http://www.who.int/mediacentre/news/statements/2016/emergency-committee-zika-microcephaly/en/>. 2016. Accessed 10 Sept 2016.
2. Ministry of Health, Singapore. First case of locally transmitted Zika virus infection. [https://www.moh.gov.sg/content/moh\\_web/home/pressRoom/pressRoomItemRelease/2016/first-case-of-locally-transmitted-zika-virus-infection.html](https://www.moh.gov.sg/content/moh_web/home/pressRoom/pressRoomItemRelease/2016/first-case-of-locally-transmitted-zika-virus-infection.html). 2016. Accessed 10 Sept 2016.
3. Ministry of Health, Malaysia. Kenyataan akhbar KPK 3 September 2016 – Kes Zika jangkitan tempatan pertama di Malaysia.

- <https://kpkesehatan.com/2016/09/03/kenyataan-akhbar-kpk-3-september-2016-kes-zika-jangkitan-tempatan-pertama-di-malaysia/>. 2016. Accessed 10 Sept 2016.
4. Ministry of Health, Malaysia. Kenyataan akhbar situasi terkini Zika di Malaysia 22 September 2016. [http://www.moh.gov.my/index.php/database\\_stores/attach\\_download/337/792](http://www.moh.gov.my/index.php/database_stores/attach_download/337/792). 2016. Accessed 25 Sept 2016.
  5. Brooks JT, Friedman A, Kachur RE, Laflam M, Peters PJ, Jamieson DJ. Update: interim guidance for prevention of sexual transmission of Zika virus — United States, July 2016. *MMWR Morb Mortal Wkly Rep*. 2016; 65: 745-7. doi: <http://dx.doi.org/10.15585/mmwr.mm6529e2>.
  6. Johansson MA, Mier-y-Teran-Romero L, Reefhuis J, Gilboa SM, Hills SL. Zika and the risk of microcephaly. *N Engl J Med*. 2016; 375: 1-4.
  7. de Araújo TVB, Rodrigues LC, de Alencar Ximenes RA, de Barros Miranda-Filho D, Montarroyos UR, de Melo AP, Valongueiro S, Souza WV, Braga C, Brandao Filho SP, Cordeiro MT. Association between Zika virus infection and microcephaly in Brazil, January to May, 2016: preliminary report of a case-control study. *Lancet Infect Dis*. 2016. doi: 10.1016/S1473-3099(16)30318-8.
  8. Ministry of Health, Malaysia. “Updated Zika alert” dan arahan pentadbiran untuk pemantauan dan pengurusan jangkitan virus Zika 11 September 2016. <https://kpkesehatan.files.wordpress.com/2016/09/updated-zika-alert-arahan-pentadbiran-final-11-sept-2016.pdf>. 2016. Accessed 26 Sept 2016.
  9. World Health Organization. Outbreak communication. Best practices for communicating with the public during an outbreak. WHO. Singapore. [http://www.who.int/csr/resources/publications/WHO\\_CDS\\_2005\\_32web.pdf](http://www.who.int/csr/resources/publications/WHO_CDS_2005_32web.pdf). 2005. Accessed 10 Sept 2016.
  10. Fu KW, Liang H, Saroha N, Tse ZT, Ip P, Fung IC. How people react to Zika virus outbreaks on Twitter? A computational content analysis. *Am J Infect Control*. 2016. doi: 10.1016/j.ajic.2016.04.253.
  11. Fazleena A. Zika: Health Ministry to meet National Fatwa Council over medical advice for pregnant victims. *The NST online*. 2016. Accessed 20 Sept 2016.
  12. Fong LF. Subra: No abortions allowed for Zika-infected pregnant mothers. *The Star online*. <http://www.thestar.com.my/news/nation/2016/09/06/subra-on-abortions-for-those-with-zika/>. 2016. Accessed 20 Sept 2016.
  13. Centers for Disease Control and Prevention. Morbidity and mortality weekly report. Update: interim guidance for health care providers caring for women of reproductive age with possible Zika virus exposure — United States. [http://www.cdc.gov/mmwr/volumes/65/wr/mm6512e2.htm#F2\\_down](http://www.cdc.gov/mmwr/volumes/65/wr/mm6512e2.htm#F2_down). 2016. Accessed 20 Sept 2016.
  14. World Health Organization. Pregnancy management in the context of Zika virus infection: interim guidance update. 13 May 2016. WHO/ZIKV/MOC/16.2 Rev.1. [http://who.int/iris/bitstream/10665/204520/1/WHO\\_ZIKV\\_MOC\\_16.2\\_eng.pdf?ua=.1](http://who.int/iris/bitstream/10665/204520/1/WHO_ZIKV_MOC_16.2_eng.pdf?ua=.1). 2016. Accessed 20 Sept 2016.
  15. Aiken ARA, Aiken CE, Trussell J. In the midst of Zika pregnancy advisories, termination of pregnancy is the elephant in the room. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2016. doi: 10.1111/1471-0528.14296.

### Herd Immunity or Heard Not of Immunity?

Anis Siham Zainal Abidin, Masri Mohamed, Mazidah Nordin, Nor Azizah Abu, Noor Shafina Mohd Noor, Faisal Mohd Fadzli, Mohammed Fauzi Abdul Rani

Faculty of Medicine, Universiti Teknologi MARA (UiTM), Selangor, Malaysia

#### INTRODUCTION

The Lancet published this early report by Andrew Wakefield et al on February 28th, 1998; “12 children (mean age 6 years [range 3–10], 11 boys) were referred to a paediatric gastroenterology unit with history of normal development followed by loss of acquired skills, including language, together with diarrhoea and abdominal pain.... Onset of behavioural symptoms was associated, by the parents, with measles, mumps, and rubella vaccination in eight of the 12 children” [1].

This article flipped the concrete evidence-based success story of vaccination into an emotionally charged and debatable topic of the century. It was only after a decade of much larger studies which failed to replicate their findings that it became evident that there was no association between Measles, Mumps and Rubella (MMR) vaccination and autism. While it is well known that scientific investigations can be wrong but what is unacceptable here is the fraudulent research practice, in this case, the presentation of wrong data, and the lead author’s overwhelming undeclared conflict of interest. The aftermath could not be more devastating, Lancet withdrew the paper fully and the loss of his license to practice medicine in the UK in 2010.

#### THE WIDER IMPLICATION

The MMR-Autism link saga to the medical world meant that more research, time and money were poured to refute the study and also to expose the fraud but the repercussions however were not only confined to the medical profession. The greatest damage was the appalling tangential increase in vaccine refusal among parents worldwide which fuelled the measles outbreak

across the United Kingdom (UK), United States and Canada in the year 2008 and 2009. UK for example, saw a drop in vaccination rates from 87.4 percent to 79.9 percent in the year 2000-01 and 2003-04 respectively and not surprisingly, a dramatic increase in measles cases in the UK in the year 2007-08, which was equal to the combined total measles notifications for the past decade [2].

As the news coverage on the controversy intensified, and coupled with advancement of technology in the social media network, the public perception on vaccination has suddenly changed, the most successful health revolution in the 20th century is now at stake. Seemingly increasing public distrust and confusion over the safety of vaccination were echoed and mischievously elaborated geographically, reaching out to most of the third world countries including Malaysia causing the dreaded domino effect of declining immunisation rates in many countries including our own.

#### THE BLIGHT ON OUR SUCCESS

We began our free national immunisation programme for Diphtheria, Pertussis and Tetanus in 1958, the vaccination for Tuberculosis, Polio and Measles were gradually added into the immunization schedule between the years 1960 to 1980s. Malaysia has done very well since, based on the latest Millenium Development Goal (MDG) report in 2015, we have reached almost full coverage for one-year-old intake of the Measles, Mumps and Rubella (MMR) vaccine. The rate of intake of this vaccine was initially 70.1% in 1990, with massive improvement to 94.3% (2008) and 95.2% in 2013 [19], validated by a recent study in 2016 from a rural clinic in Sabah at 98.5% [29].

Lurking behind this success however is the rising trend of parents refusing to vaccinate their children. We now notice an increase in the number of vaccine refusal from 470 cases in 2013 to 648 in the following year and 1292 in 2015. Among the states in Malaysia, Kedah state recorded the highest number of vaccine rejection cases with steady rise from 239 cases in 2014 to 318 cases a year later.

Why is this happening? Data from the state of Kedah health statistics suggested that the major cause for the refusal was the concern regarding the vaccine contents and their religious permissibility (*halal*). This is supported by a cross-sectional study in 2013 done in Perak that showed the main reasons for parental immunisation refusal were preference to alternative treatment (75%), assumption that vaccines have no effect (37.5%) and apprehension on the vaccine contents (25%), other reasons included not being informed regarding vaccination from health practitioners, information from family members and media, religious influence, personal belief and long waiting time in the clinic [18]. In this study the refusal rate was 8 per 10,000 children per year and immunisation defaulter rate was 30 in 10,000 children per year. Vaccine refusal could also be caused by deferral which could be due to either ill infants or parents missing the schedule or appointments [17].

The number of vaccine preventable diseases has also showed steady increment for the past few years, in tandem with the decrement of immunisation rates. Measles cases in Malaysia has quadrupled from 195 cases in 2013 (6.6 cases per million population) to 794 cases up till September 2016 (34.7 per million population). This is certainly a blight on our success and it pushes us off track from the MDG target of global measles elimination by 2015.

### APPREHENSION OF VACCINE CONTENT

Certain chemicals are present as ingredients in the vaccines to ensure safety and effectiveness of the final products. These substances naturally exist in the environment and only become toxic if they reach or exceed a certain threshold.

Among chemicals used in vaccine preparation include thimerosal (mercury), which is an organic compound containing ethylmercury. Its primary role is

to prevent bacterial and fungal contamination and has been used as vaccine preservative since 1930's [30].

Virtually all vaccines are now mercury-free; and even if present its potential harmful effect is almost negligible as the chemical content in the vaccine is extremely low.

Aluminium is another compound used in vaccine preparation. It acts as an adjuvant to enhance the immune response to the vaccine antigen [30]. Exposure to aluminium from vaccines is well below the current minimum risk level of 2.0 mg/kg per day [30]. Interestingly, the content of aluminium is higher in breast milk compared to vaccines [31] as well in certain medications such as antacids [31].

However, another reason of apprehension that is being used as bone of contention by anti-vaccination campaigners is the permissibility (*halal*) of the vaccine contents.

### THE ISLAMIC VIEWPOINT

The objectives of Islamic law (*maqasid shariah*) are the preservation of five fundamental elements in a person; religion, life, lineage, intellect and property. Correspondingly, the maxim of Islamic law (*Qawaid al Fiqh*) adheres to the principle of avoiding harm, thus taking steps towards maintenance of health and this includes vaccine administrations to prevent serious and life-threatening illnesses among children are in accordance to these principles.

As stated earlier, the main religious consternation regarding vaccination among Muslims parents revolves around the issue of permissibility (*halal*) of the vaccine contents. In this regard, many scholars in Islamic Jurisprudence have in fact issued clear ruling (fatwa) regarding the permissibility of most vaccines used as part of national immunisation programmes worldwide, including Bacillus Calmette-Guerin (BCG), Hepatitis B, Diphtheria, Tetanus, Pertussis and Rubella vaccines [20-24].

Differences of opinion however does exist among the scholars regarding vaccines that have substances derived from pork, which are forbidden (*haram*) in Islam, being used during their manufacturing process. As an example, for the production of oral polio and rotavirus vaccines, trypsin enzyme of porcine origin is used during production to



dissociate the virus from cultured cells, but it is later removed through the process of microfiltration. The use of this substance however has led the Malaysian Fatwa Committee National Council of Islamic Religious Affairs in 2008 to issue a ruling that the use of Rotavirus vaccine is forbidden, other religious considerations by the council include the availability of an alternative trypsin source and the absence of an urgent state (*darurah*) for its use. But other opinion does exist which can be considered to be more in tune with the spirit of Islam that discourages complexity in performance of religious duties, the ruling from the European Council of Fatwa & Research in 2003 led by Yusuf al-Qardhawi. He concluded that the use of oral polio vaccine was permissible based on the following reasons; the negligible amount of trypsin used in the vaccine preparation, the fact that trypsin is filtered and thus not detectable in the final vaccine, and finally what is forbidden (*haram*) is made permissible in the state of necessity. As a result of this ruling, many Muslim countries such as Saudi Arabia, Bahrain, Yemen, Qatar, Iraq, Morocco, Sudan and Pakistan [20] have incorporated Rotavirus vaccine that uses porcine trypsin in their national immunisation programmes.

## THE WAY FORWARD

This requires efforts by all relevant stakeholders, government and non-government, to reverse the trend we see locally as well as worldwide. One great stride forward was the WHO approved Global Vaccine Action Plan, a framework to prevent millions of deaths by 2020 through more equitable access to existing vaccines for all peoples in all communities [30]. The aims here are to strengthen routine immunisation to meet vaccination coverage target, accelerate control of vaccine-preventable diseases as well spur research for development of new and improved vaccines [30].

Healthcare providers are undoubtedly the front liners in educating the parents and clarifying any doubts which may prohibit vaccine adherence among them. We know that counseling parents with clear information about the risks and benefits of vaccines, and taking advantage of clinical consultation visits for explanation of immunisation are among the most effective strategies suggested to achieve this [31]. In Malaysia, forums and educational talks to the general

public are actively organised by the Malaysian Ministry of Health and other non-governmental organisations to reach for these parents at all levels and localities. Besides that, social media is also very effective and is a borderless educational platform to reach the community.

Finally, the history of vaccination had been a great success story of the last century, Measles vaccination alone has been estimated to have helped save 17.1 million lives in the year 2000 [27]. Lack of knowledge on the issue compounded with contradictory information in social media have led to the disruption of herd immunity that previously had been the gate keeper in protecting our children from vaccine-preventable disease. We must do all we can to ensure it remains a success.

## REFERENCES

1. Wakefield AJ, Murch SH, Anthony A, Linnell J, Casson DM, Malik M, Berelowitz M, Dhillon AP, Thomson MA, Harvey P, Valentine A. Retracted: Ileal-lymphoid-nodular hyperplasia, non-specific colitis, and pervasive developmental disorder in children. *Lancet*. 1998; 351(9103): 637–41.
2. Thompson G. Measles and MMR statistics. House of commons, social and general statistics section. SN/SG/2581. 10 Sept 2009.
3. DeStefano F, Chen RT. Negative association between MMR and autism. *Lancet*. 1999; 353(9169): 1987–8.
4. Taylor B, Miller E, Farrington CP, Petropoulos MC, Favot-Mayaud I, Li J, Waight PA. Autism and measles, mumps, and rubella vaccine: No epidemiologic evidence for a causal association. *Lancet*. 1999; 353(9169): 2026–9.
5. Dales L, Hammer SJ, Smith NJ. Time trends in autism and in MMR immunization coverage in California. *JAMA*. 2001; 285(9): 1183–5.
6. Murch SH, Anthony A, Casson DH, Malik M, Berelowitz M, Dhillon AP, Thomson MA, Valentine A, Davies SE, Walker-Smith JA. Retraction of an interpretation. *Lancet*. 2004; 363(9411): 750.
7. Horton R. A statement by the editors of The Lancet. *Lancet*. 2004; 363(9411): 820–1.

8. Eggertson L. Lancet retracts 12-year-old article linking autism to MMR vaccines. *CMAJ*. 2010; 182(4): E199–200.
9. Wakefield AJ, Murch SH, Anthony A, Linnell J, Casson DM, Malik M, Berelowitz M, Dhillon AP, Thomson MA, Harvey P, Valentine A. Retraction-ileal-lymphoid-nodular hyperplasia, nonspecific colitis, and pervasive developmental disorder in children. *Lancet*. 2010; 375(9713): 445.
10. Godlee F. The fraud behind the MMR scare. *BMJ*. 2011; 342: d22.
11. Deer B. Wakefield's "autistic enterocolitis" under the microscope. *BMJ*. 2010; 340: c1127.
12. Deer B. How the case against the MMR vaccine was fixed. *BMJ*. 2011; 342: c5347.
13. Deer B. Secrets of the MMR scare. How the vaccine crisis was meant to make money. *BMJ*. 2011; 342: c5258.
14. Deer B. Secrets of the MMR scare. The Lancet's two days to bury bad news. *BMJ*. 2011; 342: c7001.
15. Deer B. Autism research: what makes an expert? *BMJ*. 2007; 334: 666–7.
16. Opel DJ, Diekema DS, Marcuse EK. Assuring research integrity in the wake of Wakefield. *BMJ*. 2011; 342: d2.
17. Othman N, Yusof M, Yanty AN. Immunisation status in hospitalized infants: Reasons for incomplete immunization. *MJMHS*. 2005; 1(1): 53–9.
18. Lim WY, Amar-Singh HS, Jeganathan N, Rahmat H, Mustafa NA, Mohd Yusof FS, Rahman R, Itam S, Chan CH, N-Julia MS. Exploring immunisation refusal by parents in the Malaysian context. *Cogent Medicine*. 2016; 3(1): 1142410.
19. Malaysia's Millennium Development Goals Report 2015. United Nations Malaysia.
20. Irwan MS. Pelalian: Kaedah pencegahan penyakit menurut perspektif Islam. *InfoSihat* June 2014. [http://www.infosihat.gov.my/infosihat/media/powpoint/S/pdf/01\\_pelalian\\_jakim.pdf](http://www.infosihat.gov.my/infosihat/media/powpoint/S/pdf/01_pelalian_jakim.pdf). Accessed 10 Sept 2016.
21. Shaikh MS. Vaccination: Personal Choice vs. Public Interest. Institute of Islamic Understanding Malaysia 2016. <http://www.iais.org.my>. Accessed 10 Sept 2016.
22. Qardhawi Y. What is the Islamic point of view regarding vaccinating children against specific diseases? *Islamopedia Online* 2010. <http://www.islamopediaonline.org/fatwa/what-islamic-point-view-regarding-vaccinating-children-against-specific-diseases>. Accessed 12 Sept 2016.
23. Grabenstein JD. What the World's religions teach, applied to vaccines and immune globulins. *Vaccine*. 2013; 31(16): 2011–23.
24. Nordin MM, Ismail SA, Chan LJ. The Islamic Perspectives of Immunisation. In: *Immunisation controversies. What you really need to know*. Malaysia: Brightside Solution Sdn Bhd. 2015; 62-104.
25. Department of Statistics, Ministry of Health Malaysia.
26. Amira M, Atilia A, Syahirah F, Ahmad Z, Rashidah M, Masri M. Perception of parents towards vaccination in children. Unpublished data.
27. Seth B, Margaret C, Christopher E, Anthony F, Anthony L, Joy P. Global vaccine action plan 2011-2020. [http://www.who.int/immunization/global\\_vaccine\\_action\\_plan](http://www.who.int/immunization/global_vaccine_action_plan). Accessed 15 Oct 2016.
28. Keeton VF, Chen AK. Immunization updates and challenges. *Curr Opin Pediatr*. 2010; 22(2): 234-40.
29. Barbara E Eldred, Angela J Dean, Treasure M McGuire and Allan L Nash. Vaccine components and constituents: responding to consumer concerns. *Med J Aust* 2006; 184 (4): 170-5.
30. Agency for Toxic Substances and Disease Registry. Toxicological profile for aluminium. Atlanta: US Department of Health and Human Services, Public Health Service, 2008. <http://www.atsdr.cdc.gov/toxprofiles/tp22.pdf>
31. National Health and Medical Research Council. *The Australian Immunisation Handbook*. 8th ed. Canberra: NHMRC. 2003. <http://immunise.health.gov.au/handbook.htm>

### Medical Education in Malaysia: The Evolving Curriculum (Part 1)

Rokiah Ismail

Faculty of Medicine, Universiti Teknologi MARA (UiTM), Selangor, Malaysia

#### THE TRADITIONAL CURRICULUM

Medical education in Malaysia has evolved in the past 50 years since independence. This paper highlights the various stages of curriculum development that were made to meet the needs of the developing country.

Malaya was under British rule between the 18<sup>th</sup> and the 20<sup>th</sup> Centuries. British Malaya as it was then known comprised of the Peninsular States and the Straits Settlements of Penang, Malacca and Singapore. Western medicine was introduced to the then Malaya in Singapore, with the setting up of the first medical school in 1907. It was called the Government Medical School and later became known as the King George VII College of Medicine in Singapore. In 1910 the first batch of seven male Medical graduates received their Licentiate in Medicine and Surgery (LMS) [1].

In 1949, the University of Malaya was established, based in Singapore, with a branch set up in Kuala Lumpur in 1959. In 1961, both governments of Singapore and Malaya agreed and passed legislation in Parliament to make the Kuala Lumpur Campus an autonomous body known as the University of Malaya; with its own medical school and teaching hospital. Thus in 1962, the government approved the setting up of the University of Malaya Medical Faculty, together with its teaching hospital, in the Klang Valley. The Faculty became fully functional in 1964 with the first intake of 64 medical students. After the hospital was built, the whole complex was named the “University of Malaya Medical Centre (UMMC)”, with facilities for undergraduate medical teaching, hospital services, the nursing school and other ancillary services put in place [2].

Professor Thumboo John Danaraj who was then Professor of Medicine in the Medical Faculty at

the University of Singapore, was appointed as Founding Dean of the Medical Faculty, University of Malaya in Kuala Lumpur [3]. With his appointment, the process of “head hunting” and appointment of academic staff began together with the selection of potential students for the first academic session.

It was mandatory that the Faculty get relevant and competent professionals to start the ball rolling. These medical academicians came from different parts of the globe, including Sri Lanka, Canada, Singapore and the UK (Figure 1).



**Figure 1** The founding teachers: Faculty of Medicine, University of Malaya, 1965 (Courtesy of the late Prof. TJ Danaraj).

With these academicians on board, the toiling of planning and designing of the medical curriculum started since the first batch of medical students was scheduled to enter the medical school in 1964.

Globally, the medical curriculum followed the traditional didactic teaching of basic sciences comprising of anatomy, physiology and biochemistry



in the first year of undergraduate medical course. In the second year, the subjects of pathology, pharmacology, medical microbiology and parasitology were introduced. This was interspersed with topics on communicable diseases and principles of social and preventive medicine (SPM). The thrust of the undergraduate curriculum then was in the various aspects of issues related to social and preventive medicine. This was deemed to be important because the newly formed Malaysia, for the most part, was still mostly rural.

Professor Danaraj, having had experience as an academician in Singapore, felt that the didactic teaching of “dry” basic science subjects may not be perceived as interesting and relevant by the medical students. Thus, early on in the undergraduate medical curriculum, he introduced the clinical correlation classes (CCC) with clinical cases brought to the auditorium to demonstrate the physical signs and correlate them with basic science topics that were learnt during the previous week (Figure 2). This made the preclinical students understand the importance of basic medical science subjects in order to be able to explain the symptoms and the development of physical signs when disease occurs.



**Figure 2** Clinical Auditorium, University of Malaya Medical Faculty. Clinical integration with patients starts in year 1 (CCC) (1967) [Courtesy of the late Prof. TJ Danaraj].

This was perhaps the earliest change in the curriculum to facilitate the teachers to think about possibilities of making basic science “dry topics” more interesting to the students. This gradual introduction of clinical medicine into basic science “preclinical years” and *vice versa* in the clinical years was perhaps the beginning of integrated teaching and the evolution of the undergraduate medical curriculum in Malaysia in the late 1980’s.


The clinical years begin from years 3 to 5 with rotations in general medicine, surgery, paediatrics orthopedics and obstetrics and gynaecology. In the clinical years, the integration of basic sciences in the form of clinico-pathological case (CPCs) discussions in the final year, sets the stage for future developments in the undergraduate curriculum. The clinical postings and the CPCs were meant to expose the students to develop their critical and analytical thinking skills during their clinical clerkships. Thus, learning to make reasonable diagnoses based on patho-physiological processes that had occurred, with minimal investigative procedures. This was meant to prepare them for their general medical service as medical officers in the rural areas, and also providing them with the basics for future career development.

## **NATIONAL STRATEGIES TO IMPROVE HEALTH CARE FOR THE POPULATION**

Let us now look at the needs of the country then, and how the medical schools were tasked by the Government to contribute towards improving the health services in the then rural Malaysian society.

During the British Administration of the then Malaya, the legacy left behind by the British was a network of health services that extended to the really remote parts of Peninsular Malaya [4], as depicted in Figure 3.

When Malaya had her independence in 1957, the health programs were somewhat coordinated although there was a gross deficiency of doctors to run the district hospitals and the general hospitals. Healthcare then was provided at best by the hospital assistants (now known as medical assistants, MA).



**Figure 3** Network of Government run Health Care Services (Legacy from the British Rule of Malaya). a) Rural health unit. b) Replica of the original photo (Courtesy of the late Prof. TJ Danaraj).

During this period the teaching of medicine closely followed the British medical education system that was practiced in the UK. In those early days, the teaching of medicine was by apprenticeship with some knowledge of basic sciences to explain the symptoms.

Then came the didactic (traditional); and scientific discipline model. This preceptor-ship had advantages especially when there was as yet no formal structured curriculum mapping. To this day, clinical mentoring and preceptor-ship is practiced to some extent in the clinical ward rounds with bedside teaching. The concept of mentoring and development of clinical acumen was very apt in clinical practice; both during the undergraduate days and continues in the world of medical academia to this day. This is an art that is slowly dying with the advent of investigative medical practices.

While doctors have to know how to use modern investigative tools, clinical acumen is still required, to be able to make reasonable diagnosis and institute treatment; to be able to determine what investigations are appropriate and when referrals are necessary. This is so because government-sponsored medical graduates face compulsory service that may be in rural areas where there is scarce advanced investigative tools to aid them in making the diagnosis.

In the 1990's with the inevitable trend of producing more specialists, it was deemed necessary for the Ministry of Health to ensure that there will be enough primary care providers and general physicians who would approach patients in a holistic manner.

This was tasked to the universities to take the lead to develop programs to train medical officers as generalists and family and primary care physicians.

Medical schools in Malaysia, in developing their medical curriculum, need to address these issues and tailor-make the curriculum to suit the healthcare needs of the country.

**To be continued in Part 2: The Blended Curriculum**

**REFERENCES**

1. Rashid FA. Biography of Malay Doctors 1900-1957 Malaya and Singapore.
2. <http://www.ummc.edu.my/introduction.asp>. Accessed 24 Nov 2016.
3. [https://en.wikipedia.org/wiki/University\\_Malaya\\_Medical\\_Centre](https://en.wikipedia.org/wiki/University_Malaya_Medical_Centre). Accessed 24 Nov 2016.
4. Malaysian Health System Review 2013. Health systems in transition. Asia Pacific observatory on health systems and policies. Ministry of Health, Malaysia. 2013; 1(3).

### Detection of Sarcocyst in Meat from Local Markets in Selangor: Highlighting the Use of Crude Sample in PCR

Azdayanti Muslim, Putri Shafinaz Sharudin, Atiqah Yunus, Norhabsah Omar, Alieya Zakaria, Norshafiqah Mohamad

Faculty of Medicine, Universiti Teknologi MARA, Selangor, Malaysia

#### Received

25<sup>th</sup> February 2016

#### Received in revised form

19<sup>th</sup> August 2016

#### Accepted

19<sup>th</sup> September 2016

#### Corresponding author:

**Azdayanti Muslim,**  
Parasitology Discipline,  
Faculty of Medicine,  
Sungai Buloh Campus,  
Universiti Teknologi MARA (UiTM),  
Sungai Buloh,  
47000 Selangor,  
Malaysia.  
Tel: +6012-574 0014,  
Fax: +6 03-6126 7428  
Email: azdayanti@salam.uitm.edu.my

#### ABSTRACT

**Introduction:** *Sarcocystis* spp. are obligate intracellular protozoan parasites which cause meat-borne parasitic disease. In Malaysia, sarcocystosis is seen as a potential emerging food-borne zoonosis after a series of large outbreak of human infections. Humans acquire infection either by ingestion of cyst in raw or undercooked infected meat or from sporocysts in contaminated food and water. The goal of this study is to identify the presence of *sarcocystis* parasites in meat of cattle, buffaloes, sheep and goats collected from local markets in Selangor, Malaysia. **Methods:** A total of 64 skeletal muscles samples (57 cattle, 2 buffaloes, 4 goats and 1 sheep) were collected from local markets. The samples were cut randomly into three pieces, squashed firmly between two glass slides and then examined microscopically for the presence of cysts. **Results:** Three samples of meat (4.69 %) from cattle (1), buffalo (1) and sheep (1) were found to be positive for cysts. The cysts were confirmed by PCR as *sarcocystis* sp. **Conclusion:** The results showed low prevalence of *Sarcocystis* infection in meat collected from local markets. However, since there is a transmission among the livestock, extra precaution should be taken in consideration to prevent the spreading of sarcocystosis from animals to human.

**KEYWORDS:** Sarcocystosis, meat, local markets, Selangor

## INTRODUCTION

*Sarcocystis* spp. are obligate intracellular protozoan parasite which can cause meat-borne parasitic disease alongside with other parasites, *Toxoplasma gondii*, *Taenia* spp. and *Trichinella* spp [1]. This protozoan needs two hosts; definitive (carnivores & omnivores) and intermediate host (herbivores) to complete its life cycle. Humans can serve as both definitive and intermediate host. Therefore, there are two types of sarcocystosis inflicting man; intestinal and muscular sarcocystosis, depending on the type of species [2]. Decades ago, this parasitic infection received less attention. However, the interest on them has re-ignited especially within the past years after a series of large human outbreaks have been reported in Malaysia [3-7].

Intestinal sarcocystosis is acquired when humans eat raw or undercooked infected beef or pork with cyst stage of the parasite [1]. The infected person

may present with fever, nausea, abdominal pain, vomiting, diarrhoea and weight loss. However, the infection is often asymptomatic and often self-limited [8]. Studies in livestock in Malaysia have shown that *Sarcocysts* have been identified in cattle [9, 10], sheep [11] and water buffaloes [10, 12]. In addition, recent study has shown high *Sarcocystis* infection with 61.8 % prevalence rate in goat [13]. Most of the infected animals are asymptomatic and identified by incidental finding at necropsy. Nevertheless, some pathological changes such as fever, haemorrhages, low production of milk and meat, abortion and death have been recorded associated with the infection in livestock [2].

Diagnosis of sarcocystosis involves identification of muscular sarcocystosis in tissue samples and is usually made by detection of macroscopic or microscopic cysts in the skeletal muscle, heart, tongue, esophagus, and diaphragm. Squash method is a reliable and easy technique but should be in parallel with other methods such as

histological examination, and digestion technique. Recently molecular detection by PCR is used widely especially for the determination of the species [9, 13, 14].

Globally, lack of awareness about the importance of cooking meat properly and the habit of eating raw meat among some of the societies leads to an increasing prevalence of zoonotic sarcocystosis. In Malaysia, sarcocystosis is being increasingly recognized and consumption of meat is very common. Most of the studies in Malaysia were focusing more on imported or local livestock taken from slaughterhouse. Hence, the study was carried out to identify the presence of *Sarcocystis spp.* in meat from fresh local market in Selangor, hence providing some information regarding the status of sarcocystosis parasitic infections among local breed livestock.

**METHODS**

In this study, a total of 64 muscle samples were collected randomly from cattle (n=57), buffaloes (n=2), goats (n=4) and sheep (n=1). All samples were acquired from local fresh food market in the vicinity of Selangor from July-August 2015. Each sample, weighing approximately 250g, was immediately placed in plastic bag and transported to the parasitology laboratory at the Faculty of Medicine, Universiti Teknologi MARA (UiTM) and stored at 4°C before further examination. Each sample was cut into three small pieces and was squeezed firmly between two glass slides. The slides were observed under light microscope at x4 and x10 magnifications for the presence of microscopic cysts. Positive meat samples were cut into smaller pieces and were placed into the garlic crushing instrument for juice extraction according to Latif and his colleagues [15]. The juice was then collected into 1.5 ml Eppendorf tube and centrifuged at 5000 rpm for 10 minutes. After discarding the supernatant, the sediments were then smeared on glass slides and were dipped into absolute methanol for 5 minutes then in 1% Giemsa stain for 24 hours. Then the slides were examined under x100 magnification to detect the presence of bradyzoite.

PCR was carried directly using the remaining juice from the three positive meat samples according to manufacturer’s protocols for TOYOBO KOD FX

Neo from crude sample. An approximately 1µl juice from each sample was pipetted directly onto the prepared TOYOBO KOD FX Neo PCR master mix (total solution: 50 µl) for amplification of D2 region of the 18S rRNA of *Sarcocystis sp.* Forward primer (5’-GGAAGCGATTGGAACC -3’) and reverse primer (5’ CCTTGGTCCGTGTTTCA -3’) [16] were used. PCR was performed using Mastercycler Gradient (Eppendorf, Germany). Reaction mixture consisted of 200 pM of each primer, 25µl of 2x PCR buffer for KOD FX Neo, 10 µl dNTP, 1µl crude sample (juice) and topped with nuclease-free water to a volume of 50µl. The thermal profile for the amplification is shown in Table 1. PCR product was analysed by agarose gel electrophoresis, viewed and analyzed using GelDocXr +3 System (Bio-Rad Laboratories, USA) and MyImage Analysis Software (Thermo Scientific, USA). The expected size of the PCR product was 350bp-450 bp.

**Table 1** The thermal profile for the amplification of DNA using TOYOBO KOD FX Neo.

	Temperature	Time	Cycle X
<b>Initial denaturation</b>	94 °C	2 min	1
<b>Denaturation</b>	95 °C	30 s	} 40
<b>Annealing</b>	55 °C	30 s	
<b>Extension</b>	72 °C	1 min	
<b>Final cycle</b>	72 °C	10 min	1

**RESULTS**


Of 64 meat samples collected, three (3) samples were found to be positive (4.69 %) with microscopic cysts; 1 from cattle (1.75 %), 1 from buffalo and 1 sheep. Of the four (4) goat samples, none of them were found to be infected with *Sarcocystis* parasites. The image of the microscopic cyst is shown in Figure 1.




**Figure 1** Microscopic sarcocyst under light microscopy with x10 magnification



Banana shaped with thin cyst wall bradyzoite was seen in the sample of cattle (Figure 2). While a lot of metrocytes (immature bradyzoites) were found from the sample of buffalo and sheep. PCR confirmed the parasite as *Sarcocystis sp.* (Figure 3).



**Figure 2** Bradyzoites (banana shaped with nucleus) under light microscopy x100 magnification



**Figure 3** DNA amplification result using TOYOBO KOD FX Neo from crude sample.

## DISCUSSION

Our study shows that, the infection rate of *Sarcocystis sp.* is very low with only 4.6 % infection rate in meat samples collected from local market. However, the result was contradictory with the previous studies in meat producing animals in Malaysia using the same method. The most recent study reported the prevalence rate of 86.0 %, 61.0 % and 28.0 % in sheep, goat and cattle respectively [13]. In 2013, 40.8 % cattle and

buffalo were found to be infected with sarcocystosis with high infection rate in the esophagus and skeletal muscle [10]. We believed that, the positive rate might be higher if the samples were taken from different parts as well.

In addition, bradyzoites were seen only in cattle sample. Bradyzoite is highly infective and can trigger the infection once consumed. Metrocytes were seen in another two samples and according to Dubey [8], it could be indicated as recently acquired infection. Interestingly, our study confirmed the *Sarcocystis* using TOYOBO KOD FX Neo. This maybe the first study which indicated the successfulness of using PCR from crude samples, without extraction of the DNA. Unfortunately, the PCR was negative in sheep sample, even though it was positive by microscopic examination. The result was not surprising since it could be due to the lack of DNA (only metrocytes were seen) or other technical errors. In 2015, Kutty and team in their comparison study suggested that more than one test were essential for the identification of muscular sarcocystosis [17]. Further comparison study especially on the sensitivity of TOYOBO KOD FX NEO with other molecular techniques need to be done in the future.

Unfortunately sequencing was not done in this study; hence the exact species could not be determined. Nevertheless, studies on meat producing animals in Malaysia and neighboring countries have shown that the predominant species infecting them were *S. bovicanis* [13] and *S. cruzi* [9] in cattle, *S. capracanis* in goat [13], *S. levinei* and *S. fusiformis* [18] in water buffalo and *S. ovicanis* in sheep [13].

## CONCLUSIONS

Although the infection rate of sarcocystosis was found low in this study, extra precautions must be taken to prevent the spreading of this infection as it is one of the emerging zoonotic diseases worldwide. Our community needs to be notified about the meat-borne zoonotic diseases, specifically sarcocystosis. Health education to the public is important to prevent its transmission. Consumption of a well-cooked meat is one of the best approaches to prevent sarcocystosis.

**Conflict of Interest**

Authors declare none.

**Acknowledgement**

We are grateful to the meat sellers at fresh local markets in Selangor for their kind assistance in collecting samples from cattle, goat, sheep and buffalo. Special appreciation goes to Prof Dr Bahaa Latef for his contribution in reviewing the paper. This study was part of Year 2 Elective Project, MBBS Program Faculty of Medicine UiTM (Sg. Buloh Campus).

**REFERENCES**

1. Dorny P, Praet N, Deckers N, Gabriel S. Emerging food-borne parasites. *Vet Parasitol.* 2009; 163(3): 196-206.
2. Fayer R, Esposito DH, Dubey JP. Human Infections with *Sarcocystis* Species. *Clin Microbiol Rev.* 2015; 28(2): 295-311.
3. Tappe D, Stich A, Langeheinecke A, von Sonnenburg F, Muntau B, Schafer J, Slesak G. Suspected new wave of muscular sarcocystosis in travellers returning from Tioman Island, Malaysia, May 2014. *Euro Surveill.* 2014; 19: 21.
4. Esposito DH, Stich A, Epelboin L, Malvy D, Han PV, Bottieau E, Da Silva A, Zanger P, Slesak G, van Genderen PJ, Rosenthal BM. Acute muscular sarcocystosis: an international investigation among ill travelers returning from Tioman Island, Malaysia, 2011–2012. *Clin Infect Dis.* 2014; 59(10):1401-10.
5. A Abubakar S, Teoh BT, Sam SS, Chang LY, Johari J, Hooi PS, Lakhbeer-Singh HK, Italiano CM, Omar SF, Wong KT, Ramli N, Tan CT. Outbreak of human infection with *Sarcocystis nesbitti*, Malaysia. 2012. *Emerg Infect Dis* 2013; 19(12): 1989-91.
6. Tappe D, Ernestus K, Rauthe S, Schoen C, Frosch M, Muller A, Stich A. Initial patient cluster and first positive biopsy findings in an outbreak of acute muscular *Sarcocystis*-like infection in travelers returning from Tioman island, Peninsular Malaysia, in 2011. *J Clin Microbiol.* 2013; 51(2): 725-6.
7. Esposito DH, Freedman D, Neumayr A, Parola P. Ongoing outbreak of an acute muscular *Sarcocystis*-like illness among travellers returning from Tioman Island, Malaysia, 2011–2012. *Euro Surveill.* 2012; 17(45).
8. Dubey JP. Foodborne and waterborne zoonotic sarcocystosis. *Food and Waterborne Parasitology.* 2015; 1(1): 2-11.
9. Ng YH, Fong MY, Subramaniam V, Shahari S, Lau YL. Short communication: Genetic variants of *Sarcocystis cruzi* in infected Malaysian cattle based on 18S rDNA. *Res Vet Sci* 2015; 103: 201-4.
10. Latif B, Vellayan S, Heo CC, Kutty MK, Omar E, Abdullah S. High prevalence of muscular sarcocystosis in cattle and water buffaloes from Selangor, Malaysia. *Trop biomed.* 2013; 30(4): 699-705.
11. Norlida O, Zarina M, Latif B, Abo R, Al-Sultan I. A histopathological appraisal of ovine *Sarcocystosis* in Kelantan. *J Adv Biomed Pathobiol Res.* 2012; 2(4): 137-42.
12. Dissanaikie AS, Kan SP. Studies on *Sarcocystis* in Malaysia. *Z Parasitenkd.* 1978; 55(2): 127-38.
13. Latif B, Kutty MK, Muslim A, Hussaini J, Omar E, Heo CC, Rossle NF, Abdullah S, Kamarudin MA, Zulkarnain MA. Light microscopy and molecular identification of *Sarcocystis* spp. in meat producing animals in Selangor, Malaysia. *Trop Biomed.* 2015; 32(3): 444-52.
14. Lau YL, Chang PY, Tan CT, Fong MY, Mahmud R, Wong KT. *Sarcocystis nesbitti* infection in human skeletal muscle: possible transmission from snakes. *Am J Trop Med Hyg.* 2014; 90(2): 361-4.
15. Latif B, Al-Delemi JK, Mohammed BS, Al-Bayati SM, Al-Amiry AM. Prevalence of *Sarcocystis* spp. in meat-producing animals in Iraq. *Vet Parasitol.* 1999; 84(1-2): 85-90.
16. Wünschmann A, Armien AG, Reed L, Gruber AD, Olias P. *Sarcocystis calchasi*-Associated Neurologic Disease in a Domestic Pigeon in North America. *Transbound Emerg Dis.* 2011. 58(6): 526-30.
17. Kutty MK, Latif B, Muslim A, Hussaini J, Daher AM, Heo CC, Abdullah S. Detection of sarcocystosis in goats in Malaysia by light microscopy, histology, and PCR. *Trop Anim Health Prod.* 2015; 47(4): 751-6.
18. Huong LT, Uggla A. *Sarcocystis dubeyi* n. sp. (Protozoa: Sarcocystidae) in the water buffalo (*Bubalis bubalis*).