UNIVERSITI TEKNOLOGI MARA

COPY NUMBER VARIATIONS OF 'ORANG ASLI' (NEGRITO) FROM PENINSULAR MALAYSIA

SITI SHUHADA MOKHTAR

+

Thesis submitted in fulfillment of the requirements for the degree of **Master of Science**

Faculty of Medicine

April 2015

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student Student I.D. No. Programme Faculty Thesis Tittle Siti Shuhada Binti Mokhtar
2010692186
Master of Science (MD 780)
Medicine
Copy Number Variations of 'Orang Asli'(Negrito) From Peninsular Malaysia

Suhalfolder

Signature of Student Date

: April 2015

ABSTRACT

Copy number variation (CNV) has been acknowledged as a major contributor to the human genome diversity. This variability covers approximately about 15% of the entire human genome. To date, it is reported that CNV plays an important role in the association of common and complex diseases and also in determining the phenotypes. Nevertheless, CNV data from diverse populations is still limited and not saturated especially population from South East Asia. This study represents the first investigation of CNV in the indigenous populations from Peninsular Malaysia, of particular interest, the Negrito. Ninety-seven Negrito samples were recruited, of which 50 unrelated Negrito samples were genotyped using the Affymetrix SNP 6.0 microarray. CNVs were then called by two independent algorithms namely, Genotyping Console and Nexus CNV. Subsequently the CNVs being called by both algorithms were considered stringent hence included to the subsequent analyses. A total of 643 stringent CNVs were identified, comprising 217 gains and 426 losses. These stringent CNVs were then matched with the publicly available datasets including DGV, HapMap3 and SGVP, and identified 57 putative novel and Negrito specific CNVs, consisting of 28 CNVRs. These included 15 gains and 13 losses. Analysis of gene ontology revealed that genes within these CNVs were enriched in the immune system (p < 0.001). In view of the small population size, relative isolation and semi-nomadic lifestyles of this community, it is speculated that these CNVs may be attributed to recent local adaptation of Negritos from Peninsular Malaysia. Nonetheless other factors such as genetic drift should not be ruled out. This study offers a preliminary effort at an extent to which rare variants shape risk of common disease. More saturated CNV map is needed and should be carried out through large-scale next generation sequencing.

TABLE OF CONTENTS

CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR'S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENT	v
TABLE OF CONTENTS	vi
LIST OF TABLES	ix
LIST OF FIGURES	xi
LIST OF ABBREVIATIONS	xiii
CHAPTER ONE: INTRODUCTION	
1.1 Background of Study	1
1.2 Problem Statement	1
1.2 Problem Statement	4
1.3 Significance of Study	4

1.4 Research Objectives	5

CHAPTER TWO: LITERATURE REVIEW

2.1	.1 Human Genome	
	2.1.1 The Human Genome Project	6
	2.1.2 The Human Genome Diversity Project (HGDP)	7
	2.1.3 The International HapMap Project	7
	2.1.4 The Singapore Genome Variation Project (SGVP)	8
	2.1.5 The 1000 Genome Project	9
2.2	Deoxyribonucleic acid (DNA)	10
2.3	Structural Variations in Human Genome	12
2.4	Copy Number Variations	14
2.5	Copy Number Variation Detection Method	17
	2.5.1 Fluorescence in situ hybridization (FISH)	17
	2.5.2 Array comparative genomic hybridization (Array CGH)	18
	2.5.3 Quantitative PCR (qPCR)	18

	2.5.4 Microarray Technology	19
2.6	Population Genetics	21
2.7	Orang Asli from Peninsular Malaysia	22
2.8	Negrito of Peninsular Malaysia	25

CHAPTER THREE: METHODOLOGY

3.1	Study Design		27
3.2	Ethics	Approval	29
3.3	Samp	le Recruitment	30
3.4	Extrac	ction of DNA from whole blood sample	32
3.5	Quant	itative Analysis - Concentration and purity measurement of	32
	Extra	cted DNA	
3.6	Quali	ative Analysis – Electrophoresis	33
3.7	DNA	Genotyping with Affymetrix SNP 6.0 Platform	33
	3.7.1	Digestion	36
	3.7.2	Ligation	39
	3.7.3	Polymerase Chain Reaction (PCR)-Stage A	42
	3.7.4	Polymerase Chain Reaction (PCR)- Stage B	44
	3.7.5	PCR Product Purification	44
	3.7.6	Fragmentation	45
	3.7.7	Fragmentation QC Gel Electrophoresis	47
	3.7.8	Labeling	47
	3.7.9	Hybridization	49
	3.7.10	Washing and Staining	51
	3.7.11	Scanning	52
3.8	Genot	yping Data Analysis	52
	3.8.1	Affymetrix Genotyping Console Version 4.1 Software	52
	3.8.2	Nexus CNV Version 7.0 Software	54
	3.8.3	Stringent and Population Specific CNVs Analysis	56
3.9	Real-	Fime Quantitative PCR	56
	3.9.1	Optimization of Primers by Polymerase Chain Reaction	58