

VOLUME 14 NO. 1

JUNE 2017

ISSN 1675-7009

SCIENTIFIC RESEARCH JOURNAL

Institute of Research Management and Innovation

Synthesis, Spectral Characterisation and Antimicrobial Properties of Cu(II) and Fe(II)
Complexes with Xanthone

Rabuyah Ni, Mohammad Isa Mohamadin, Vivien Jong Yi Mian

Bioadsorption of Multiple Heavy Metal Ions by *Rhizophora Apiculate* sp. and *Elaeis
Guineensis* sp.

**M.B. Nicodemus Ujih, Mohammad Isa Mohamadin, Millaa-Armila Asli,
Bebe Norlita Mohamed**

Microwave Assisted Synthesis and Characterisation of Trinuclear Zinc(II) Schiff Base
Complexes Derived from *m*-phenylenediamine and Salicylaldehyde

Karimah Kassim, Muhamad Azwan Hamali

Identification of Pathogenic Bacteria Isolated from Raw and After Sand Filtration Water at
Lubok Buntar Water Treatment Plant

**Nur Hafizah Zakaria, Husnul Azan Tajarudin, Mohd Sharizal Mohd Sapongi
Mohamad Fared Murshed**

Analysing Population Structure of *Elaeis Oleifera* Germplasm using Model-Based Approach
Programme STRUCTURE

**Wan Nurhayati Wan Hanafi, Farida Zuraina Mohd Yusof, Rajinder Singh,
Ahmad Kushairi Din, Rajanaidu Nookiah, Maizura Ithnin**

SCIENTIFIC RESEARCH JOURNAL

Chief Editor

Hamidah Mohd Saman
Universiti Teknologi MARA, Malaysia

Assistant Chief Editor

Yazmin Sahol Hamid
Universiti Teknologi MARA, Malaysia

International Editors

R. Rajakuperan, B.S.Abdur Rahman University, India
Park Hee-Kyung, Korea Advanced Institute of Science and Technology, Korea
Vasudeo Zambare, South Dakota School of Mines and Technology, USA
Greg Tan, University of Notre Dame, Australia
Pauline Rudd, National Institute for Bioprocessing Research & Training, Dublin, Ireland

Editorial Board

Nor Ashikin Mohamed Noor Khan, Universiti Teknologi MARA, Malaysia
Yahaya Ahmad, University of Malaya, Malaysia
Faredia Ahmad, Universiti Teknologi Malaysia, Malaysia
Abdul Rahman Mohd. Sam, Universiti Teknologi Malaysia, Malaysia
Mohd Nizam Ab Rahman, Universiti Kebangsaan Malaysia, Malaysia
Faieza Hj. Buyong, Universiti Teknologi MARA, Malaysia
Judith Gisip, Universiti Teknologi MARA, Malaysia
Ahmad Hussein Abdul Hamid, Universiti Teknologi MARA, Malaysia
Baljit Singh Bhathal Singh, Universiti Teknologi MARA, Malaysia
Alias Mohd. Saman, Universiti Teknologi MARA, Malaysia

Journal Administrators

Khairul Nurudin Ahnaf Khaini, Universiti Teknologi MARA, Malaysia
Nurul Iza Umat, Universiti Teknologi MARA, Malaysia

© UiTM Press, UiTM 2017

All rights reserved. No part of this publication may be reproduced, copied, stored in any retrieval system or transmitted in any form or by any means; electronic, mechanical, photocopying, recording or otherwise; without prior permission in writing from the Director of UiTM Press, Universiti Teknologi MARA, 40450 Shah Alam, Selangor Darul Ehsan, Malaysia. E-mail: penerbit@salam.uitm.edu.my

Scientific Research Journal is a journal by Institute of Research Management & Innovation (IRMI), Universiti Teknologi MARA, Bangunan Wawasan, Level 3, 40450 Shah Alam, Selangor Darul Ehsan, Malaysia. E-mail: irmiuitm@salam.uitm.edu.my

The views, opinions and technical recommendations expressed by the contributors and authors are entirely their own and do not necessarily reflect the views of the editors, the publisher and the university.

SCIENTIFIC RESEARCH JOURNAL

Institute of Research Management & Innovation (IRMI)

Vol. 14 No. 1

June 2017

ISSN 1675-7009

1. **Synthesis, Spectral Characterisation and Antimicrobial Properties of Cu(II) and Fe(II) Complexes with Xanthone** 1
Rabuyah Ni
Mohammad Isa Mohamadin
Vivien Jong Yi Mian

2. **Bioadsorption of Multiple Heavy Metal Ions by *Rhizophora Apiculate sp.* and *Elaeisis Guineensis sp.*** 15
M.B. Nicodemus Ujih
Mohammad Isa Mohamadin
Millaa-Armila Asli
Bebe Norlita Mohamed

3. **Microwave Assisted Synthesis and Characterisation of Trinuclear Zinc(II) Schiff Base Complexes Derived from *m*-phenylenediamine and Salicylaldehyde** 29
Karimah Kassim
Muhamad Azwan Hamali

4. **Identification of Pathogenic Bacteria Isolated from Raw and After Sand Filtration Water at Lubok Buntar Water Treatment Plant** 41
Nur Hafizah Zakaria
Husnul Azan Tajarudin
Mohd Sharizal Mohd Sapingi
Mohamad Fared Murshed
5. **Analysing Population Structure of *Elaeis Oleifera* Germplasm using Model-Based Approach Programme STRUCTURE** 53
Wan Nurhayati Wan Hanafi
Farida Zuraina Mohd Yusof
Rajinder Singh
Ahmad Kushairi Din
Rajanaidu Nookiah
Maizura Ithnin

Microwave Assisted Synthesis and Characterisation of Trinuclear Zinc(II) Schiff Base Complexes Derived from *m*-phenylenediamine and Salicylaldehyde

Karimah Kassim¹, Muhamad Azwan Hamali²

¹*Institute of Science, Universiti Teknologi MARA,
40450 Shah Alam, Selangor, Malaysia*

²*Faculty of Applied Sciences, Universiti Teknologi MARA,
40450 Shah Alam, Selangor, Malaysia*

E-mail: ¹karimah@salam.uitm.edu.my,

²azwan_hamali@yahoo.com

Accepted: 31 May 2017

Received: 26 January 2017

ABSTRACT

A series of Schiff base ligand, SALMPD, and its mono- and trinuclear Zn(II) metal complexes were synthesised from m-phenylenediamine and salicylaldehyde in alcoholic solution. The synthesis of ligand and mononuclear complex were synthesised using conventional condensation method, while the trinuclear complex was done using microwave-assisted synthesis method. The structure of each compound was elucidated by elemental analysis, infrared and ¹H NMR spectroscopy. The infrared spectrum of SALMPD shows a strong azomethine (C=N) band at 1621.62cm⁻¹, indicates the formation of the ligand. Upon complexation of the mononuclear complex, the C=N infrared band shifted and the disappearing of the phenolic hydrogen signal in ¹H NMR suggesting the chelation between Zinc(II) metal ion and ligand took place when azomethine and phenolic hydrogen deprotonated. The trinuclear complex, Zn₃(SALMPD) obtained was consist of two moieties of mononuclear Zn(SALMPD), which act as ligands that chelating to the third Zn(II) metal ion through oxygen atom due to the shifting of M-O infrared band from 575.12-540.53cm⁻¹, which serves as a coordination site for the metal ion.

Keywords: *schiff base, m-phenylenediamine, salicylaldehyde, zinc, multinuclear*

INTRODUCTION

Many studies on Schiff bases has been conducted extensively which considered to be a 'privileged ligands' due to its stability and well-designed structures, as well as their ease in synthesising [1]. Ligands consisting of nitrogen (N), oxygen (O) or sulphur (S) donor atom are found to be biologically actives. This contributes to a wide range of biological application such as antimicrobial [2], anticancer [3] and antifungal [4]. On top of that, these ligands coordinate with almost all metal ions, capable of stabilising metals in various oxidation states to form complexes [5].

Metal ions are able to coordinate to an aromatic diamine based ligand on their relative *ortho*, *meta* or *para* position directly [6]. This is due to the presence of O, N, N, O tetradentate donor atom which positioned at the two axial open site, providing four coordination site for the metal to provide support to the ligand [7].

Multinuclear complexes have attracted particular interest due to its capability to coordinate more than one metal ion. This feature is biologically important because the complexes synthetically resemble ranges of metalloprotein. Their usage in the biological modeling of the metal active sites of the metalloprotein picks the interest of modern chemist to synthesis and characterisation of polynuclear complexes [8-9]. In order to prepare multinuclear complexes, a simple metal ion complex is firstly synthesised. This simple ion complex will serve as ligands for another metal ion [10].

The conventional method of synthesising multinuclear complexes is time-consuming and thus, replacing the conventional method with microwave assisted synthesis could overcome this problem. Microwave assisted synthesis has been developed and been explored broadly by the researcher. The irradiation that comes from the microwave not only shorten the reaction time, it also increased yield and eco-friendly, due to minimum usage of solvent [11-12].

In this paper, mono- and trinuclear zinc(II) complex were synthesised from Schiff base ligand derived from *m*-phenylenediamine and salicylaldehyde. The structures were elucidated using elemental analyses, FTIR and NMR.

MATERIALS AND METHOD

Chemicals and Materials

M-phenylenediamine (MPD), salicylaldehyde, zinc(II) acetate dihydrate, and triethylamine used in this research were purchased from Sigma Aldrich, while the solvent such as methanol and ethanol were analytical reagent grade purchased from Merck. All chemicals were used without further purification.

Instrumental

To synthesise the proposed trinuclear Zn(II) complex, Anton Paar Monowave 400 microwave was used. Elemental analyses were conducted on Flash EA 110 Elemental Analyzer. Infrared spectra were recorded using in the wavelength region of 4000-450 cm^{-1} using Perkin Elmer 1600 Spectrometer. The proton NMR spectra for ligand solution and complexes dissolved in D_6 -DMSO and CDCl_3 respectively were obtained using Bruker Avance 300 MHz. The melting point of the ligand and complexes were determined in a one-end capillary tube using BÜCHI Melting Point B-545 apparatus.

Synthesis

The overall reaction for the syntheses Schiff base and its mono- and trinuclear zinc complexes is represented in Figure 1.

Synthesis of Ligand, SALMPD

The Schiff base ligands were prepared according to the published literature method [13]. *M*-phenylenediamine (5mmol, 0.54g) dissolved in 15mL MeOH was added dropwise to (10mmol, 1.22g) salicylaldehyde in 15mL MeOH solution. The mixture was refluxed for three hours with continuous stirring. The yellow precipitate was filtered off and washed with cold MeOH. The obtained ligand was then recrystallised using EtOH and dried in a vacuum desiccator.

Synthesis of Mononuclear Complex, Zn(SALMPD)

The synthesised SALMPD ligand (3mmol, 0.94g) was dissolved in EtOH, to which was added with equimolar ethanolic solution of $\text{Zn}(\text{CH}_3\text{COO})_2 \cdot 2\text{H}_2\text{O}$. The mixture was refluxed for three hours. The resulting yellowish precipitate was filtered and washes with cold EtOH and dried in a vacuum desiccator.

Synthesis of Trinuclear Complex, $\text{Zn}_3(\text{SALMPD})$

The $\text{Zn}_3(\text{SALMPD})$ complex was synthesised using the method as described by [14] with slight modification. The previously synthesised mononuclear complex (1mmol, 0.37g) was mixed equimolar with $\text{Zn}(\text{CH}_3\text{COO})_2 \cdot 2\text{H}_2\text{O}$ in 5mL EtOH with the addition drops of triethylamine. The microwave condition was set at speed of a 1000rpm and heat up to 160°C. The reaction took place for 15 minutes before the yellow precipitates were filtered and rinsed with cold ethanol. The product was dried in a vacuum desiccator.




Figure 1: The synthesis routes of SALMPD, Zn(SALMPD) and Zn₃(SALMPD)

RESULT AND DISCUSSION

Synthesis of Schiff Bases

SALMPD ligand was prepared through condensation of *m*-phenylenediamine and salicylaldehyde in EtOH, with the ratio of 2:1. The ligand, which is the precursor for the formation of mononuclear zinc complex, was checked using elemental analysis and the physical properties are shown in Table 1.

The elemental analysis of the ligand is confirmed to be similar as previously reported [15]. From Table 1, the corresponding data of Zn(SALMPD) and Zn₃(SALMPD) suggested that these complexes are dimers.

Table 1: Analytical data of the ligand, mono- and trinuclear Zn complex

| Comp. | Emp for. (molc. Mass) | Yield (%) | Melting point (°C) | Elemental analysis Calc. (Found) (%) | | |
|--------------------------|---|--------------|--------------------------|---|----------------|----------------|
| | | | | C | H | N |
| SALMPD | C ₂₀ H ₁₆ N ₂ O ₂ (316.36) | 93.45 | 110 | 75.93 (76.09) | 5.10 (5.07) | 8.86 (9.15) |
| Zn(SALMPD) | C ₂₀ H ₁₄ N ₂ O ₂ Zn (379.72) | 85.08 | >300 | 63.26 (60.39) | 3.72 (3.84) | 7.38 (5.07) |
| Zn ₃ (SALMPD) | C ₄₀ H ₂₈ N ₄ O ₄ Zn ₃ (820.80) | 67.25 | >300 | 58.53 (53.35) | 2.95 (3.30) | 6.83 (6.01) |

Infrared Spectroscopy

The important vibrational modes of the functional group for the ligand and its respective complexes are described in Table 2 and Figure 2. A strong absorption band at 1621.62cm^{-1} for ligand SALMPD is assigned to be the azomethine ($\text{C}=\text{N}$) functional groups. This band was shifted to a lower frequency in complex $\text{Zn}(\text{SALMPD})$ by approximately 20cm^{-1} . The shifting indicates that the coordination between the free ligand and metal ion through azomethine nitrogen was a success [8]. As for the tri-nuclear complex, the $\text{C}=\text{N}$ absorption band only show small shifting since there are no new atoms coordinates at the azomethine nitrogen.

Another strong band from the ligand was detected at 1275.28cm^{-1} , assigned to be the stretching vibration of the $\text{C}-\text{O}$ of the phenolic group. Similarly, the band shifted to a lower frequency at 1148.23cm^{-1} indicates the chelation of the metal ion to the ligand through oxygen atom of the phenolic group [16].

The shifting of $\text{C}=\text{N}$ and $\text{C}-\text{O}$ vibrational band in $\text{Zn}(\text{SALMPD})$ and $\text{Zn}_3(\text{SALMPD})$ leads to the formation of a new band in the low the wavelength region. The weak intensity band observed at 609.71cm^{-1} and 607.87cm^{-1} respectively were found to be the $\text{M}-\text{N}$ bond, while the vibrational band 575.12cm^{-1} from $\text{Zn}(\text{SALMPD})$ attributes the present of $\text{M}-\text{O}$ stretching bond. As for complex $\text{Zn}_3(\text{SALMPD})$, the $\text{M}-\text{O}$ vibrational band was shifted to 540.53cm^{-1} , confirming the coordination of the third metal ion with another two mononuclear complex $\text{Zn}(\text{SALMPD})$.

Table 2: Infrared spectral data of ligand, mono- and trinuclear Zn complex

| Comp. | Assignments (cm^{-1}) | | | | |
|------------------------------|----------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | $\nu(\text{C}=\text{N})$ | $\nu(\text{C}=\text{C})$ | $\nu(\text{C}-\text{O})$ | $\nu(\text{M}-\text{N})$ | $\nu(\text{M}-\text{O})$ |
| SALMPD | 1621.62 | 1459.52 | 1275.28 | - | - |
| $\text{Zn}(\text{SALMPD})$ | 1606.43 | 1439.85 | 1148.23 | 609.71 | 575.12 |
| $\text{Zn}_3(\text{SALMPD})$ | 1604.95 | 1440.26 | 1147.27 | 607.87 | 540.53 |




Figure 2: FTIR spectra of free ligand (SALMPD), Zn(SALMPD) complex, Zn₃(SALMPD)

¹H NMR Spectroscopy

The ¹H NMR spectra of the free ligand, SALMPD and its mono- and trinuclear zinc complex were recorded in D₆-DMSO and CDCl₃ respectively and its chemical shifts are given in Table 3. The spectra of SALMPD ligand showed a single peak on the low field region at 13.14 ppm indicated the presence of phenolic hydroxyl proton. This peak is not visible in the mono- and trinuclear complex, Zn(SALMPD) and Zn₃(SALMPD) suggesting that chelation of the metal ion occurred when the hydroxyl deprotonated. The peak observed at 8.71 ppm in free ligand SALMPD is attributed to azomethine (HC=N) proton and as the shifting of the azomethine chemical shift in Zn(SALMPD) and Zn₃(SALMPD) towards upfield region suggested the complexation between ligand and metal ion.

Table 3: Chemical shifts (ppm) for ^1H NMR of ligand, mono- and trinuclear Zn complex

| Comp. | Chemical shifts, δ (ppm) | | |
|--------------------------|---------------------------------|------|-------------|
| | O-H | HC=N | Ar. protons |
| SALMPD | 13.14 | 8.71 | 7.50-6.98 |
| Zn(SALMPD) | - | 8.68 | 7.50-6.61 |
| Zn ₃ (SALMPD) | - | 8.67 | 7.44-6.61 |

CONCLUSION

A series of mono- and trinuclear zinc complexes derived from *m*-phenylenediamine and salicylaldehyde has been successfully synthesised and the structure of these compounds has been elucidated using elemental analysis, melting point, infrared and ^1H NMR spectroscopy. More structural data such as magnetic susceptibility, molar conductivity, and thermogravimetric analysis are required to determine the compound's magnetism, geometrical shape, as well as to determine the presence of counter-ion in the complexes. Future work on anticancer activity will be studied in vitro using cell line. Previous studies hypothesised that complexation of Schiff base ligand enhances the biological properties of the free ligand while decreasing the toxicity effect from the metal ion. Other than that, microwave irradiation technique provided a quick, clean, and effective synthesis.

ACKNOWLEDGEMENT

The authors gratefully acknowledge the help of Universiti Teknologi MARA (UiTM) in providing the facilities and the Malaysian Ministry of Higher Education for the research grant fund (600-RMI/FRGS 5/3 (0068/2016).

REFERENCES

- [1] R. C. Maurya, H. Singh, A. Pandey, and T. Singh, 2001. Metal Chelates of Bioinorganic and Catalytic Relevance: Synthesis, Magnetic and Spectral Studies of Some Mononuclear and Binuclear Oxovanadium (V) and Dioxotungsten (VI) Complexes Involving Schiff Bases Derived from 4-Butyryl-3- Methyl-L-Phenyl-2-Pyrazolin-5-One and Certain Aromatic Amines, *Indian Journal of Chemistry*, Vol. 40(October), pp. 1053–1063.
- [2] O. Taheri, M. Behzad, A. Ghaffari, M. Kubicki, G. Dutkiewicz, A. Bezzatpour, H. Nazari, A. Khaleghian, A. Mohammadi, M. Salehi, 2014. Synthesis, Crystal Structures and Antibacterial Studies of Oxidovanadium(IV) Complexes of Salen-Type Schiff Base Ligands Derived from Meso-1, 2-Diphenyl-1, 2-Ethylenediamine, *Transit. Met. Chem.*, Vol. 39(2), pp. 253–259. <https://doi.org/10.1007/s11243-014-9798-9>
- [3] M. Damercheli, D. Dayyani, M. Behzad, B. Mehravi, and M. S. Ardestani, 2015. New Salen-Type Manganese(III) Schiff Base Complexes Derived from Meso-1,2-Diphenyl-1, 2-Ethylenediamine: in Vitro Anticancer Activity, Mechanism of Action, and Molecular Docking Studies, *J. Coord. Chem.*, Vol. 68(9), pp. 1500–1513. <https://doi.org/10.1080/00958972.2015.1027697>
- [4] M. M. Abo-Aly, A. M. Salem, M. A. Sayed, and A. A. Abdel Aziz, 2015. Spectroscopic and Structural Studies of The Schiff Base 3-Methoxy-N-Salicylidene-O-Amino Phenol Complexes With Some Transition Metal Ions and Their Antibacterial, Antifungal Activities, *Spectrochim. Acta - Part A Mol. Biomol. Spectrosc.*, Vol. 136(PB), pp. 993–1000. <https://doi.org/10.1016/j.saa.2014.09.122>
- [5] K. C. Gupta and A. K. Sutar, 2008. Catalytic Activities of Schiff Base Transition Metal Complexes, *Coord. Chem. Rev.*, Vol. 252(12–14), pp. 1420–1450. <https://doi.org/10.1016/j.ccr.2007.09.005>

- [6] R. Hernandez-Molina, A. Mederos, P. Gili, S. Dominguez, F. Lloret, J. Cano, M. Julve, C. Ruiza Perez, X. Solans, 1997. Dimer Species in Dimethyl Sulfoxide-Water (80:20 W/W) Solution of N,N'-Bis(Salicylideneimine)-M-Phenylenediamine (H₂sal-M-Phen) and Similar Schiff Bases With Cu^{II}, Ni^{II}, Co^{II} And Zn^{II}. Crystal Structure of [Co₂(sal-m-phen)₂]CHCl₃, *J. Chem. Soc., Dalt. Trans.*, (22), pp. 4327–4334. <https://doi.org/10.1039/A702151H>
- [7] P. G. Cozzi, 2014. Metal-Salen Schiff Base Complexes in Catalysis: Practical Aspects, *Chem. Soc. Rev.*, Vol. 33(7), pp. 410–421. <https://doi.org/10.1039/B307853C>
- [8] S. N. Abu Bakar, H. Bahron, and K. Kassim, 2010. Synthesis and Characterization of a Novel Schiff Base Derived from 2,4,6-Trimethyl-M-Phenylenediamine with O-Vanillin and Its Metal Complexes, *CSSR 2010 - 2010 International Conference on Science and Social Research*, pp. 463–466. <https://doi.org/10.1109/CSSR.2010.5773821>
- [9] D. Dey, Kaur, G., Ranjani, A., Gayathri, L., Chakraborty, P., Adhikary, J., Pasan, J., Dhanasekaran, D., R. Choudhury, A., A. Akbarsha, M., Kole, N., Biswas, B., 2014. A Trinuclear Zinc–Schiff Base Complex: Biocatalytic Activity and Cytotoxicity, *Eur. J. Inorg. Chem.*, Vol. 2014(21), pp. 3350–3358. <https://doi.org/10.1002/ejic.201402158>
- [10] A. J. Abdulghani and A. M. N. Khaleel, 2013. Preparation and Characterization of Di-, Tri-, and Tetranuclear Schiff Base Complexes Derived from Diamines and 3,4-Dihydroxybenzaldehyde, *Bioinorg. Chem. Appl.*, Vol. 2013. <https://doi.org/10.1155/2013/219356>
- [11] M. Chakraborty, S. Baweja, S. Bhagat, and T. Chundawat, 2012. Microwave Assisted Synthesis of Schiff Bases: A Green Approach, *Int. J. Chem. React. Eng.*, Vol. 10(1). <https://doi.org/10.1515/1542-6580.2973>
- [12] C. P. Gharu, 2014. Green and Efficient Microwave Assisted Synthesis of Schiff Bases and Hydroxyl Derivatives of 1, 3, 4-Thiadiazole Containing N-Methyl Piperazine Moiety and their Antimicrobial and Antioxidant Potential, *Chem Sci Trans.*, Vol. 3(4), pp. 1310–1317. <https://doi.org/10.7598/cst2014.904>

- [13] M. M. El-Ajaily, M. M. Abou-Krishna, A. M. Etorki, F. S. Alassbaly, and A. A. Maihub, 2013. Schiff base derived from phenylenediamine and salicylaldehyde as precursor techniques in coordination chemistry, *J. Chem. Pharm. Res.*, Vol. 5(12), pp. 933–938.
- [14] N. Karaboecek, S. Karaboecek, and F. Kormali, 2007. Mono-, Di-and Trinuclear Copper (II) Complexes of a Schiff Base Ligand, 2-{(E)-[(6-{{[(1E)-(2-hydroxyphenyl) methylene] amino} pyridin-2-yl) imino]-methyl} phenol, *Turkish J. Chem.*, Vol. 31(3), pp. 271–277.
- [15] M. M. El-Ajaily, M. M. Abou-Krishna, A. M. Etorki, F. S. Alassbaly, and A. A. Maihub, 2013. Schiff base derived from phenylenediamine and salicylaldehyde as precursor techniques in coordination chemistry, *J. Chem. Pharm. Res.*, Vol. 5(12), pp. 933–938.
- [16] P. Jeslin Kanaga Inba, B. Annaraj, S. Thalamuthu, and M. A. Neelakantan, 2013. Cu(II), Ni(II), and Zn(II) Complexes of Salan-Type Ligand Containing Ester Groups: Synthesis, Characterization, Electrochemical Properties, and In Vitro Biological Activities., *Bioinorg. Chem. Appl.*, Vol. 2013. <https://doi.org/10.1155/2013/439848>

GUIDELINES FOR SUBMISSION OF ARTICLES

The Scientific Research Journal (SRJ) (ISSN: 1675-7009) is an international refereed journal, jointly published by the Institute of Research Management and Innovation (IRMI) and the UiTM Press, Universiti Teknologi MARA (UiTM), Malaysia. This journal was launched in the hope of stimulating quality research projects in science and technology related areas. Researchers are strongly encouraged to use this journal as a platform for publishing and disseminating their research findings to the members of academia and the community at large.

The SRJ publishes research papers that address significant issues in the fields of science and technology including life science, natural science, health science and physical science. It provides a balanced presentation of articles and solicits contributions from the fields of: Engineering, Computer Science, Mathematical Sciences, Energy, Environmental Science, Materials Science, Physics, Chemistry, Astronomy, Pharmacy, Medicine and Applied Sciences.

Emphasis is place on direct and clearly understood communication, originality and scholarly merit. Only original papers will be accepted and copyright of published papers will be vested in the publisher.

Submitted manuscripts should be written in English with production-quality figures. Manuscripts should be typed using Times New Roman (point 12), single-line spacing and wide margins (2.54 cm left, right, top and bottom). The manuscript should include the title of the paper; the name, address, contact numbers and email(s) of the correspondence author; a short abstract of between 200 and 300 words, which clearly summaries the paper with 4-5 keywords. Submission should be limited to a maximum of 20 typed pages.

Tables should be included within the text where appropriate and must be numbered consecutively with Arabic numerals and have titles that precede the table. Similarly, all figures must be numbered and a detailed caption should be provided below each figure. Figures should be embedded within the text where appropriate. Glossy photographs when required

should be scanned to a suitable resolution (1200 dpi), which enables quality reproduction.

References should be numbered in ascending order and cited within square brackets, e.g. [1], [3-5], in the main body of the text. References included at the end of the manuscript are to be in the order that they appear in the main body of text. Some entry samples are as follows:

- [1] A. B. Author, 2000. *Title of Book*, ABC Press, Kuala Lumpur, Malaysia.
- [2] C. D. Author and E. F. Author, 1999. Title of Paper, *Journal Name*, Vol. 10, pp. 32-45.
- [3] G. H. Author, 2006. Title of the conference paper, in Proceedings of the 2000 IEEE International Symposium on Circuits and Systems, Geneva, Switzerland, pp. 100-105

Manuscripts submitted to the journal will be initially screened by the Chief Editor, to determine appropriateness. Only those manuscripts considered of a sufficiently high standard will proceed to undergo a double blind review.

Upon completion, the manuscripts will be passed to an editorial board member for appraisal of their value. Additionally, they will be reviewed by an expert in that discipline.

For further information, please contact:

Professor Dr. Hamidah Mohd Saman

Chief Editor

Scientific Research Journal

Institute of Research Management and Innovation

Universiti Teknologi MARA

40450 Shah Alam, Selangor,

MALAYSIA

E-mail: hamid929@salam.uitm.edu.my

Or e-mail SRJ's Secretariat: khairul2950@salam.uitm.edu.my



ISO 9001:2008 Certificate No. KLR 0404089

PENERBIT  **PRESS**
UNIVERSITI TEKNOLOGI MARA

ISSN 1675-7009

