ANTIMICROBIAL ACTIVITY OF Anacardium occidental LEAVES EXTRACT AGAINST SELECTED GRAM POSITIVE AND GRAM NEGATIVE BACTERIA

MARYAM NAJIHAH BINTI ZAINUDDIN

Final Year Project Report Submitted in
Partial Fulfillment of the Requirements for the
Degree of Bachelor of Science (Hons.) Biology
in the Faculty of Applied Sciences
Universiti Teknologi MARA

JANUARY 2017

ABSTRACT

ANTIMICROBIAL ACTIVITY OF Anacardium occidentale LEAVES EXTRACT AGAINST SELECTED GRAM POSITIVE AND GRAM NEGATIVE BACTERIA

Anacardium occidentale is the tropical tree that can be used as traditional medicine for treating variety of disease. The present study was carried out to determine the phytochemical compound and anitimicrobial activity of methanol and hexane of the leaves of A. occidentale. In phytochemical screening, there were five tests done to detect the presence of saponins, tannins, phenols, flavanoids and alkaloids. The results obtained demonstrate that in A. occidentale leaves contained the same phytochemical compounds which were saponins, tannins, phenols and flavanoids. The antimicrobial activity of leaves extract was determined by the disc diffusion method. The results obtained revealed that A occidentale leaves was potential against gram positive and gram negative bacteria. There were four concentration used (500 mg/ml, 250 mg/ml, 125 mg/ml and 65.5 mg/ml) for the leaves extract. The result showed that A. occidentale was effective against Bacillus subtilis, Klebsilla pneumoniae, Pseudomonas aeruginosa and Staphylococcus aureus. Concentration 500 mg/ml was the best concentration for A. occidentale leaves to inhibit the choosen bacteria. In conclusion, the leaves of A. occidentale was potential in inhibiting bacteria with significant value at p<0.05 in methanol extract.

TABLE OF CONTENTS

| В. | | | | | | PAGE |
|-------------------|------------|----------------------|---------------------------------------|------------|---------|---------|
| AC | KNOW | LEDGEMENTS | 94 37 | , , | H 4 | iii |
| TABLE OF CONTENTS | | | and an an an | | | iv |
| | | ABLES | | 9 II 8 | | vi |
| | | FIGURES | * * * * * * * * * * * * * * * * * * * | | 10 | vii |
| | | BBREVIATIONS | | | 18 | viii |
| ABSTRACT | | | * * 5 | | B B B | x |
| | STRAK | | u g u | | | xi |
| 8 200 | | | 8 8 | 8 | | |
| | | 18 A 9 | | | | |
| CH | APTER | R 1: INTRODUCTIO | N | | | |
| 1.1 | Backs | ground of Study | | 19 1901 | | 1 |
| 1.2 | Proble | em Statement | | | a na | 2 |
| 1.3 | Signif | ficance of the Study | | | at M | 3 |
| 1.4 | Objec | tive of the Study | * | | | 4 |
| | | | | | | a 5 |
| | | | 2 | | | |
| 25- | | R 2 : LITERATURE I | REVIEW | | | W 100 m |
| 2.1 | | ardium occidentale | | | | 5 |
| | 2.1.1 | Common type | | e e | | 7 |
| | 2.1.2 | Dwarf type | | | g. | 8 |
| | Extra | | | 9 | | 8 |
| 2.3 | - | chemical | | * | Bite | 10 |
| | | Alkaloids | | | | 11 |
| | | Flavanoids | | | | 12 |
| | 2.3.3 | Phenols | | | | 13 |
| 15 | 2.3.4 | Tannins | 40 | | | 13 |
| | 2.3.5 | Saponins | 2 2 10 g | | 348 | 14 |
| 2.4 | Bacte | | a 2 | 2 | N | 15 |
| | 2.4.1 | Staphylococcus aurer | | | 9 | 15 |
| | 2.4.2 | Pseudomonas aerugi | nosa | | | 17 |
| | 2.4.3 | Klebsilla pneumonia | e | E 8 | | 17 |
| | 2.4.4 | Bacillus subtilis | | | | 18 |
| | | | | | | |
| | A Marian - | A MERITORAL OF | O. T | | | |
| | | R 3 : METHODOLO | GY | * | 20 | |
| 3.1 | Mater | | ¥F | | H N | 1.0 |
| | 3.1.1 | | | | 1 | 19 |
| | 3.1.2 | | × | | | 19 |
| | 3.1.3 | Apparatus | | | 19 | 19 |

| 3.2 | Metho | ods | 20 |
|-----|--------|------------------------------------|----|
| | 3.2.1 | Sample collection | 20 |
| | 3.2.2 | Preparation of sample | 20 |
| | 3.2.3 | Extraction procedure | 21 |
| | 3.2.4 | Phytochemical | 21 |
| | | 3.2.4.1 Test for flavonoids | 21 |
| | | 3.2.4.2 Test for alkaloids | 22 |
| | | 3.2.4.3 Test for tannins | 22 |
| | | 3.2.4.4 Test for saponins | 22 |
| | | 3.2.3.5 Test for phenols | 22 |
| | 3.2.5 | | |
| | | 3.2.5.1 Nutrient broth (NB) | 23 |
| | | 3.2.5.2 Nutrient agar (NA) | 23 |
| | 3.2.6 | | 23 |
| | 3.2.7 | - | 24 |
| 3.3 | Data . | Analysis | 25 |
| | | | |
| | | | |
| CH | APTE | R 4: RESULTS AND DISCUSSION | |
| 4.1 | Perce | ntage Yield | 26 |
| 4.2 | | ochemical Screening | 28 |
| 4.3 | | nicrobial Activity | 32 |
| | | | |
| | | | |
| CH | APTE | R 5: CONCLUSION AND RECOMMENDATION | |
| 5.1 | Conc | lusion | 38 |
| 5.2 | Recor | mmendation | 39 |
| | | | 1 |
| | | | |
| CIT | ED RE | EFERENCES | 40 |
| API | PENDI | CES | 45 |
| CU | RRICU | JLUM VITAE | 54 |

LIST OF FIGURES

| FIGURE | TITLE | PAGE |
|--------|--|------|
| 4.2.1 | Phytochemical screening of methanol of A. occidentale leave extract: (a) saponins, (b) tannins, (c) phenols, (d) flavanoid and (e) alkaloid. | |
| 4.2.2 | Phytochemical screening of hexane of A. occidentale leave extract: (a) saponins, (b) tannins, (c) phenols, (d) flavanoid and (e) alkaloid. | 29 |
| 4.3.1 | Antimicrobial activity of methanol of A. Oocidentale leave extract against: (a) P. aeruginosa, (b) B. subtilis, (c) K. Pneumoniae, and (d) S. aureus | |
| 4.3.2 | Antimicrobial activity of hexane of A. occidentale leave extract against: (a) K. pneumoniae, (b) P. aeruginosa, (c) S. aureus and (d) B. subtilis | 33 |
| 4.3.3 | The antimicrobial activity of methanol leaves extract of A. occidentale against several bacteria species | 36 |
| 4.3.4 | The antimicrobial activity of hexane leaves extract of A. occidentale against several bacteria species | 37 |