

**ISOLATION AND IDENTIFICATION OF POTENTIAL CELLULOSE
BACTERIAL DEGRADER**

NURDINA FATIHAH BINTI AMINUDDIN

**Final Year Project Report Submitted in
Partial Fulfillment of the Requirements for the
Degree of Bachelor of Science (Hons.) Biology
in the Faculty of Applied Sciences
Universiti Teknologi MARA**

JANUARY 2017

ABSTRACT

ISOLATION AND IDENTIFICATION OF POTENTIAL CELLULOSE BACTERIAL DEGRADER

The cellulolytic bacteria can produce cellulase that degrade the cellulose and have the low possibility in causing environmental pollution and can be used in bioconversion technology to produce energy from biomass waste. The aims of this study are to isolate and screen the potential cellulose bacterial degrader. It is also to determine the optimum pH, temperature and incubation period for the growth of bacteria and to characterize the isolated bacteria by using biochemical test. The results of this study showed that 4 bacteria were isolated from soil of hot spring and able to growth on the carboxymethyl cellulose (CMC) agar. All the 4 bacteria show positive results which show the clearing zone around the colonies. However, only two of the bacteria which are ST2 and ST3 that shows high cellulase activity. Biochemical test revealed that both bacteria ST2 and ST3 were facultative anaerobic, gram variable and positive for citrate test. The optimum temperature, pH, incubation period for ST2 bacteria were 40°C, pH 10 and 6 days respectively. On the other hand, the optimum temperature, pH, incubation period for ST3 bacteria were 40°C, pH 8 and 6 days respectively. As a conclusion, ST2 and ST3 bacteria can be classified as mesophilic rod bacteria and have potential to degrade the cellulose.

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATIONS	viii
ABSTRACT	ix
ABSTRAK	x
CHAPTER 1: INTRODUCTION	
1.1 Background Study	1
1.2 Problem Statement	2
1.3 Significance of the Study	2
1.4 Objective of Study	3
CHAPTER 2: LITERATURE REVIEW	
2.1 Cellulose	4
2.2 Bacteria Cellulose Degradar	6
2.3 Physiological and Adaptive Thermophilic Bacteria	7
2.4 Cellulase	8
2.5 Application of Cellulase	9
2.5.1 Textile and Detergent Industries	10
2.5.2 Energy Industry	11
2.6 Advancing Method for Identification of Bacteria	12
2.6.1 Polymerase Chain Reaction and DNA Sequencing	12
2.6.2 OmniLog Identification System	13
CHAPTER 3: METHODOLOGY	
3.1 Material	14
3.1.1 Raw Material	14
3.1.2 Chemical	14
3.1.3 Apparatus	15
3.2 Methods	15
3.2.1 Preparation of Nutrient Agar	17
3.2.2 Preparation of Nutrient Broth	17
3.2.3 Preparation of Trace Element	17
3.2.4 Preparation of Mineral Salt Medium (MSM) Broth	17
3.2.5 Sample Collection	18
3.2.6 Isolation of Bacteria	18
3.2.7 Screening of Potential Cellulose Bacteria	18

3.2.8	DNS Method	19
3.2.9	Optimization for Bacterial Growth	19
3.2.9.1	pH	19
3.2.9.2	Temperature	20
3.2.9.3	Incubation Period	20
3.2.10	Morphological Observation	20
3.2.11	Biochemical Test	20
3.2.11.1	Methyl Red and Voges Proskauer	21
3.2.11.2	Indole Test	22
3.2.11.3	Catalase Test	22
3.2.11.4	Citrate Test	22
3.2.11.5	Glucose and Lactose Fermentation	23
3.2.11.6	Starch Hydrolysis Test	23
CHAPTER 4: RESULTS AND DISCUSSION		
4.1	Pre-screening for Cellulose Degrading Bacteria	24
4.2	The Cellulolytic Potential of Cellulase Potential Bacteria	26
4.3	Effect of pH for Optimization Bacterial Growth	27
4.4	Effect of Temperature for Bacterial Growth	28
4.5	The Effect of Incubation Periods for Bacterial Growth	30
4.6	Colony Characterization	31
4.7	Biochemical Characterization	33
CHAPTER 5: CONCLUSION AND RECOMMENDATIONS		35
CITED REFERENCES		37
APPENDICES		41
CURRICULUM VITAE		44

LIST OF TABLES

TABLE	TITLE	PAGE
4.1	The total cellulose activity produced by the bacteria	26
4.2	The growth of bacteria based on different pH	28
4.3	The growth of bacteria based on temperature	29
4.4	The growth of bacteria for incubation period	31
4.5	The morphological characterization of bacteria	32
4.6	Biochemical characterization of bacteria	34