

**CHEMICAL CONSTITUENTS FROM ETHYL ACETATE
EXTRACTS OF ROOT BARK OF *GARCINIA MANGOSTANA***

SHAHRUL AZWAN BIN ABDUL KHALIL

**Final Year Project Report Submitted in
Partial Fulfilment of the Requirement for the
Degree of Bachelor of Science (Hons.) Chemistry
in the Faculty of Applied Science
Universiti Teknologi MARA**

JANUARY 2017

ABSTRACT

CHEMICAL CONSTITUENTS FROM ETHYL ACETATE EXTRACTS OF ROOT BARK OF *GARCINIA MANGOSTANA*

Mangosteen fruit was widely used as medical purpose such as medicine to treat abdominal pain, diarrhea and infected wound. This was due to the presences of active compounds such as xanthones. Hence, root barks of *Garcinia mangostana* was chosen for further analysis. Detailed study on the cold extraction of ethyl acetate extract from root barks of *Garcinia mangostana* have resulted on the isolation of two phenolic compound, β -mangostin (**15**) and tovopyrifolin C (**46**). The structure of these compounds were identified and elucidated by using spectroscopy methods such as ^1H NMR, ^{13}C NMR, infrared (IR), EIMS, COSY, HMQC and HMBC.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	i
TABLE OF CONTENT	ii
LIST OF TABLES	iv
LIST OF FIGURES	v
LIST OF ABBREVIATIONS	vi
ABSTRACT	viii
ABSTRAK	ix
CHAPTER 1 INTRODUCTION	
1.1 Background of study	1
1.2 Botanical aspect of <i>Garcinia mangostana</i>	2
1.2.1 Family of <i>Clusiaceae</i>	2
1.2.2 The Genus of <i>Garcinia</i>	3
1.2.3 The Species of <i>Garcinia mangostana</i> L.	4
1.3 Problem statement	6
1.4 Significance of study	6
1.5 Objectives of study	6
CHAPTER 2 LITERATURE REVIEW	
2.1 Chemical constituents in <i>Garcinia mangostana</i>	7
2.2 Previous study on xanthenes	7
CHAPTER 3 METHODOLOGY	
3.1 Plant material	18
3.2 Previous study on xanthenes	18
3.2.1 Infrared Spectroscopy (IR)	18
3.2.2 Gas Chromatography-Mass Spectra (GC-MS)	18
3.2.3 Melting point	18
3.2.4 Nuclear Magnetic Resonance (NMR)	19
3.2.5 Ultra Violet Spectroscopy (UV)	19
3.3 Chromatographic method	19
3.3.1 Column chromatography	19
3.3.2 Thin Layer Chromatography (TLC)	19
3.4 Dyeing Reagent for TLC	20
3.4.1 Iron (III) Chloride solution	20
3.5 Extraction and isolation of chemical composition from root bark of <i>Garcinia mangostana</i>	21
3.5.1 Isolation of β -mangostin	21
3.5.2 Isolation of Tovopyrofolin C	23

CHAPTER 4 RESULTS AND DISCUSSIONS	
4.1 Isolation of chemical constituents from <i>Garcinia mangostana</i>	24
4.1.1 Characterization of β -mangostin	25
4.1.2 Characterization of Tovopyrifolin C	33
CHAPTER 5 CONCLUSION AND RECOMMENDATIONS	44
CITED REFERENCES	45
CURRICULUM VITAE	49

LIST OF FIGURES

Figures	Caption	Page
1.1	<i>G. mangostana</i> tree	5
1.2	<i>G. mangostana</i> roots	5
4.1	Compounds obtained from root bark of <i>Garcinia mangostana</i>	24
4.2	EIMS spectrum of β -mangostin (15)	28
4.3	IR spectrum of β -mangostin (15)	29
4.4	^1H NMR spectrum of β -mangostin (15) (400 MHz, CDCl_3)	30
4.5	^{13}C NMR spectrum of β -mangostin (15) (100 MHz, CDCl_3)	31
4.6	DEPT spectrum of β -mangostin (15) (100 MHz, CDCl_3)	32
4.7	EIMS spectrum of tovoxyrifolin C (46)	36
4.8	IR spectrum of tovoxyrifolin C (46)	37
4.9	^1H NMR spectrum of tovoxyrifolin C (46) (500 MHz, acetone-D)	38
4.10	COSY spectrum of tovoxyrifolin C (46) (500 MHz, acetone-D)	39
4.11	^{13}C NMR spectrum of tovoxyrifolin C (46) (125 MHz, acetone-D)	40
4.12	DEPT spectrum of tovoxyrifolin C (46) (125 MHz, acetone-D)	41
4.13	HMQC spectrum of tovoxyrifolin C (46)	42
4.14	HMBC spectrum of tovoxyrifolin C (46)	43