## PRELIMINARY SCREENING OF ALKALOID BY USING THIN LAYER CHROMATOGRAPHY (TLC) ANALYSIS FROM SEDIMENT OF Manihot esculenta (TAPIOCA)

**NURUL ATIQAH BINTI AZMI** 

Final Year Project Submitted in
Partial Fulfillment of the Requirements for the
Degree of Bachelor of Science (Hons.) Biology
In the Faculty of Applied Sciences
Universiti Teknologi MARA

**JANUARY 2019** 

This Final Year Project Report entitle "Preliminary Screening of Alkaloid by Using Thin Layer Chromatography (TLC) Analysis From Sediment Of Manihot esculenta (Tapioca)" was submitted by Nurul Atiqah binti Azmi, in partial fulfillment of the requirement for the Degree of Bachelor of Science (Hons.) Biology, in the Faculty of Applied Science and was approved by

Dr. Rosli bin Noormi
Supervisor
Faculty of Applied Science
Universiti Teknologi MARA
Cawangan Negeri Sembilan
Kampus Kuala Pilah
7200, Kuala Pilah, Negeri Sembilan

\_\_\_\_\_

Siti Norazura binti Jamal Project coordinator Faculty of Applied Science Universiti Teknologi MARA Cawangan Negeri Sembilan Kampus Kuala Pilah 7200, Kuala Pilah, Negeri Dr. Aslizah binti Md Aris Head of Biology School Faculty of Applied Science Universiti Teknologi MARA Cawangan Negeri Sembilan Kampus Kuala Pilah 7200, Kuala Pilah, Negeri Sembilan

Date:		

## **TABLE OF CONTENTS**

		PAGE
ACF	KNOWLEDGMENT	iii
	BLE OF CONTENT	iv
	Γ OF TABLES	vi
LIST OF PLATES LIST OF ABBREVIATIONS		vii ix
	TRAK	xi
CHA	APTER 1: INTRODUCTION	
1.1	Background Study	1
1.2		3
1.3	$\mathcal{E}$	4
1.4	Objectives of the Study	5
CHA	APTER 2: LITERATURE REVIEW	
2.1	Manihot esculenta (Tapioca)	
	2.2.1 Background of Tapioca	6
	2.2.2 Secondary metabolite in Tapioca	7
	2.2.3 Alkaloid in tapioca	9
2.2	Primary metabolite	10
2.3	Secondary metabolite	12
2.4	Alkaloid	13
2.5	Thin layer chromatography	15
2.6	Vanillin reagents	16
CHA	APTER 3: METHODOLOGY	
3.1	Materials	
	3.1.1 Raw materials	18
	3.1.2 Chemicals	18
	3.1.3 Apparatus	18
3.2	Methodology	13
	3.2.1 Preparation of sample	19
	3.2.2 Extraction of alkaloid	20
	3 2 3 Detection of alkaloid	21

CHA	APTER 4: RESULT AND DISCUSSION	
4.1	Identifying of Manihot esculenta (Tapioca)	24
4.2	Comparison between different ratio of solvent	
	(Chloroform:Hexane)	25
4.3	Comparison concentrations of Tapioca	29
4.4	Unknown compound of alkaloid detection	
	from sediment of Manihot esculenta	32
СНА	APTER 5: CONCLUSION AND DISCUSSION	36
CITI	ED REFERENCES	38
APP	ENDICES	44
CUR	RICULUM VITEA	47

## **ABSTRACT**

## PRELIMINARY SCREENING OF ALKALOID BY USING THIN LAYER CHROMATOGRAPHY (TLC) ANALYSIS FROM SEDIMENT OF Manihot esculenta (TAPIOCA)

Manihot esculenta or commonly known as tapioca is widely distributed around northern Brazil and it is the third largest source of carbohydrates in the tropics. It is indigenous to Brazil and is cultivated in most tropical parts of the Americas. After the arrival of the Spanish and Portuguese, the crop spread all over the tropical world especially in Africa, where it is now an important everyday staple and provide up to half of all calories consumed. The sediment of the tapioca always treated as the waste material from local factory in Jelai. In additional, this research will help people especially in rural area to produce new potential product in future from this sample such as snack because of its benefit for maintenance good health with the low cost of production. Moreover, it is also helps in reduce the potential of environment pollution due to abundance of this sediment. Thus, this study was aimed to detect the presence of alkaloid from sediment of Manihot esculenta that has been collected from local factory located in Jelai, Negeri Sembilan. The sample was collected, dried and extract by using 1:4 (Chlorofrom: Hexane) ratios. After that, they were sonicated by using sonicator for 15 minutes and centrifuged for 10 minutes with 6000 rpm at 4 °C. TLC method was used to run this sample with various ratios of chloroform and haxane which were 1:4, 4:1, 2:3, and 3:2. Difference concentration of tapioca was used 0.375mg/ml, 0.75mg/ml, 1.5mg/ml, 3mg/ml, 4.5mg/ml, 6mg/ml, 7.5mg/ml and 9mg/ml. Therefore, the best ratio showed was 1:4 ratio with 9mg/ml concentration was the best concentration after has been sprayed with vanillin reagent that expressed the clear band on the TLC plate. Beside, 8 unknown compound of alkaloid has been detected as well from 1:4 and 2:3 ratio. Unknown A, B, C, D, E, F, G, H, and I with Rf value of 0.15, 0.45, 0.78, 0.85, 0.08, 0.40, 0.53, 0.23 and 0.75 respectively. Last but not least, the results of this research investigation provide benefits for scientist and researcher for the identification of alkaloid and its concentration study in future by using for High Performance Liquid Chromatography (HPLC) analysis.