

**EXTRACTION AND PHYTOCHEMICAL SCREENING OF
MURRAYA KOENIGII LEAVES**

MUHAMMAD ZULFARIS BIN HARUN

**BACHELOR OF SCIENCE (Hons.) CHEMISTRY
FACULTY APPLIED SCIENCE
UNIVERSITI TEKNOLOGI MARA**

JANUARY 2020

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATION	viii
ABSTRACT	ix
ABSTRAK	x
Chapter 1	1
INTRODUCTION	1
1.1 Background of the study	1
1.2 Problem statement	3
1.3 Significance of Study	4
1.4 Objectives	4
Chapter 2 LITERATURE REVIEW	5
2.1 M.koenigii (L.) Spreng.	5
2.1.1 Taxonomic status and plant description	5
2.2 Preliminaries phytochemical screening	7
Chapter 3 METHODOLOGY	10
3.1 Leaves sample	10
3.2 Extraction and isolation of <i>Murraya koenigii</i>	10
3.3 Chromatographic Method	11
3.3.1 Column chromatography	11
3.3.2 Thin layer chromatography analysis	11
Chapter 4	13
Result and discussion	13
4.1 Extraction of <i>Murraya koenigii</i>	13
4.2 Preliminary phytochemical screening analysis	18

Chapter 5	23
CONCLUSION & RECOMENDDATION	23
CITED REFERENCES	24
CURRICULUM VITAE	27

LIST OF TABLES

Table	Caption	Page
4.1.1	Yield% from the soaking of pet ether, ethyl acetate and methanol	13
4.1.2	Yield% of each sample fraction from pet ether extraction	14
4.1.3	Yield% of each sample fraction from ethyl acetate extraction	15
4.1.4	Yield% of each sample fraction from methanol extraction	16
4.2.1	Phytochemical chemical analysis on column extracted sample of pet ether	18
4.2.1	Phytochemical chemical analysis on column extracted sample of ethyl acetate	19
4.2.1	Phytochemical chemical analysis on column extracted sample of methanol	20

ABSTRACT

EXTRACTION AND PHYTOCHEMICAL SCREENING OF *M.KOENIGII* LEAVES

Murraya Koenigii is a tropical tree originated from India and widely distributed across Southeast Asia and its leaves is a well-known traditional medicine. The genus of this species contains many of medicinal benefit such as anti-cancer, antimicrobial and anti-inflammatory. The leaves were soaked with petroleum ether, ethyl acetate and methanol respectively. Extraction were carried out using column chromatograph. After that, the fraction sample then were tested using phytochemical screening to determine the presences of alkaloid, flavonoid, terpenoid and tannin.