

UNIVERSITI TEKNOLOGI MARA

**ANTI-APOPTOTIC MECHANISM OF
URSODEOXYCHOLIC ACID (UDCA)
ON HYPOXIC CARDIOMYOCYTES**

NOORUL IZZATI BINTI HANAFI

MSc

February 2017

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student : Noorul Izzati binti Hanafi

Student I.D. No. : 2013685536

Programme : Master of Science Medicine (Biochemistry) – MD751

Faculty : Medicine

Thesis : Anti-Apoptotic Mechanism of Ursodeoxycholic Acid
(UDCA) on Hypoxic Cardiomyocytes

:

Signature of Student

Date : February 2017

ABSTRACT

Ursodeoxycholic acid (UDCA), the most hydrophilic bile acid is used as a therapeutic agent in liver related diseases and to eliminate hydrophobic bile acid induced apoptosis in liver. Recently, studies suggested the potential used of UDCA in treating heart related diseases. UDCA is reported cardioprotective against the development of ischemia. However, the mechanism of action in UDCA-cardioprotection is not clearly understood. Therefore, this study aimed to determine the anti-apoptotic mechanisms of UDCA on cardioprotection using an *in vitro* hypoxic model of neonatal rat cardiomyocytes. Rat heart from newborn (0-2 days old) was isolated for primary cell culture of cardiomyocytes. Hypoxia was induced by using CoCl_2 and hypoxic chamber. Cardiomyocytes were incubated with UDCA (pre-UDCA and post-UDCA) and co-incubated with FTY720 (S1P receptor agonist), PTX ($\text{G}\alpha_i$ inhibitor). The treated cardiomyocytes were subjected for proliferation assay (MTS assay), beating assessment assay, protein expression (aSMase and nSMase, Hif-1 α , caspase-3 and caspase-9, ERK and Akt), ROS generation assay and gene expression (*Hif-1 α* , *smpd1*, *smpd2*, *caspase-3*, *caspase-9*). The data were analyzed by using sample paired t-test and One-way ANOVA. MTS assay revealed that UDCA was not toxic to cardiomyocytes even at high concentration (250 μm). Results showed that CoCl_2 activates Hif-1 α expression, reduces cell viability; reduce beating rate, upregulates nSMase protein, increases ROS production, downregulates ERK and Akt protein expression. Interestingly, pre-UDCA treatment significantly abolished the negative effects of CoCl_2 on cell viability, beating rate, ROS production, ERK and Akt expression in cardiomyocytes. Treatment with PTX partially inhibits the protection of UDCA against CoCl_2 negatives effects on beating rate of cardiomyocytes. Meanwhile, FTY720 showed similarity with UDCA action in downregulating *smpd1* gene expression and upregulates *caspase-9* gene expression. In conclusion, the current data suggests that UDCA mechanism is mediated partially through $\text{G}\alpha_i$ -coupled receptor dependent and independent pathways in protecting cardiomyocytes against hypoxia. This study provides an insight of anti-apoptotic mechanism of UDCA on hypoxic cardiomyocytes.

ACKNOWLEDGEMENTS

Assalamualaikum w.b.t

First of all, I would like to thank to ALLAH S.W.T for His bless that enable me to complete this study without any problems. I am sincerely thankful to my supervisor Dr. Siti Hamimah binti Sheikh Abdul Kadir for her guidance for me to fulfill this study.

I also would like to express my appreciation to my co-supervisors, Dr. Julina Md Noor and Dr. Rosfaiizah Siran and also Dr. Nora Julianna Osman for their support and advice to complete the study.

Acknowledgement must also go towards Ministry of Higher Education (MOHE) and UiTM for funding this project through Fundamental Research Grant Scheme (FRGS) FRGS 5/3/FST (47/2011), and the Research Acculturation Grant Scheme (600-RMI/RAGS/2012/UITM/ST04/1) and (RAGS/1/2014/SKK01/UITM/1).

I would like also to take this opportunity to thank both of my parents, Mr. Hanafi Hassan and Mdm. Rosliala Yaakub and also my siblings for their trust and love to ensure that I completed this study.

I would like to express my gratitude to Institute of Medical Molecular and Biotechnology (IMMB), Faculty of Medicine, UiTM Sungai Buloh for the facilities provided throughout the experiments. Thanks to all IMMB staff for being helpful whenever I need help in the laboratory.

Last but not least, I would like to thank all my lab-mates for being supportive and helpful throughout my study. Thank you very much everyone.

TABLE OF CONTENT

	Page
CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR'S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENTS	v
TABLE OF CONTENT	vi
LIST OF TABLES	x
LIST OF FIGURES	xi
LIST OF PLATES	xiv
LIST OF SYMBOLS	xv
LIST OF ABBREVIATIONS	xvi
CHAPTER ONE INTRODUCTION	1
1.1 Research Background	1
1.2 Problem Statement	3
1.3 Scope of Study	3
1.4 Research Objectives	4
1.5 Research Question	4
1.6 Hypothesis	4
1.7 Significance of The Study	4
1.8 Limitation of The Study	4
CHAPTER TWO LITERATURE REVIEW	6
2.1 Hypoxia	6
2.1.1 Hypoxia as an Early Event of Ischemia Heart Disease	7
2.1.2 Hypoxia Signaling	8
2.2 Apoptosis	15
2.2.1 Reactive Oxygen Species (ROS)	16
2.2.2 Caspase Activity	16
2.2.3 Intracellular Kinases Enzymes	17
2.3 Bile Acid	18