

UNIVERSITI TEKNOLOGI MARA

**RELATIONSHIP BETWEEN TIMING OF THE
FIRST ZYGOTIC CLEAVAGE WITH
CYTOSKELETAL STRUCTURES AND AMINO
ACID METABOLIC PROFILES IN VITRIFIED
MOUSE EMBRYOS**

WAN HAFIZAH BINTI W. JUSOF

PhD

November 2016

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student : Wan Hafizah Binti W. Jusof
Student I.D. No. : 2010429698
Programme : Doctor of Philosophy (Medicine) – MD990
Faculty : Medicine
Thesis Title : Relationship between Timing of the First Zygotic
Cleavage, Cytoskeletal Structures and Amino Acid
Metabolic Profiles in Vitrified Mouse Embryos

Signature of Student :

Date : November 2016

ABSTRACT

Timing of the first zygotic cleavage has been used as a marker of embryo developmental competence and subsequent viability. Previous studies showed that embryos that cleaved early had higher developmental viability. However, the factors contributing to timing of the first zygotic cleavage are unknown. Energy production from mitochondria, nucleus and cytoskeletal organization might be some of the factors involved. Amino acid metabolic profiles might also relate with timing of the first zygotic cleavage as it has been reported to have significant relationships with embryo viability. Thus, the present study was designed to investigate the relationship between timing of the first zygotic cleavage, amino acid metabolic profiles, mitochondria, nucleus and cytoskeletal organization of mouse embryos with subsequent viability. Embryos were retrieved from superovulated ICR mice, 28 hours after hCG injection. At this point of time, 2-cell stage embryos were categorized as early-cleaving (EC), while zygotes with two pronuclei as late-cleaving (LC) embryos. Embryos were cultured overnight in M16 medium supplemented with 3% Bovine Serum Albumin (BSA) in a humidified carbon dioxide (CO₂) incubator. For Experiment 1, both EC and LC embryos were divided into control and treatment groups. For control group, 2-cell stage embryos were cultured until the blastocyst stage. For treatment group, embryos were vitrified by EFS40 or EFS20/40 method for 1 hour and warmed. The vitrified-warmed embryos were cultured until the blastocyst stage. The number of surviving embryos and their development to the blastocyst were observed and counted. For Experiment 2, 2-cell stage embryos were divided into control (non-vitrified) and treatment (vitrified) groups. Embryos in both control and treatment groups were fixed in 4% paraformaldehyde and immunostained to visualize the localization and intensities of mitochondria, actin, tubulin and nucleus. Finally, the embryos were mounted on slides and examined under a Confocal Laser Scanning Microscope. The structures intensity were analyzed by LAS-AF-Lite Software. For Experiment 3, EC and LC embryos were cultured individually in 4 µl drops of KSOM/AA medium supplemented with 3% BSA. The embryos were transferred every 24 hours to fresh 4 µl drops of KSOM/AA + 3% BSA until the blastocyst stage. The spent culture medium was analyzed by Ultra Performance Liquid Chromatography (UPLC) for amino acids metabolic profile. Results showed that non-vitrified and vitrified EC embryos had significantly higher developmental viability and higher cryosurvivability after vitrification by EFS40 and EFS20/40 method. Confocal analysis showed that non-vitrified and vitrified EC embryos had a significantly higher densities of mitochondria, actin and nuclear chromatin compared with non-vitrified and vitrified LC embryos, which appear to result in more efficient cell division, and therefore greater developmental competence. Amino acids metabolic profile showed that EC embryos had a significantly lower amino acids turnover compared to LC embryos in both Day-2 to Day-3 and Day-3 to Day-4 of cultures. These findings suggest that the higher developmental viability of EC embryos was significantly attributed by lower metabolic activity. In conclusion, timing of the first zygotic cleavage was associated with mitochondria, nucleus and cytoskeletal ultrastructure and amino acids metabolic profile that affect subsequent developmental viability and cryosurvival of embryos.

ACKNOWLEDGEMENT

The experience during my Ph.D journey is unforgettable. Taking advantage of this opportunity, I would like to thank those people who made this thesis possible.

First and foremost, I would like to express my sincerest appreciation to my main supervisor Professor Dr. Nor Ashikin Mohamed Noor Khan for the continuous support of my Ph.D study and research, for her patience, understanding, motivation, inspiration and immense knowledge. Her guidance helped me in all the time of research and thesis writing. I could not have imagined having a better or friendlier supervisor for my Ph.D study. Besides my main supervisor, I also would like to express my honest appreciation to my co-supervisor Professor Dr. Mohd Hamim Rajikin that always be there to help, guide and support me. He is very attentive, communicative and extremely helpful. I have learned a lot from him including critical thinking and academic writing skills. My deepest thanks should also go to my second co-supervisor Associate Professor Dr. Nuraliza Abdul Satar, for her encouragement and insightful comments.

Writing this thesis was not the lonely experience it could have been because of cherished friends who provided enthusiasm and empathy. I thank my fellow labmates in embryo group, Norhazlin Jusoh @ Mohd Yusof, Razif Dasiman and Fazirul Mustafa for the stimulating discussions we have had. The group has been a source of friendships as well as good advice and collaboration. My sincere thanks also go to Puan Nor Shahida Abdul Rahman and Puan Salina Othman, for helping me in confocal study. I am also grateful to Haji Ismail Muit from Research Instruments for generously sharing his time, knowledge and expertise in amino acids profiling analysis. He has played a major role in making me understand the concept and uses of UPLC system.

I gratefully acknowledge the funding sources that made my Ph.D. work possible. I was given a full time study leave by Universiti Kuala Lumpur - Royal College of Medicine Perak (UniKL-RCMP) for 4 years and my study fees was sponsored by Skim Tindakan Pensyarah fellowship from Majlis Amanah Rakyat (MARA). My work was also supported by the Fundamental Research Grant Scheme (FRGS) [600-RMI/ST/FRGS 5/3/SFT (71/2010)], Research Acculturation Grant Scheme [600-RMI/RAGS 5/3 (89/2013)], Cumulative Impact Factor Initiative [600-RMI/DANA 5/3/CIFI (135/2013)], the Ministry of Higher Education, Malaysia and the UiTM Excellence Fund [600-RMI/ST/DANA 5/3/Dst (335/2011)].

Last but not the least, I would like to thank the unconditional love and encouragement provided by my family served as a secure anchor during the hard and easy times; my lovely husband Muhammad Iqbal bin Samadi, my beloved sons and daughter Muhammad Irfan Haziq, Muhammad Isyraf Hakim, Muhammad Imran Harith and Nur Irdina Hana for their countless sacrifices they have made, for all their love and encouragement, full support and understanding, my parents Haji W. Jusof W. Muda and Hajah Hasnah Husain, for raising me with love and supporting me spiritually throughout my life, my brothers and sisters who always help me in many ways. Thank you very much.

TABLE OF CONTENTS

	Page
CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR'S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENT	v
TABLE OF CONTENTS	vi
LIST OF TABLES	xi
LIST OF FIGURES	xiii
LIST OF SYMBOLS	xvi
LIST OF ABBREVIATIONS	xvii
CHAPTER ONE: INTRODUCTION	1
1.1 General Introduction	1
1.2 Problem Statement	4
1.3 Objectives	4
1.4 Significance of Study	5
CHAPTER TWO: LITERATURE REVIEW	6
2.1 Preimplantation Embryo Development	6
2.1.1 Cell Cycle in Early Development	10
2.2 Embryo Metabolism	12
2.3 Embryo Assessment and Selection Methods	14
2.3.1 Morphological Grading	15
2.3.2 Timing of the First Zygotic Cleavage	21
2.3.3 Metabolomic Analysis	22
2.3.3.1 Carbohydrate metabolism	23
2.3.3.2 Oxygen consumption	24
2.3.3.3 Amino acids	24
2.4 Major Organelles in Embryonic Development	28