UNIVERSITI TEKNOLOGI MARA

IN VITRO ANTIPROLIFERATIVE ACTIVITY OF THE METHANOL EXTRACT OF *BAUHINIA PURPUREA* LEAVES

NURFATIHAH NASIHA BINTI BAHAGIA 2008401734

Dissertation submitted in partial fulfillment of the requirement for Bachelor of Pharmacy (Hons)

Faculty of Pharmacy

2012

ACKNOWLEDGEMENT

Alhamdulillah, first of all I would like to thank God as for His blessing, I was able to complete my final year project and thesis writing. Thank to all people who have helped and inspired me during completing this project. Foremost, I would like to express my sincere gratitude to my supervisor, Madam Noor Jannah Bt Yob for the continuous support, patience, motivation, enthusiasm, immense knowledge and providing me with an excellent atmosphere for doing research. Her guidance helped me in all the time of research and writing of this thesis.

My sincere thanks also go to Professor Dr Mohd Zaki bin Salleh and Dr Teh Lay Kek for giving their permission to use PROMISE Laboratory to complete this project. Not to be forgotten, I would like to thank postgraduate students Mr Mohd Salleh and Miss Erda Syerena Bt Rosli who were always willing to help and give their best suggestions. My research would not have been possible without their helps.

Besides, I would like to thank lab assistants and staffs of PROMISE Laboratory of UiTM Puncak Alam who giving me the necessary knowledge about equipments and chemicals in the laboratory. Without their guidance, I would not be able to complete this project smoothly. Finally thanks to my fellow friends, for their support and sharing their knowledge that they have. I am grateful having them as their encouragement strengthens my spirit. Last but not the least, I would like to thank my family members who were always supporting and encouraging me with their best wishes. My endless appreciation goes to my beloved father, Mr Bahagia bin Sidek and mother, Madam Zah binti Awi for supporting me spiritually throughout my life. They were always there cheering me up and stood by me through the good times and bad. I would never have been able to finish my dissertation without the guidance of my supervisor, lecturers, postgraduate students, lab assistants, help from friends, and support from my family members. Thank you all.

TABLE OF CONTENTS

		Page
TITI	LE PAGE	
APP	ROVAL	
ACKNOWLEDGEMENTS		ii
TABLE OF CONTENTS		iv
LIST OF TABLES		vii
LIST OF FIGURES		viii
LIST OF EQUATIONS		ix
LIST OF ABBREVIATIONS		X
ABSTRACT		xi
CHA	APTER ONE (INTRODUCTION)	
1.1	Background of the study	1
1.2	Statement of problem	3
1.3	Objectives	4
1.4	Hypothesis	4
CHA	APTER TWO (LITERATURE REVIEW)	
2.1	Cancer	5
	2.1.1 Liver cancer	7
	2.1.2 Breast cancer	8
	2.1.3 Colon cancer	8
2.2	Herbal remedies	10

ABSTRACT

Background: Nowadays, there is an increasing burden of cancer in developed countries. The most common cancers in the world are liver cancer, breast cancer and colon cancer. Previous finding and latest studies show that *Bauhinia purpurea* or in Malay known as 'pokok tapak kerbau' have potential inhibition property to human cancer cells lines due to the new compounds which are named Bauhiniastatin 1, 2, 3, and pacharin. The purpose of this study is to evaluate antiproliferative activities of Bauhinia purpurea leaves methanolic extract on cancer cells lines and not affect normal cells lines. This study used four types of cell lines consists of Hep G2 (human hepatocarcinoma cells lines) MCF-7 (human breast adenocarcinoma cell line), HCT 116 (colorectal carcinoma cells lines) and WRL 68 (normal cells lines). Methods: Cells were plated in 96-well plate and incubated in conditions of 95% O2, 5% CO2 at 37°C. Five different concentrations of Bauhinia purpurea leaves methanolic extract were tested on cells lines; 0.1, 1, 10, 100, 1000 ug/mL and 3-[4,5-dimethylthioazol-2-yl]-2-5-diphenyltetrazolium bromide (MTT) assay was used to measure cell viability. The effects of different concentrations on percentage of cell viability and calculation of median inhibitory concentration (IC50) were determined by plotting graph percentage of cell viability versus log concentration of Bauhinia purpurea leaves methanolic extract. Results: Bauhinia purpurea leaves methanolic extract has potential antiproliferative activity against Hep G2 cancer cells lines, MCF-7 cancer cells lines, HCT 116 cancer cells lines by producing median inhibitory concentration IC50 of 143.3 ug/mL, 167.7 ug/mL, 104.2 ug/mL respectively. Conclusion: Methanolic extract of Bauhinia purpurea leaves exhibit potential anticancer activity and can be used as preliminary study to do further research on antiproliferative activity of Bauhinia purpurea leaves.

Keyword: Bauhinia purpurea leaves, cancer, methanolic extract, antiproliferative