DIVERSITY OF FRESHWATER FISHES AT LOWER STREAM OF TUARAN RIVER, SABAH

NURUL NADZIRAH BINTI YUSMAN

Final Year Project Report Submitted in Partial Fulfillment of the Requirements for the Degree of Bachelor of Science (Hons.) Biology In the Faculty of Applied Sciences Universiti Teknologi MARA

University Teknology MARA

JANUARY 2016

TABLE OF CONTENTS

			PAGI	
ACKNOWLEDGEMENTS				
TABLE OF CONTENTS				
LIST	OF TA	BLES	vi viii x	
LIST	OF FIG	GURES		
LIST	OF AB	BREVIATIONS		
ABSTRACK				
ABS	TRAK		xii	
СНА	PTER 1	: INTRODUCTION		
1.1				
1.2		em Statement	4	
1.3	Signif	icance of the Study	5	
1.4	Object	tives of the Study	5	
СНА	PTER 2	: LITERATURE REVIEW		
2.1		gy of Fishes	6	
		General Morphology and Characteristics	7	
		Taxonomy and Classification	9	
		Distribution and Abundance	10	
2.2		water Fishes and Their Diversity	11	
2.3		water Fishes in Malaysia	12	
	2.3.1	Factors Affecting Fish Diversity and Distribution	13	
2.4	Previo	ous Studies on Freshwater Fish Species in Sabah	15	
~~~.				
		S: METHODOLOGY	17	
3.1	Study		17	
3.2			19	
		Raw Materials	19	
		Chemicals	21	
2.2	3.2.3	Apparatus	21 23	
3.3	Methodology		23	
	3.3.1	Fish sampling and capturing method	24	
	3.3.2	Samples collection Wester physical chamical parameters (WPP)	24	
	3.3.3	Water physico-chemical parameters (WPP)	25	
	3.3.4	Species preservation	25	
	3.3.5	Species identification	25	
	3.3.6	Specimens depositions	23	

## LIST OF TABLES

TABLE	TITLE	PAGE
1.1	Villagers with reported river usage	4
2.1	External features of fish and their functions	7
4.1	The characteristics of each station at the lower stream of Tuaran River	29
4.2	The water physico-chemicals parameters of water measured in the Tuaran River	32
4.3	List of fish species caught at three different stations	36
4.4	Morphometric measurements and meristic counts of <i>Ambassis</i> interrupta	41
4.5	Morphometric measurements and meristic counts of Arius maculatus	43
4.6	Morphometric measurements and meristic counts of Barbonymus gonionotus	45
4.7	Morphometric measurements and meristic counts of Caranx sexfasciatus	47
4.8	Morphometric measurements and meristic counts of Gerres abbreviatus	49
4.9	Morphometric measurements and meristic counts of Leiognathus equulus	51
4.10	Morphometric measurements and meristic counts of Liza subviridis	53
4.11	Morphometric measurements and meristic counts of Lobocheilus bo	55

## LIST OF FIGURES

FIGURE	TITLE	PAGE
2.1	Generalized outline of fish	7
3.1	Sampling stations and their coordinates at the lower stream of Tuaran River	17
3.2	An overview of Station 1	18
3.3	An overview of Station 2	18
3.4	An overview of Station 3	19
3.5	Raw materials that were used; cast net, hand gloves, meter rule, cool boxes, distilled water and plastic jars	20
3.6	10% formalin and 75% ethanol	21
3.7	The Hydrolab set and the GPS tracker	22
3.8	Handling Hydrolab for obtaining reading on the river physic- chemical parameters	24
3.9	The Fresh-water Fishes of North Borneo identification manual by Chin and Inger (2002).	25
3.10	Flowchart of the study	28
4.1	Sand mining activities observed at Station 2 and 3	31
4.2	The proportion of fish families	37
4.3	Ambassis interrupta	40
4.4	Arius maculatus	42
4.5	Barbonymus gonionotus	44

#### ABSTRACT

# DIVERSITY OF FRESHWATER FISHES AT LOWER STREAM OF TUARAN RIVER, SABAH

Sabah is rich with freshwater fishes and about 168 species can be found. Many localized studies of freshwater fishes focusing on areas such as Lahad Datu, Kota Marudu, Ranau and Kinabatangan had been done since the past few years, nevertheless, no further comprehensive studies have been made. When narrowing down the scope in Tuaran, the available information on the fish fauna at Tuaran River is only limited to locations of tagal and a few human activities such as sand mining have posed many threats to the aquatic living. This made it very difficult to gain any information related to the fish fauna available. The study was conducted with the purposes to provide morphological data and hence determine the diversity of freshwater fishes found at lower stream of Tuaran River, Sabah. All of the fish samples were captured by using cast nets along three selected stations. Throughout this study, a total of 225 individuals of 15 species and 12 families were captured which the family Cyprinidae gave the highest percentage. All of the fish specimens were measured and identified and the data were recorded for future references on the morphological data. The diversity index of the freshwater fishes ranged from 1.67 to 1.85. An exotic species Oxveleotris marmorata (Ketutu) which is the top five most expensive fish in Malaysia was also captured during this study which shows that the lower stream of the Tuaran River might be the potential habitat for this exotic species along with other valuable species such as Pangasius hypopthalmus (Patin) and Tilapia sp..