

VARIATION IN FIBER PROPERTIES WITH DIFFERENT HEIGHT LEVEL IN
KEDONDONG (*Dacryodes*)

By

ANAS BIN RIDUAN

Final Project Submitted in Partial Fulfillment for the Diploma in Wood Industries
Faculty of Applied Science
Universiti Teknologi MARA Pahang

MARCH 2004

ACKNOWLEDGEMENT

I would like to take this opportunity to express my special thank to ALLAH S.W.T for his blessing and favour in guiding me towards the complete my final project. As I expected, I'm glad that I was able to complete this project on time, as developments are taking place continuously, which adds my interest in this project and help me to make it such as a fascinating one.

First of all, I would like to express my special appreciation to my advisor Associate Prof. Dr. Suhaimi Mohammad for his encouragement, guidance and assisting in order to finish my research as final was compulsory to get my diploma.

Finally to every body who was involved directly and indirectly in the completion my final project, I wish success their life and thanks a lots.

TABLE OF CONTENTS

	Page
APPROVAL SHEET	i
DEDICATION.....	ii
ACKNOWLEDGEMENTS.....	iii
LIST OF TABLES.....	iv
LIST OF GRAPH.....	v
LIST OF ABBREVIATIONS.....	vi
ABSTRACT.....	vii
ABSTRAK	viii

CHAPTER	Page
I 1.0 INTRODUCTION.....	1-2
II 2.0 LITERATURE REVIEW.....	3
2.1 Field characteristic of Kedondong.....	3
2.2 Brief Diagnoses of Burseraceae Genera.....	3-4
2.3 Botanical explanation of <i>Dacryodes</i>	4-5
2.4 Uses	5
2.5 Importance of Fiber Properties	5
2.5.1 Fiber Length	5
2.5.2 Cell Wall Thickness	6
2.5.3 Runkle Ratio	6
2.5.4 Coefficient of suppleness	7
III 3.0 MATERIAL AND METHODS.....	8
3.1 Material Preparation	8-10
3.2 Determination of fiber morphology.....	11-12
Procedure.....	
IV 4.0 RESULT AND DISCUSSION.....	13-18
V 5.0 CONCLUSION	19
REFERENCES.....	20
APPENDICES.....	21-26
VITA.....	27

LIST OF TABLE

PAGES

TABLE 1.....	21-22
TABLE 2.....	23-24
TABLE 3.....	25-26

**Variation in Fiber Properties With Different Height Level In Kedondong
(*Dacryodes*)**

By

ANAS BIN RIDUAN

MARCH 2004

ABSTRACT

Variation of fiber morphology in Kedondong (*Dacryodes*) were ascertained. The variation of fiber properties of Kedondong with three different portions from three were also determined. The samples representing with base, middle and top of this tree. The result are indicated that the average of this threes for fiber morphology of Kedondong were suitable for pulp and paper industries according to their fiber morphology were achieved the target and the standard.