UNIVERSITI TEKNOLOGI MARA

HYPOGLYCEMIA: AWARENESS AND CHALLENGES FACED BY PATIENTS WITH DIABETES

SITI NURAISHAH BINTI HISHAMMUDIN

Dissertation submitted in partial fulfilment of the requirements for the degree of Bachelor of Pharmacy (Hons.)

FACULTY OF PHARMACY

JUNE 2014

ACKNOWLEDGEMENT

In the name of Allah S.W.T, the Most Gracious and the Most Merciful.

Alhamdulillah, all praises to Allah S.W.T who blessed me with health and strength to complete this research study. First and foremost, I take this opportunity to express my profound gratitude and deep regards to my supervisor, Dr. Mahmathi Karuppannan for her exemplary guidance, monitoring and constant encouragement throughout the course of this research study. Without her valuable help and guidance, this research study would not have been successful.

Sincere thanks to all my friends especially Ainul, Ayuni, Khaleyda, Izzati and others for their kindness and moral support in completing this research study. Thanks for the friendship and memories.

Last but not least, I would also like to thank my family who encouraged and fully supported me in every trial that came in my way. I thank them too for giving me not just financial, but moral and spiritual support.

To those who indirectly contributed in this research, your kindness means a lot to me. Thank you very much.

SITI NURAISHAH BINTI HISHAMMUDIN, JUNE 2014.

TABLE OF CONTENT

ACKNOWLEDGEMENT	4
LIST OF TABLES	8
LIST OF ABBREVIATIONS	9
CHAPTER ONE	10
INTRODUCTION	10
1.1 Overview	10
1.2 Problem statement	11
1.3 Rational of study	12
1.4 Objectives	12
CHAPTER TWO	13
LITERATURE REVIEW	13
2.1 Definition of Hypoglycemia	13
2.2 Epidemiology	13
2.2.1 Morbidity and mortality	15
2.3 Classification of Hypoglycemia	16
2.4 Signs and symptoms	17
2.5 Causal factors	19
2.6 Diagnosis	21
2.7 Treatment	22
2.8 Challenges	24
2.8.1 Hypoglycemia unawareness	24
2.8.2 Fear	25
2.8.3 Lack of knowledge	26

CHAPTER ONE

INTRODUCTION

1.1 Overview

One of the most feared complications of diabetes treatment is hypoglycemia. Regrettably, the threat and incidence of hypoglycemia case is increased day by day in efforts to accomplish normal blood glucose level among diabetic patients. Incorrect administration of insulin or anti-diabetic medications, excessive physical activity, skipping a meal or decreased appetite may cause drops in blood glucose level. Low blood glucose level can have an effect on all the organs in the body, especially the brain (McConnell, 2007). There may be barriers in the management of this condition and this could potentially increase the risk on incidence and hence, affect patients' quality of life.

Possibly, hypoglycemia is a severe and deadly complication of diabetes treatment and become a major cause of morbidity as well as mortality (Cryer, 2008). If misdiagnosed, this complication can be life-threatening where it can lead to serious morbidity such as major vascular event like stroke, acute cardiac failure, myocardial ischemia, myocardial infarction and arrhythmias especially among diabetic patients (Frier, 2008). Additionally, based on the study made before, the fear of hypoglycemia could get worse because of strict metabolic control implementation and this issue should need further research for finding a solution (Diabetic Hypoglycemia, 2008).

Other author suggested that, the cause and impact of hypoglycemia unawareness including factors such as psychosocial and self-management should need further research too (Thomas Kubiak, 2006). Furthermore, incidence of severe hypoglycemia in type 2 diabetes patients treated with insulin may be imprecise since many cases are treated by non-emergency medical services like family and friends at home or other than hospitals (Frier, 2008). Henceforth, the threat and inaccurate incidence of hypoglycemia become a major limiting factor for diabetes mellitus glycemic control (Briscoe & Davis, 2006).

1.2 Problem statement

The incidence of hypoglycemia unawareness is rising and frequency of untreated mild to moderate low blood sugar events may progress into severe event of hypoglycemia thus, become a threat. Recent studies also said that patients with Type 2 diabetes may develop hypoglycemia unawareness syndrome as found in the patients with Type 1 diabetes (Stephen Davis, 2004). This statement indicates that diabetic patients are prone to lose awareness towards symptoms of hypoglycemia in some cases and thus become a challenge for them to take care of themselves.

The rising of diabetes prevalence also raises the risk of developing hypoglycemia unawareness. The problem regarding to the diminished of hypoglycemia awareness level among diabetic patients can lead to morbidity and mortality if the matter is not solved.