UNIVERSITI TEKNOLOGI MARA

THE EFFECT OF HIGH TEMPERATURE AND pH ON RADICAL SCAVENGING ANTIOXIDANT ACTIVITY AND TOTAL PHENOLIC CONTENTS IN TUALANG HONEY

NOOR MARINAH BINTI SAHRI

Dissertation submitted in partial fulfillment of the requirements for the Bachelor of Pharmacy

Faculty of Pharmacy

JUNE 2014

ACKNOWLEDGEMENT

In the name of Allah, the Most Beneficent, the Most Merciful

Alhamdulillah, all praises to Allah for the strengths and His blessing in allowing the completion of this thesis. I am really grateful that I was able to finish this coursework successfully on time.

I would like to express my deepest appreciation to my supervisor, Dr. Syaliza Omar who has been very patient and helpful in guiding me from the beginning until I managed to complete this thesis. Her invaluable support and kindness will be remembered for a long time to come.

Besides that, I would like to extend my thanks to all master students and laboratory officers who helping me throughout laboratory work and demonstrated on how to use related instruments correctly. I also take this opportunity to give my thanks to all my acquaintances that have been directly or indirectly helping my works.

Last but not least, my gratitude goes to my beloved parents and family members for their prayers, endless love and encouragement. Without all of them, it might be impossible to finish this project. All I can say is thank you very much to everyone and may Allah repay you for kindness.

TABLE OF CONTENTS

TITLE P	AGE
APPROV	AL SHEET
	VLEDGEMENTii
TABLE OF CONTENTS	
	ABBREVATIONSv
LIST OF FIGURESvi	
ABSTRA	CTvii
СНАРТЕ	JR 11
INTRODUCTION	
1.1	Background of research
1.2	Problem statement
1.3	Significance of study
1.4	Research questions
1.5	Objectives4
CHAPTER 2	
LITERATURE REVIEW5	
2.1	Honey5
2.2	Composition of honey6
2.3	Medical use of honey
2.4	Antioxidant property of honey8
2.5	Phenolic compounds as antioxidant in honey9
2.6	Effect of heat and pH treatment to antioxidant activity11
2.7	Methods for testing antioxidant activity

ABSTRACT

Tualang honey has been selected for this study due to the health benefits and the possibility of it been added to foods and beverages. The objectives of this study are to determine the stability of antioxidant compounds in Tualang honey to high temperature and pH and to determine whether the phenolic compounds in Tualang honey contribute to antioxidant activity. Antioxidant activity of Tualang honey behaved varyingly after exposed to different temperature and pH. The stability of phenolic content at certain conditions leads to high antioxidant properties that give health beneficial to the consumer. Tualang honey has best antioxidant activity at neutral pH and at temperature 70 ©C. The exposure of Tualang honey toward these two conditions elevated the radical scavenging activity and correlated to the stability of phenolic content in Tualang honey. At elevated temperature and pH exposure, the antioxidant activity started to decline due to the changes in antioxidant compound behavior that responsible for antioxidant activity in Tualang honey. Antioxidant activity in Tualang honey also declined over time after exposure to different temperature due to degradation of some compounds in Tualang honey that act as radical scavenger.

CHAPTER 1

INTRODUCTION

1.1 Background of research

Free radicals are chemical species with an unpaired electron and they are very reactive (Devasagayam et al., 2004; McCord, 2000). Free radicals can be formed during metabolization of oxygen (Alzahrani et al., 2012). There are several types of free radical which are superoxide radical, hydroxyl radical, and peroxyl radical (Devasagayam et al., 2004).

The imbalance of free radicals production and antioxidants level will cause oxidative stress and lead to certain chronic diseases (Kishore et al., 2011; Mohamed et al., 2010). The examples of the chronic diseases are cardiovascular disease (Štajner et al., 2014), Alzheimer's disease (Wang et al., 2013) and diabetes (Saeidnia et al., 2013). Free radicals move through the cells and disrupted the structures of deoxyribonucleic acid (DNA). It also cause cross-linkage of proteins, formation of aging pigments, lipid peroxidation of the membrane cell and malfunction of mitochondria (Wickens, 2001).

The free radical effects can be neutralized by antioxidants. Antioxidant is the substance that capable to inhibit or delay the oxidation of the substrate (Tilak et al., 2004). Antioxidants occur naturally in human body (endogenous) or by consumed through diets