UNIVERSITI TEKNOLOGI MARA

PHYTOCHEMICAL STUDY OF LOPHOPETALUM JAVANICUM

SITI AISYAH BT ZALFAKHAR

Dissertation submitted in partial fulfilment of the requirements for the Bachelor of Pharmacy (Hons.)

Faculty of Pharmacy

ACKNOWLEDGEMENT

First and foremost, Alhamdulillah and thank you to Allah who allowed me to completed this thesis on time.

My special appreciation to my supervisor, Dr Humera Naz for her support, advice and generous guidance for completing this thesis. It was pleasurable and great working with her.

I am grateful to Miss Shazwani Shariffudin for her valuable advices and intellectual guidance during the laboratory work during this research. Besides, I would like to express my gratitude to all the staff of Atta-ur-Rahman institute for natural product discovery for their continuous support throughout this research work.

I also want to express my gratitude to my family and friends for their continuous support and urged me to do better. There are my sources of motivation. Last but not least, I would also like to thank to everyone who help me directly or indirectly throughout the process of this thesis.

TABLE OF CONTENTS

APPROVAL SHEET	
ACKNOWLEDGEMENTii	
TABLE OF CONTENTSiii	
LIST OF TABLEv	
LIST OF FIGURESv	
LIST OF ABBREVIATIONSvi	
ABSTRACTvii CHAPTER 1	
CHAPTE	ER 1
INTROD	OUCTION
1.1	Introduction
1.2	Research Question
1.3	Problem Statement
1.4	Significance of the study
1.5	Objectives of the study5
1.6	Hypothesis of the study
CHAPTER 26	
LITERATURE REVIEW6	
2.1	The family Celastraceae6
2.2	Lophopetalum javanicum
2.3	Phytochemical study of Celastraceae family
2.4	Phytochemical study of <i>Lophopetalum</i> 27
2.5	Biological activities of Celastraceae family
2.6	The genus Lophopetalum
CHAPTER 3	
MATERIALS AND METHODS	
3.1	Materials
3 1 1	Plant material

ABSTRACT

Lophopetalum javanicum or also known as Perupok by Malaysian is commonly found in Laos, Thailand, Malaysia, Vietnam, India and Cambodia. The bark of this plant is traditionally used as a constituent of a dart poison and the timber is commonly used for paneling and plywood. This plant is a member of Celastraceae family which have many biological activities such as stimulant, sedation, antibacterial, insect repellant and anti-tumour (Alarcón et al. 2015). Only a few species of the genus Lophopetalum have been subjected for phytochemical study thus Lophopetalum javanicum species is the most suitable candidate for this research. Fraction no. 4 of chloroform extract was selected and column chromatography of this fraction yielded seven sub fractions. Out of seven sub fractions, fraction no. 5 was chosen for isolation and purification by using preparative TLC. Once the purity of the isolated compound was confirmed, the compound was subjected to ¹H-NMR for identification of the chemical constituent. However, from this research, NMR analysis only showed the presence of long chain alcohol.

CHAPTER 1

INTRODUCTION

1.1 Introduction

Medicinal plants are continuously used by millions of people as medicines for diseases treatment and maintaining the world health since ancient times. According to World Health Organization (WHO), there are approximately 80% people worldwide dependent on the medicinal plants for some facial expression. The number of higher plants presence worldwide are about 270,000 but only small number has been studied scientifically and properly for pharmacological use (Ansari et al. 2015). The primary source for bioactive organic compounds isolation is the plants chemical diversity. The plants are anticipated to supply certain bioactive organic compounds to help in the development of "lead compound" for a treatment of diseases.

The Celastraceae species supply a large number of bioactive metabolites and it is widely used in the traditional medicine (Núñez et al. 2004). *Lophopetalum javanicum* is a plant species that is classified in the Celastraceae family. The bark is commonly used as the dart poison constituent (Md. Reyad-ul-Ferdous et al. 2014). By proper adjustment and