UNIVERSITI TEKNOLOGI MARA

DIVERSITY OF ENDOPHYTIC FUNGI ISOLATED FROM ASYMPTOMATIC COCOA POD AND THEIR POTENTIAL AGAINST Phytophthora palmivora

NUR FARAHIN BINTI SAHAT

Final year project report submitted in partial fulfilment of the requirement for the degree of Bachelor of Science (Hons.) Plantation Technology and Management

Faculty of Plantation and Agrotechnology

January 2015

DECLARATION

This Final Year Project is a partial fulfilment of the requirements for a degree of Bachelor of Science (Hons.) Plantation Technology and Management, Faculty of Plantation and Agrotechnology, Universiti Teknologi MARA.

It is entirely my own work and has not been submitted to any other University or higher education institution, or for any other academic award in this University. Where use has been made of the work of other people it has been fully acknowledged and fully referenced.

I hereby assign all and every rights in the copyright to this Work to the Universiti Teknologi MARA ("UiTM"), which henceforth shall be the owner of copyright in this Work and that, any reproduction or use in any form or by any means whatsoever is prohibited without a written consent of UiTM.

Candidate's signature : Date: Name:

I hereby declare that I have checked this project and in my opinion, this project is adequate in terms of scope and quality for the award of the degree of Bachelor of Science (Hons.) Plantation Technology and Management, Faculty of Plantation and Agrotechnology, Universiti Teknologi MARA.

Signature:
Name of Supervisor:
Position:
Date:

TABLE OF CONTENT

D) D) A(L) L) L) A) A)	EDICATION ECLARATION CKNOWLEDGEMENT ABLE OF CONTENT IST OF TABLES IST OF FIGURES IST OF ABBREVIATIONS BSTRACT BSTRAK	ii iii iv v vii viii x xi xiii	
~		1	
1	1 Packground of Study	1	
1.	2 Problem Statement	2	
1	3 Research Question	3	
1.	4 Objective Of Study	3	
1.	5 Significance of Study	4	
CHAPTER TWO LITERATURE REVIEW			
2.	1 Black Pod Rot (BPR) of Cocoa	5	
	2.1.1 Causal Pathogen	5	
	2.1.2 Life Cycle of the Pathogen	5	
•	2.1.3 Disease symptoms	8	
2.	2 Available Control Methods	9	
	2.2.1 Chemical Control	10	
	2.2.2 Cultural Practices	11	
	2.2.3 Biological Control	11	
2	2.2.4 Integrated Fest Management	12	
2.	2.3.1 Introduction of Endonbytic	13	
	2.3.1 Introduction of Endophytic. 2.3.2 Potential of endophytic fungi as biological control	14	
	21012 I Otomina of endopsiyae samp as offologieas constor		
CHAPTER THREE MATERIALS AND METHODS 15			
3.	1 Finding Endophytic Fungi from the Isolation Cocoa Pod	15	
	3.1.1 Pod Sampling	15	
	3.1.2 The Detection of endophytic fungi	16	
	3.1.3 Isolation of endophytic fungi	17	
	3.1.4 Statistical analysis	20	
	5.1.8 In Vitro Screening of Endophytic Fungi against P.	01	
	3 1 0 Culture P. nalmiyora	21	
	3.1.7 Culture F. paintivora	22	
	3 1 11 Culture Filtrate	25	
	STITE Outdre Finance	20	

ABSTRACT

DIVERSITY OF ENDOPHYTIC FUNGI ISOLATED FROM ASYMPTOMATIC COCOA POD AND THEIR POTENTIAL AGAINST Phytophthora palmivora

By

NUR FARAHIN BINTI SAHAT

January 2015

Chairman : Dr. Zaiton Binti Sapak

Faculty : Plantation and Agrotechnology

The cocoa (*Theobroma cacao L*) is the most important crop in Malaysia, producing an average yield of approximately 9,249 tonnes of cocoa beans and 33.1% of total production. Malaysian Cocoa Board has 13 stations to collect daily prices information: Tawau, Lahad Datu and Tenom in Sabah, Sabak Bernam, Bagan Datoh, Ayer tawar, Batu Pahat, Muar and Jerantut in Peninsular Malaysia and Kuching, Sarikei and Sibu in Sarawak. In Malaysia, the crop is hallowed by being basically disease free, suffering from but one major disease, Black Pod Rot (BPR), caused by *Phytophthora palmivora*. This disease can cause major loss in cocoa production and have a great effect to economic importance to the Malaysian Cocoa industry.

Endophytic fungi are able to play important functions in their plant hosts. The early studies with endophytic fungi were conducted in laboratory by isolating the cocoa pod .The use of endophytic fungi should be preferable as one of biological control agents as they are internal colonizers that are able to compete with *P. palmivora*. The potential of endophytic fungi colonization for protection against disease has given possibility for research. Understanding the life stage of endophytes can help to design management controlling of BPR. The objectives of this study were, (i) to evaluate the potential of isolate endophytic fungi from asymptomatic cocoa pod and (ii) to determine the potential of endophytic fungi as biological control against *P. palmivora*.

Endophytic fungi were detected and isolated from the pods of cocoa on different areas at different ages. Observations of pod sections under a light microscope showed the presence of fungal endophytes, stained blue with Lactophenol cotton blue (LCB). The fungal cells were observed intra and intercellular in pods from asymptomatic cocoa.There was significant difference between the numbers of Potential Endophytic Fungi found in Malaysian Cocoa Board, Jengka Pahang (MCBJP) with 30% than endophytic isolated from cocoa pod collected from the Malaysian Cocoa Board, Hilir Perak (MCBHP) with 70%. Endophytic fungus more prominent in immature cocoa pods as compared to mature for both collected fungi samples of MCBJP and MCBHP.

The isolated fungi were screened in vitro for their potential properties towards *P. palmivora*. Three fungal endophytes coded as F3, F6 and F20 were shown to have potential as bio control agents based on their percentage inhibition of radial growth (PIRG) in dual culture and culture filtrate. The PIRG of F3, F6 and F20 from dual culture are 83.06%, 63.29% and 68.47%. Respectively, the PIRG for culture filtrate for F3, F6 and F20 are 87.76%, 72.08% and 89.33%. Based on PIRG, F3 has the high potential to control *P. palmivora* disease on cocoa.