REMOVAL OF CHROMIUM (VI) FROM AQUEOUS SOLUTION BY BIOSORPTION OF RICE HUSK (RH) AND MODIFIED RICE HUSK (MRH)

RAZIN SAAD MD NOR

Final Year Project Report Submitted in Partial Fulfillment of the Requirements for the Degree of Bachelor of Science (Hons.) Applied Chemistry in the Faculty of Applied Sciences Universiti Teknologi MARA

MAY 2010

This final year project report entitled "**Removal of Chromium (VI) from Aqueous Solution by Biosorption of Rice Husk (RH) and Modified Rice Husk (MRH)**" was submitted by Razin Saad Md. Nor, in partial fulfillment of the requirements for the Degree of Bachelor Of Science (Hons.) Applied Chemistry, in the Faculty of Applied Sciences, and was approved by

Puan Zitty Sarah Ismail

Supervisor

B.Sc.(Hons.) Environmental Chemistry

Faculty of Applied Sciences

Universiti Teknologi MARA

40450 Shah Alam

Selangor

Ms. Sabring M. Yahaya Project Coordinator B.Sc.(Hons.) Applied Chemistry Faculty of Applied Sciences Universiti Teknologi MARA 40450 Shah Alam Selangor

Dr. Siti Halimah Sarijo Head of Programme B.Sc. (Hons.) Applied Chemistry Faculty of Applied Sciences University Teknologi MARA 40450 Shah Alam Selangor

ACKNOWLEDGEMENTS

In the name of Allah, the Most Gracious, the Most Merciful,

All the praise to Allah, for His Mercy has given me patience and strength to complete this work.

My heartfelt thanks go to Puan Zitty Sarah Ismail, for accepting me as her student project and for her sincere advice, time, guidance, patience and understanding within the one year of completing this thesis.

I am also indebted to Faculty of Applied Sciences; especially to member of Applied Chemistry program including lectures, laboratory assistants, and others staff for the technical support and guidance upon completing the project experiments.

I also would like to express my gratitude to my fellow friends, for their understanding, moral support, and all the sweet memories that we shared while doing thesis project together in the laboratory.

Last but not least, my utmost appreciation to my family for their support; moral and financial, and understanding.

Razin Saad Md Nor

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATIONS	viii
ABSTRACT	ix
ABSTRAK	Х

CHAPTER 1 INTRODUCTION

1.1	Background	1
1.2	Problem statement	4
1.3	Significant of study	6
1.4	Objective	8
1.5	Scope of study	9

CHAPTER 2 LITERATURE REVIEW

2.1	Heavy	Heavy metal pollution		
2.2	.2 Chromium		11	
	2.2.1	Properties of chromium	13	
	2.2.2	Carcinogenicity of hexavalent chromium	14	
	2.2.3	Sources and pathway of hexavalent chromium	16	
2.3	Biosorption		18	
	2.3.1	Defination	18	
	2.3.2	Biomass	19	
	2.3.3	Pretreatment of biomass	20	
	2.3.4	Effect of pH, biosorbent dose and contact time on biosorption	23	
2.4	Rice h	usks	28	
	2.4.1	Rice husks as biosorbent for heavy metal removal	30	
	2.4.2	Others application of rice husks in Malaysia	34	
		•		

ABSTRACT

REMOVAL OF CHROMIUM (VI) FROM AQUEOUS SOLUTION BY BIOSORPTION OF RICE HUSK (RH) AND MODIFIED RICE HUSK (MRH)

Presence of toxic heavy metal contaminants in aqueous streams is one of the major environmental issues these days. Chromium (VI) is one of the toxic heavy metals that have been major concern because it is carcinogenic and associated with many health hazards. Biosorption is the ability of certain biological material to remove the heavy metals from aqueous streams. Rice husks are the good low cost adsorbent as it contain 70 - 85% of organic matter (lignin, cellulose, sugars, etc). The adsorption of Cr (VI) from aqueous solution using raw rice husk (RH) and HNO₃ modified rice husk (MRH) was investigated in this study using the various parameter such as contact time, biosorbent dosage and pH at optimum condition. As the time increasing the percentage of Cr (VI) uptake was increased until it reach equilibrium time and the optimum uptake capacity of Cr (VI) for contact time parameter was found at 150 min for RH (57.9%) and 180 min for MRH (73.7%). For biosorbent dosage parameter, as the absorbent dosage increased the percentage of Cr (VI) removal was increased. The optimum biosorbent dosage for RH (68.4%) and MRH (82.3%) was found at 80 g/L. The optimum pH condition for RH (71.4%) and MRH (76.4%) was revealed as the highest removal of Cr (VI) was occurring at pH 2 and pH 4. The comparison for both RH and MRH for effectiveness of Cr (VI) removal was revealed as MRH has better performance than RH.