

UNIVERSITI TEKNOLOGI MARA

**WATER QUALITY, CLIMATIC
FACTORS AND PLANT TRAITS AS
PREDICTORS TO THE INSECT'S
ABUNDANCE IN THE PADDY FIELD**

NORAZLIZA BINTI ROSLY

Thesis submitted in fulfillment
of the requirements for the degree of
Master of Science


Faculty of Plantation and Agrotechnology

May 2015

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the result of my own work, unless otherwise indicated or acknowledged as referenced work. The thesis has not been submitted to any other academic institution or non-academic institution for any other degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student	:	Norazliza Binti Rosly
Student I.D No.	:	2010985549
Programme	:	Master in Science (Research)
Faculty	:	Plantation and Agrotechnology
Thesis Title	:	Water Quality, Climatic Factors and Plant Traits as Predictors to the Insect's Abundance in the Paddy Field
Signature of Student	:	
Date	:	May 2015

ABSTRACT

Paddy (*Oryza sativa* L.) was cultivated in many countries. The major problem in paddy field is insect pests. Due to that, pesticides were widely used. However, heavily usage can cause environmental pollution, loss of non-target pests, insecticides resistance and residual toxicity. Therefore, beneficial insects should be used as an alternative method. Understanding habitat preferences and factors that can reflect populations of beneficial insects are urgently needed. Following that, the intention of this study is to assess the environmental factors that reflect the abundance of beneficial insects in the paddy field at Sungai Burong, Tanjung Karang, Selangor. A total of 1,257 individuals of insects consisted of five orders (Odonata, Hemiptera, Coleoptera, Lepidoptera and Diptera), two sub-orders (Anisoptera and Zygoptera) and six families (Gerridae, Coccinellidae, Staphylinidae, Pyralidae, Cicadellidae and Chironomidae) found throughout two cropping periods. The compositions and diversity of the insects varied throughout the sampling period. The species richness and evenness was highest during vegetative phase. Besides, according to Mann-U Whitney and Kruskal-Wallis Tests, most of insects were influenced by localities of sampling areas. There were significance indirect relationship of insects with water temperature, Biochemical Oxygen Demand and Chemical Oxygen Demand. The plant height, insect pests, temperature, rainfall and humidity were good predictors on abundance of beneficial insects however, the R^2 values were relatively low due to significance indirectly relationship between water parameters with insects. Consideration of all factors, the total variance showed more than 78%, which indicate a strong correlation between those parameters insects in the paddy field.

TABLE OF CONTENTS

	Page
AUTHOR'S DECLARATION	ii
ABSTRACT	iii
ACKNOWLEDGEMENT	iv
TABLE OF CONTENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	x
LIST OF ABBREVIATIONS	xii
CHAPTER ONE: INTRODUCTION	
1.1 Background	1
1.2 Problem Statement	4
1.3 Significance of Study	5
1.4 Scope of Study	6
1.5 Objectives of Study	6
CHAPTER TWO: LITERATURE REVIEW	
2.1 Introduction to Paddy Cultivation in Malaysia	7
2.2 Taxonomy and Growth Stages	8
2.3 Ecosystem in Paddy Cultivation Area	15
2.4 Diversity of Insects in Paddy Area	17
2.5 Some Major Insect Pests and Damage	19
2.5.1 Stem Borers	21
2.5.2 Defoliators	23
2.5.3 Piercing-Sucking Insects	24
2.6 Predators in Paddy Fields	26
2.7 Pest Management	27
2.8 Application of Pesticides	28

2.9 Water and Paddy Cultivation	32
2.9.1 Acceptable Water Classes for Agricultural Purposes	34
2.9.2 Water Contamination due to Agricultural Activities	38

CHAPTER THREE: COMPOSITION, DIVERSITY AND DISTRIBUTION OF INSECTS IN PADDY FIELD

3.1 Background	42
3.2 Methodology	43
3.2.1 Location of Study	43
3.2.2 Sampling Design	45
3.2.3 Materials	45
3.2.4 Sampling Techniques	46
3.2.5 Statistical Analysis	46
3.2.6 Flowchart of Study	47
3.3 Results	48
3.3.1 Composition of Insects	48
3.3.2 Distribution of Insects	51
3.3.3 Diversity of Insects	53
3.4 Discussion	54
3.4.1 Composition and Diversity of Insects	55
3.4.2 Distribution of Insects	57

CHAPTER FOUR: INDIRECT RELATIONSHIP BETWEEN ABUNDANCE OF INSECTS WITH WATER PARAMETERS IN PADDY FIELD

4.1 Background	58
4.2 Methodology	59
4.2.1 Materials	59
4.2.2 Sampling Techniques	60
4.2.3 Sample Analysis	61
4.2.4 Statistical Analysis	62
4.2.5 Flowchart of Study	63
4.3 Results	68
4.3.1 The Water Quality Index of Water Sources for Paddy Field	68