

**ECTOPARASITES ON FRESHWATER FISH
Oreochromis niloticus Linnaeus, 1758 (TILAPIA MERAH)
IN FISH FARMS AT PENAMPANG, KOTA KINABALU, SABAH**

SITTY AIRA M.T AIDALA

**BACHELOR OF SCIENCE (Hons.) BIOLOGY
FACULTY OF APPLIED SCIENCE
UNIVERSITI TEKNOLOGI MARA**

JULY 2017

This Final Year Project Report entitled “**Ectoparasites on Freshwater Fish *Oreochromis niloticus* Linnaeus, 1758 (Tilapia Merah) in Fish Farms at Penampang, Kota Kinabalu, Sabah**” was submitted by Sitty Aira M.T Aidala, in partial fulfillment of the requirements for the Degree of Bachelor of Science (Hons.) Biology, in the Faculty of Applied Science, and was approved by

Ansir Bin Salim
Supervisor
B. Sc, (Hons.) Biology
Faculty of Applied Science
Universiti Teknologi MARA
88997 Kota Kinabalu
Sabah

Ajimi Bin Jawan
Project Coordinator
B. Sc, (Hons.) Biology
Faculty of Applied Science
Universiti Teknologi MARA
88997 Kota Kinabalu
Sabah

Siti Aminah Binti Tengah
@Mohammad
Programme Coordinator
B. Sc, (Hons.) Biology
Faculty of Applied Science
Universiti Teknologi MARA
88997 Kota Kinabalu
Sabah

Date: _____

ACKNOWLEDGMENTS

This thesis could not have been completed without a great deal of guidance and both mental and practical support. I would like to deeply thank those people who, during the several months in which this project lasted, provided me with everything I needed.

Foremost, I want to offer this endeavour to ALLAH Almighty for the wisdom He bestowed upon me, the strength, peace of mind and good health in order to finish this study. I would like to express deep gratitude to my supervisor Encik Ansir Salim for his guidance, encouragement and gracious support throughout the course of my work, for his expertise in the field that motivated me to work in this area and for his faith in me at every stage.

I would like to thank Encik Ajimi Jawan, my lecturer as well as the Project coordinator, for providing necessary support and permissions to conduct this study. My heartfelt thanks also goes to the lecturers who have been willingly to guide, correct and give opinions as well as help me to improve be better along my study period and along the project lasted. Besides, I also would like to thank the Laboratory assistants of KOMSAT, UiTM Kota Kinabalu for their help in the whole process of the project. I thank profusely Jabatan Perikanan Penampang, Kota Kinabalu, especially Pegawai Perikanan Penampang, Encik Tony Anus and his fellow assistants as well as the fish farmers who involved, for all their kind help and cooperation throughout my study and sampling period.

My thanks and appreciation also goes to my classmates especially, Estelle Lius, Marommie Roland Mecilus, Sherene Mylon, Debora Jolian and Maziera Safiza Mansur, who have willingly helped me out with their abilities especially along all the sampling period. I owe a deep sense of gratitude to my fellow project mates under parasitology study, Nur Deanna Kamislin, Shazanna Nur Atiqah Mulkan and Mohd Syahfiq Osman, for all the exchange of ideas, discussions and resources of references throughout the study.

My heartfelt thanks also go to my friends in UiTM Shah Alam, UiTM Sarawak, and Universiti Utara Malaysia (UUM), Abdul Rashid Bakri, Machairey Marcus and Khaidher Mohd Yasin for their unending support, advices and help to complete this study. Last but not least, with boundless love and appreciation it is my privilege to thank my family, my parents Mohd Taha Aidala, Siti Muslihat Mohd Satu and Zainab Janjani as well as my siblings for the encouragements and supports throughout my life.

Sitty Aira M.T Aidala

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vii
LIST OF FIGURES	viii
LIST OF ABBREVIATIONS	x
ABSTRACT	xi
ABSTRAK	xii
CHAPTER 1 INTRODUCTION	
1.1 Background of study	1
1.2 Significant of study	3
1.2.1 Fish management	3
1.2.2 Benefits to the society	3
1.2.3 Academic purposes	3
1.3 Objectives of study	4
CHAPTER 2 LITERATURE REVIEW	
2.1 Morphology and classification of <i>Oreochromis niloticus</i> (Tilapia Merah)	5
2.2 <i>Oreochromis niloticus</i> (Tilapia Merah) fish farming	7
2.3 Economic importance of freshwater fish <i>Oreochromis niloticus</i> (Tilapia Merah)	7
2.4 Environmental factors affects the fish	10
2.5 Water quality in fish farming of <i>Oreochromis niloticus</i> (Tilapia Merah)	10
2.6 Parasites of fish	12
2.6.1 Ectoparasites effects on fish	12
2.6.2 Ectoparasites infecting <i>Oreochromis niloticus</i> (Tilapia Merah)	13
2.7 Relationship between parasites infections with the sex and body size of the fish	14
2.7.1 Relationship between parasites infections with the sex of the fish	14
2.7.2 Relationship between parasites infections and the size body of the fish	16

ABSTRACT

ECTOPARASITES ON FRESHWATER FISH *Oreochromis niloticus* Linnaeus, 1758 (TILAPIA MERAH) IN FISH FARMS AT PENAMPANG, KOTA KINABALU, SABAH

Ectoparasites infection has been posing a threat to the fish cultivation activities in many fish farms, including Penampang area. As the matter of fact due to the ectoparasitic activity, the quality of the fish will be affected and often leads to death of fish resulting in enormous economic losses to the fish industry. The objectives of this study were to determine the type of ectoparasites on freshwater fish *Oreochromis niloticus* to determine the prevalence and intensity of ectoparasitic infections and to determine the relationship of both sex and body size with the parasite intensity of *Oreochromis niloticus* in fish farms at Penampang, Kota Kinabalu, Sabah. Sampling was done from February until June, 2017 with three times replications of sampling that range monthly per sampling. In this study, a total of 180 fish samples from two different fish farms, Putaton and Kipouvo were examined. The fish species examined was *Oreochromis niloticus* (Tilapia merah). Five Phylum of ectoparasites were found from the data observation and analysis. The ectoparasites were, skin blister roundworm from Phylum Nematoda, *Halacarid sp.* from the Phylum Acarina (mites), *Trichonida sp.* from the Phylum Protozoa, *Dactylogyrus sp.* and *Gyrodactylus sp.* from the Phylum Platyhelminthes Monogenea (Gillworms), and *Myzobdella sp.* from the Phylum Hirudinea (leeches). The prevalence and intensity of ectoparasite infection shows a significant different from both fish farms which in Putaton was slightly higher than Kipouvo. According to the statistical analysis, this study also shows that the relationship of the size of fish as well as the sex of the fish to the parasite intensity is negative (r value for the weight = - 0.104 and r value for the length = - 0.395). Throughout this study, it was observed that poor fish and water management were the main factors of the susceptibility of the fish to the parasitic infections. This study contribute to prepare data and management tool for future researchers in the aquaculture industry. As a recommendation, the distribution and abundance of ectoparasites on *Oreochromis niloticus* at Penampang should be further studied. It is also recommended that the fish farmers to continuously upgrading the water and fish management of the fish farms to maintain a hygienic and favourable condition for the fishes to be their source of economic as well as to be consumed by the society.