

**UNIVERSITI TEKNOLOGI MARA**

**REPEATED SPRINT AND  
DRIBBLING ABILITY AMONG  
MALAYSIAN UNIVERSITY SOCCER  
PLAYERS**

**ANNISAA BINTI BASAR**

Thesis submitted in fulfillment  
of the requirement for the degree of  
**Master of Science**

**Faculty of Sports Science and Recreation**

August 2015

## CONFIRMATION BY PANEL OF EXAMINERS

I certify that a Panel of Examiners has met on 25<sup>th</sup> February 2015 to conduct the final examination of Annisaa Binti Basar on her Master of Science thesis entitled "Repeated Sprint and Dribbling Ability among Malaysian University Soccer Players" in accordance with Universiti Teknologi MARA Act 1976 (Akta 173). The Panel of Examiners recommends that the student be awarded the relevant degree. The Panel of examiners was as follows:

Zulkifli Abd. Kadir  
Associate Professor  
Faculty of Sports Science and Recreation  
Universiti Teknologi MARA  
(Chairman)

Nor Anita Megat Mohd Nordin, PhD  
Associate Professor  
Faculty of Medicine, Physiology  
Universiti Kebangsaan Malaysia  
(Internal Examiner)

Zainal Abidin Zainuddin, PhD  
Associate Professor  
Faculty of Education  
Universiti Teknologi Malaysia  
(External Examiner)

**SITI HALIJJAH SHARIFF, PhD**

Associate Professor  
Dean  
Institute of Graduates Studies  
Universiti Teknologi MARA  
Date : 24<sup>th</sup> August, 2015

## ABSTRACT

The purpose of the present study was to determine the relationships between repeated sprints ability (RSA) and repeated dribbling ability (RDA) with aerobic power, leg power and blood lactate. Fifty two, well-trained soccer players (age:  $21.73 \pm 1.82$  years, height:  $171.31 \pm 5.27$  cm, weight:  $63.78 \pm 7.30$  kg, BMI:  $21.69 \pm 2.29$  kg.m<sup>-1</sup>) participated in this study were enrolled university students. All were members of top three teams (UiTM, UM, UKM) from the first division of Malaysian Higher Education Institution soccer league, 2013. Participants performed all four tests: 20 Multistages Shuttle Run Test, vertical jump, RSA, RDA and blood lactate concentration. The  $VO_{2max}$  was measured by 20MST, while leg power was measured by vertical jump. The RSA and RDA performance indices measured were total time (TT), fastest time (FT), mean time (MT) and performance decrement (PD). Both tests involved sprinting or dribbling 7 x 34.2 m with 25 seconds rest intervals (Bangsbo, 1994). Blood sample were collected from fingertip at the end of RSA and RDA tests (zero time recovery), and at 1, 3, 5, 7, 9, and 12 minutes of recovery. Significant differences were found in peak blood lactate and lactate removal rate for RSA and RDA ( $p < .05$ ). Low negative correlations were found among total time, mean time and  $VO_{2max}$  in RSA. While, moderate negative correlations were found among total time, fastest time and  $VO_{2max}$  in RDA. Significant positive correlations were found between performance indices (total time, fastest time, mean time) and lactate removal rate in RSA. As conclusion, the study suggests that  $VO_{2max}$  is correlated with selected RSA and RDA performance indices.

## TABLE OF CONTENTS

	<b>Page</b>
<b>CONFIRMATION BY PANEL OF EXAMINERS</b>	ii
<b>AUTHOR'S DECLARATION</b>	iii
<b>ABSTRACT</b>	iv
<b>ACKNOWLEDGEMENT</b>	v
<b>TABLE OF CONTENTS</b>	x
<b>LIST OF TABLES</b>	xi
<b>LIST OF FIGURES</b>	xii
<b>LIST OF ABBREVIATIONS</b>	xiii
<b>CHAPTER ONE: INTRODUCTION</b>	
1.1 Background of Study	1
1.2 Problem Statement	6
1.3 Purpose of the Study	7
1.4 Objectives of the Study	7
1.5 Hypotheses of the Study	8
1.6 Significant of the Study	9
1.7 Limitations of the Study	9
1.8 Delimitations of the Study	10
1.9 Study Assumptions	10
1.10 Operational Terms	10
1.10.1 Malaysian University Soccer Players	10
1.10.2 Repeated Sprints Ability (RSA)	10
1.10.3 Repeated Dribbling Ability (RDA)	11
<b>CHAPTER TWO: LITERATURE REVIEW</b>	
2.1 Introduction	12
2.2 Repeated Sprint Ability (RSA)	14

# CHAPTER ONE

## INTRODUCTION

### 1.1 BACKGROUND OF STUDY

Soccer is an intermittent sport whereby high intensity running interspersed with rest or submaximal work performed over long duration (90 minutes) (Alexander & Mier, 2011; Bishop & Edge, 2006; Bloomfield, Polman & O'Donoghue 2007; Sirotic & Coutts, 2007). Intermittent sports correlated with three main factors which are maximal oxygen consumption, anaerobic threshold and efficiency in the oxygen cost to generate high running speed (Michael, Joyner, & Edward, 2008). Due to the nature of the game, it can be characterized as aerobic sport performed close to anaerobic threshold (Bangsbo, Mohr, & Krstrup, 2006; Sirotic et al., 2007). Intermittent activities such as repeated sprint ability influenced both energy systems (Krstrup, Mohr, Steenberg, Bencke, Kjaer & Bangsbo, 2006; Spencer, Bishop, Dawson, & Goodman, 2005).

As mentioned earlier, soccer is a team sport which relies on high aerobic energy system (Bangsbo et al., 2006; Esposito, Impellizzeri, Margonato, Vanni, Pizzini, & Veicsteinas, 2004; Stolen, Chamari, Castagna, & Wisloff, 2005). It is largely represented by 42% walking, 20% standing, 16% jogging and 17% running, 1% sprinting and 4% other activities (Mohr, Krstrup, & Bangsbo, 2003). It also requires long distance running, which covers approximately 10,000 to 14,000 meters (Da Silva, Bloomfield & Marins, 2008) in 90 minutes game and almost 90% of energy expenditure is obtained by aerobic system (Bangsbo, 1994). High level of aerobic capacity indicates good cardiovascular endurance and fitness. It has been suggested that high  $VO_{2max}$  is required in enhancing anaerobic performance especially during intermittent activities as in repeated sprint ability (Bishop, Girard & Villanueva, 2011; Meckel, Machnai & Eliakim 2009).

Previous studies reported professional soccer players  $VO_{2max}$  could reach between 55 ml.kg.min<sup>-1</sup> to 67 ml.kg.min<sup>-1</sup> (Abrantes, Macas & Sampaio, 2004; Reilly, Bangsbo & Franks, 2000) equates between 12 to 15 in 20 meters Multistage Shuttle Run Test (MST) (Aziz, Tan & Teh, 2005). Aerobic capacity is also important in