# UNIVERSITI TEKNOLOGI MARA

# AGE STRUCTURE OF Sitophilus Oryzae USING GULTINOUS RICE, CALROSE RICE, THAI RED RICE AND PONNI RICE

### NUR SYAKIRAH BINTI MOHAMED ELIAS

Thesis submitted in fulfillment of the requirements for the degree of **Bachelor of Science (Hons.) Biology** 

**Faculty of Applied Sciences** 

January 2020

## **AUTHOR'S DECLARATION**

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Undergraduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student	:	Nur Syakirah Binti Mohamed Elias
Student I.D. No.	:	2016586999
Programme	:	Degree of Bachelor of Science (Hons.) Biology
Faculty	:	Applied Sciences
Thesis Title	:	Age Structure of <i>Sitophilus oryzae</i> using Gultinous Rice, Calrose Rice, Thai Red Rice and Ponni Rice

Signature of Student	:	
Date	:	January 2020

#### ABSTRACT

Stored product insect cause more than one fifth of total damages in rice in developing countries commonly and one of the insect that contributes to this problem is rice weevil Sitophilus oryzae. The huge amount of loss could be associated with their biology because their all four larval stage and pupal stage occurred inside the grain. Hence, causing their control and elimination are very difficult. Thus, this study focuses on the rice weevil S. oryzae age structure relationship with morphological characteristic based on the amount of nutrient used which might gain information on the preference and development of rice weevil S. oryzae. Through that, a new process can be developed later to protect the food grain in a better way. The tests were conducting by rearing this species in the four types of rice and were measured according to their age. Rice weevil S. oryzae were exposed to glutinous rice, calrose rice, thai red rice and ponni rice. Results shows there is no significant difference between each type, particularly on the 1<sup>st</sup> until 28<sup>th</sup> day range. However, the growth development of rice weevil S. oryzae are correlated with the types of nutrient used in this study. It is found that rice weevil S. oryzae which that reared in thai red rice show the fastest growth development compared to the other type of rice.

## **TABLE OF CONTENTS**

#### Page

CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR'S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENT	vii
TABLE OF CONTENTS	viii
LIST OF TABLES	ix
LIST OF FIGURES	X
LIST OF PLATES	xi
LIST OF SYMBOLS	xii
LIST OF ABBREVIATIONS	xii
LIST OF NOMENCLATURE	xiv

CHAPTER ONE: INTRODUCTION		1
1.1	Research Background	1
1.2	Problem Statement	2
1.3	Significance of Study	3
1.4	Objectives of Study	4

CHAPTER TWO: LITERATURE REVIEW		5
2.1	Introduction	5
2.2	Rice Weevils Sitophilus oryzae L.	6
2.3	Identification of Sitophilus oryzae	8
2.4	Distribution and Host of Sitophilus oryzae	8
2.5	General Biology of Sitophilus oryzae	
	2.5.1 Egg	9
	2.5.2 Larva	9
	2.5.3 Pupa	9
	2.5.4 Adult	10
2.6	Food Source of Sitophilus oryzae	11

2.7	Sitophilus oryzae Role as a Vector of Disease	11
2.8	Population study	12
2.9	Age Structure	14
	2.9.1 Life Table	14
CHA	APTER THREE: RESEARCH METHODOLOGY	16
3.1	Materials	16
	3.1.1 Raw Material	16
	3.1.2 Chemicals	16
	3.1.3 Apparatus	16
3.2	Methods	17
	3.2.1 Preparation of Apparatus	18
	3.2.2 Preparation of Material	19
	3.2.3 Collection of Sitophilus oryzae	19
	3.2.4 Identification of Sitophilus oryzae	20
	3.2.5 Rearing of Sitophilus oryzae	20
	3.2.6 Measuring The Morphological Data Characteristics	20
3.3	Statistical Analysis	21
3.4	Experimental Design	23
CHA	APTER FOUR: RESULTS AND DISCUSSION	24
4.1	The Growth Pattern in Body Parts Structure of Sitophilus oryzae	24
4.2	Comparison of P-value Based on Type of Rice	29
4.3	Factors That Influence the Age Structure of Sitophilus oryzae	31
CHA	APTER FIVE: CONCLUSION AND RECOMMENDATION	33
REF	ERENCES	34
APP	ENDICES	39
AUT	THOR'S PROFILE	47

11

2.6.1 Nutrient Content