UNIVERSITI TEKNOLOGI MARA

Canal at Sawah Sempadan Ricefield and Its Potential Health Risk" is a present

Assessment of Imazapic Residues in Surface Water and Drainage Canal at Sawah Sempadan Ricefield and Its Potential Health Risk

Mohamad Izhharuddin Bin Ismail

Project submitted in fulfillment of the requirements for the degree of

Bachelor in Environmental Health and Safety (Hons.) Faculty of Health Sciences

JULY 2015

DECLARATION

Project entitled "Assessment of Imazapic Residues in Surface Water and Drainage Canal at Sawah Sempadan Ricefield and Its Potential Health Risk" is a presentation of my original research. Wherever contributions of others are involved, every effort is made to indicate this clearly, with due reference to literature and acknowledgement of collaborative research and discussions. The project was done under guidance of Associate Professor Dr Hazilia Bt Hussain as project supervisor. It has been submitted to the Faculty of Health Sciences in partial fulfillment of requirement for the Degree of Environmental Health and Safety (Hons.).

Israel has Hashiff and my siblings for supports and encourage Students Signature,

(Mohamad Izhharuddin Bin Ismail)

Student ID: 2012279092

IC Number: 910328-07-5149

Date: 10th of June 2015

shown along this project period. Last but not least, I would lily

ACKNOWLEDGEMENT

In the name of Allah, all Praise is due to Him, the Sustainer of the Heavens and Earth and all that is within it and may His Blessings be upon the Prophet Muhammad SAW, peace be upon him. Alhamdulillah thanks to Him so that I am able to finish my final year project according to the scheduled.

Here I would like to take this opportunity to express my highest gratitude to my supervisor, Associate Professor Dr. Hazilia Bt Hussain of her guidance, support and advice that had been given along the project duration. Without her guidance this project may not be able to complete successfully.

I also want to dedicate my appreciation to Mr Erdzuam B Abd Rashid, Mr Muhammad Azwat Bin Abdullah and all postgraduate students that assists me during my laboratory analysis works and also trained me on various kind of instrumentation.

Lastly, I would like to thanks a lot to my family members especially to my father, Mr Ismail Bin Hashim and my siblings for supports and encouragement that they had shown along this project period. Last but not least, I would like to thank my dearest friends for their continuous support and sharing their experience in making this project a success

TABLE OF CONTENTS

| Chapter | 2.9 Remembran metho Title | Page Number |
|---------|---------------------------------------|-------------|
| | | |
| | Declaration . | ii |
| | Acknowledgement | iii |
| | Table of contents | iv |
| | List of tables | vii |
| | List of figures | viii |
| | List of plates | ix |
| | List of abbreviations | x |
| | Abstract | xii |
| 1 | Background Study | |
| | 1.1 Background information | 18 |
| | 1.2 Problem statement | 3 |
| | 1.3 Objective | 4 |
| | 1.4 Hypothesis | 4 |
| | 1.5 Conceptual framework | 4 |
| | 1.6 Significance of study | 6 |
| | 1.7 Scope of study | 6 |
| | | |
| 2 | Literature Review | |
| | 2.1 Terminologies | 8 |
| | 2.2 Regulations | 9 |
| | 2.3 Pesticide | 9 |
| | 2.4 Pesticide practice in paddy field | 10 |
| | 2.5 Pesticide fate | 11 |
| | 2.6 Surface water | 12 |
| | | |

ABSTRACT

ASSESSMENT OF IMAZAPIC RESIDUES IN SURFACE WATER AND DRAINAGE CANAL AT SAWAH SEMPADAN RICEFIELD AND ITS POTENTIAL HEALTH RISK

MOHAMAD IZHHARUDDIN BIN ISMAIL

Introduction: Imazapic is an herbicide under imidazolinone herbicide. This group of herbicide has about the same physical and chemical characteristics. It has characteristics of low potential to volatize. This herbicide also has low application rate and have wide selectivity of cropping system. The half-life of imazapic herbicide is 232 days and it may contain crystalline silica which is a carcinogenic substance. Block C, Sawah Sempadan, Tanjong Karang was chosen as the study area due to extensive use of this herbicide.

Methodology: Total samples collected for this study was 66 samples comprise of 2 seasons. There were 6 sampling points and for each sampling points 1 liter of surface water was collected. The same sampling point was used to collect samples on day 0, 1, 3, 5, 7, 11, and 15 of herbicide application. Standard sampling method was used to collect samples from each sampling point. Then, all amber bottles were kept in the ice box with temperature less than 4°C. Next, the samples was filtered before undergo extraction. Samples were extracted by using 500mg per 6ml SPE cartridge Bond Elut C18. Then, sample analyte was analyzed using HPLC-UV/VIS Detector with a reversed-phase column. A set of questionnaires was prepared for health risk assessment. Results: There were 60.6% of samples detected with imazapic residue in season 1 while in season 2 it was 57.6%. Based on Mann-Whitney test the result showed the *p*-value 0.445 and Kruskall-Wallis test *p*-value 0.774. For health risk assessment calculations, hazard index for non-carcinogenic health effects of male is 0.287 and female 0.254. Meanwhile, for carcinogenic health effects hazard index of male is 9.2x10⁻³ and female 5.81x10⁻³.

Conclusion: It can be concluded that there was presence of imazapic residues in surface water but the imazapic residues will not pose significant health hazards as indicated from the value of the hazard index.

Keywords: imazapic, surface water, risk assessment, hazard index.