UNIVERSITI TEKNOLOGI MARA

DEMULSIFICATION AND RHEOLOGICAL CHARACTERIZATION OF LIGHT MALAYSIAN CRUDE OIL EMULSIONS USING ELECTROCOAGULATION PROCESS

TAJNOR SURIYA BINITI TAJU ARIFFIN

MSc

April 2020

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student : Tajnor Suriya Binti Taju Ariffin

Student I.D. No. : 2014653436

Programme : Master of Science (Chemical Engineering) – EH 750

Faculty : Chemical Engineering

Thesis : Demulsification And Rheological Characterization Of

Light Malaysian Crude Oil Emulsions Using

Electrocoagulation Process

Tajnor Suriya

Signature of Student

Date : 27 April 2020

ABSTRACT

To date, most of the world's ongoing demand for crude has focused on lighter grades. However, creation of oil emulsion due to mechanical strength and natural surfactant during production are undesirable. The challenges associated with transportation and refining posed by world's oil produces lead the researchers to probe into practical demulsification methods. A great understanding of crude oil rheology and operating condition to treat emulsion is necessary. The research deals with water-in-oil (W/O) emulsion with API gravity of 38.98. The objectives of this research are to study the rheological behaviour and demulsification rate of emulsion. The rheological behaviour of crude oil emulsion has been studied by manipulating the temperature, shear rate and water volume fraction. The rheological study was conducted within a range of temperature $(30 - 90^{\circ}\text{C})$, shear rates (0.1 - 100 s-1) and volume water fraction (20%,30% and 40%). For emulsion with different water content, the rheological studies follow non-Newtonian shear thinning behaviour and were described in better way by Herschel – Bulky model. Besides that, experimental results indicated that measured viscosity of emulsion decreases significantly with temperature (dropped from 41.0 cP to 27.9 cP) while increasing water volume fraction resulted in increase in viscosity (raised from 6.14 cP to 103 cP). The increasing of temperature cause the molecules loss the chance to agglomerate and thereby breaking the ordered structure thus reduce the viscosity. Meanwhile, higher water fraction of emulsion characterizes higher elastic behaviour, subsequently increase resistance to flow then resulted in higher viscosity. The demulsification rates of water in crude oil emulsion in direct current fields were investigated under various conditions by using electrochemical cell. The separation rate of water increased along with the applied field, water content and salt concentration. The results showed that emulsion separation was governed by the applied electric field, which yield high separation efficiency up to 85.75% at 30V. Moreover, as the salt concentration reached 40000 ppm, the separation percentage achieved 88.75% with shorter time by using copper as electrode. On the other hand, the highest separation achieved by using aluminium was 62.5%. Copper showed a better performance than aluminium. This is because, copper is good electricity conductor, thus it can produced more ions during electrolysis and enhanced the separation efficiency. The findings obtained from this study proved that the demulsification process of crude oil emulsion under electric field can have a high efficiency of separation when the water-in-oil emulsions have high water volume fraction (40%) and salt concentration (40 000ppm) by using copper electrode with 30V applied voltage.

ACKNOWLEDGEMENT

Firstly, I wish to thank Allah for giving me the opportunity to embark on my Master and for completing this long and challenging journey successfully. I would like to express my gratitude and thanks to my main supervisor, Dr Erfan Mohammadian and my co-supervisor, Puan Effah Yahya for their commitment and encouragement throughout the progress of this research.

My sincere thanks also go to Dr Hazlina Husin, my former supervisor, for her kindness and guidance for helping me with this research. I always indebted for all the knowledge shared with me.

Million thanks to Faculty of Chemical Engineering who provided all the facilities that required to conduct the research and to all staffs for giving guidance and assistance to handle several equipment and conducting the laboratory works.

Last but not least, a special thanks to my beloved parents, Encik Taju Ariffin and Puan Normah for their unlimited love and support from the first until the end. Not to forget, my dearest husband. Thank you for constantly being by my side, for being my best friend and critique.

Great pleasure to all my Postgraduate friends for helping me with this project. Especially, Nurfarisha Achim, Siti Nabihah, Nurul Aida Sulaiman and Sofia Sakina. I really appreciate the helps in term of sharing journal and other reference materials from all of you. Alhamdulillah.

TABLE OF CONTENT

			Page
CONFIRMATION BY PANEL OF EXAMINERS			ii
AUTHOR'S DECLARATION ABSTRACT ACKNOWLEDGEMENTS TABLE OF CONTENTS LIST OF TABLES LIST OF FIGURES LIST OF SYMBOLS LIST OF ABBREVIATIONS			iii
			iv v
			ix
			X
			xiii
			xiv
CH	APTE]	R ONE: INTRODUCTION	1
1.1	Resear	rch Background	1
1.2	Proble	m Statement	4
1.3	Resear	rch Objectives	6
1.4	Resear	rch Scope	6
1.5	Signifi	icant of Research	7
CH	APTE	R TWO: LITERATURE REVIEW	8
2.1	Introdu	uction	8
2.2	Light (Crude Oil	9
2.3	Emuls	ion Formation	10
2.4	Rheology of Emulsion		13
	2.4.1	Theory	13
	2.4.2	Newtonian versus Non – Newtonian Liquids	14
	2.4.3	Rheological Models	16
2.5	Factors Influencing the Stability of Emulsions		18
	2.5.1	Brine Salinity	18
	2.5.2	Temperature	19
	2.5.3	Mixing Speed And Mixing Duration	20