

**UNIVERSITI TEKNOLOGI MARA**

**THE ENHANCEMENT OF INSULIN  
SENSITIVITY BY STEVIOSIDE  
FROM *Stevia rebaudiana* Bertoni IN  
*IN-VITRO* AND *IN-SILICO* MODELS**

**NABILATUL HANI BINTI MOHD RADZMAN**

Thesis submitted in fulfilment  
of the requirements for the degree of  
**Doctor of Philosophy**

**Faculty of Pharmacy**

August 2017

## ABSTRACT

The adoption of unhealthy, sedentary lifestyles has triggered the prevalence of metabolic diseases like diabetes and obesity. A condition referred to as insulin resistance has been found to be the precursor to these diseases. It commonly manifests itself in most if not all Type 2 diabetic cases. A cure is yet to be found and side effects from current drugs create complications among patients. Thus, alternative therapies from natural, plant-based products like stevioside are becoming a more preferred option. Stevioside that is extracted from *Stevia rebaudiana* Bertoni has impeccable sweetening potential, which provides an interesting aspect to its proposed antidiabetic potentials. Hence, in-depth investigations were conducted to analyse how stevioside can manifest its effects towards insulin sensitivity in *in-vitro* and *in-silico* models. Like many herbal products, scientific data on stevioside's efficacies has been scarce. Its safety of consumption was hence tested through 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay on 3T3-L1 adipocytes as an *in-vitro* model. No IC<sub>50</sub> was observed as cell viability was only slightly reduced, signifying its non-cytotoxicity towards adipocytes. Preceding the assay, the cells were tested using Oil Red O to confirm differentiation which was also successfully achieved. Stevioside was observed to increase glucose uptake in adipocytes better than a drug both in normal and insulin resistant states based on the glucose uptake assay that was conducted. Through Western blotting, expression of the phosphorylated tyrosine (pY20) protein on the insulin receptor (IR $\beta$ ) was also observed to be enhanced by stevioside. This suggests that stevioside has a high probability of interacting with the insulin receptor in improving insulin sensitivity and increasing glucose uptake. Stevioside's actions are therefore, upstream rather than downstream of the insulin signalling pathway and were confirmed through computer simulations. Prior to that, a protein model was constructed using the MODELLER software. Multiple sequence alignment of the human and mouse insulin receptor sequences was first conducted through ClustalW. A human insulin receptor 3D structure with PDBID; 3LOH was selected as the protein template to model the *Mus musculus* insulin receptor. Subsequent docking of the stevioside ligand was conducted via AutoDock Vina and had managed to reveal possible binding sites. Interestingly, stevioside was observed to share the same binding region to that of insulin on the insulin receptor. Henceforth, insulin binding was analysed through radioimmunoassay (RIA) with radioactively tagged <sup>125</sup>I-insulin; quantified using a  $\gamma$ -counter. Stevioside was seen to reduce insulin binding but not as severe as S961 treatments; that are positive controls to insulin binding inhibition. In conclusion, stevioside enhances insulin sensitivity in adipocytes by increasing glucose uptake and enhancing expressions of pY20 on IR $\beta$  of the insulin signalling pathway. Computer simulations of insulin receptor-stevioside interactions have also revealed docking of stevioside onto a site shared by insulin binding on the receptor. This was confirmed through stevioside's reduction in total insulin binding analysed through RIA. Therefore, stevioside may have a role in manifesting its effects through the insulin receptor towards improving insulin sensitivity; upstream of the insulin signalling pathway by possibly binding to the insulin receptor.

## ACKNOWLEDGEMENTS

*“Surely with difficulty is ease. With difficulty is surely ease.”*  
(Holy Quran 94:5-6)

This thesis is produced as a result of five years of hard work and has been proven to be challenging in many ways. First and foremost, I would like to thank the Lord, Allah SWT for guiding me and for the strength that He has bestowed upon me to go through this journey. It is also befitting for me to express my gratitude to some of the most wonderful people with even more wonderful virtues that they have shown me in this journey of completing this thesis.

I would like to thank my supervisor Dr Wan Iryani Wan Ismail for her support and guidance throughout the making of this thesis, who have mentored me both in life and science. The same appreciation goes to my other advisors, Dr Zainah Adam, Dr Rosmadi Yusof and Ms Safura Jaapar for their undying support and encouragements. I would like to extend my heartiest gratitude to Dr Siti Azma Jusoh for mentoring me on the Bioinformatics aspects of the project. I would also like to thank the Ministry of Education Malaysia for awarding the grant that has funded this project.

Huge thanks to all past and present members of the Cell Signalling Research Group; Suhana, Francis, Nazri, Fadzilah, Hisham, Ashraf, Naim, Hamiyah, Aida, and Faiz. Of whom have contributed to this project be it by lending a helping hand for my experiments or just being there, helping me maintain my sanity in the lab throughout the years.

I would like to reserve my most heartfelt gratitude to my family, especially to my husband Daniel, and my parents, siblings and in-laws for believing in me and for putting up with me throughout this process. They have never failed to assure me of their unconditional love. Their profound affection and support is irreplaceable for it has shown me the boundless patience anyone could ever have.

Finally, I would like to dedicate this thesis to anyone who has carried on against all odds in order to do the things they are passionate for, especially in the name of science.

# TABLE OF CONTENTS

	<b>Page</b>
<b>CONFIRMATION BY PANEL OF EXAMINERS</b>	ii
<b>AUTHOR'S DECLARATION</b>	iii
<b>ABSTRACT</b>	iv
<b>ACKNOWLEDGEMENTS</b>	v
<b>TABLE OF CONTENTS</b>	vi
<b>LIST OF TABLES</b>	xi
<b>LIST OF FIGURES</b>	xii
<b>LIST OF ABBREVIATIONS</b>	xvi
<b>CHAPTER ONE: INTRODUCTION</b>	<b>1</b>
1.1 Research Background	1
1.2 Problem Statements	2
1.3 Research Questions	3
1.4 Research Objectives	3
1.5 Significance of Study	4
1.6 Scope of Study	5
<b>CHAPTER TWO: LITERATURE REVIEW</b>	<b>7</b>
2.1 Introduction	7
2.2 Diabetes mellitus	8
2.2.1 History of Diabetes	8
2.2.2 Classification of Diabetes	9
2.2.2.1 Type 1 Diabetes	10
2.2.2.2 Type 2 Diabetes	12
2.2.2.3 Gestational Diabetes	14
2.2.3 Factors Contributing to Diabetes	15
2.2.3.1 Free Fatty Acids	15

2.2.3.2	Tumour Necrosis Factor $\alpha$	16
2.2.3.3	Serine Kinases	17
2.2.4	Pathophysiology, Symptoms and Complications of Diabetes	17
2.2.5	Mechanisms of Diabetes Progression	19
2.2.5.1	Insulin Signalling Pathway	20
2.2.5.2	Disruptions to Insulin Signalling in Insulin Resistance	22
2.2.5.3	Insulin Receptor (InsR)	24
2.2.6	Diabetes Management, Solution and Treatment	26
2.2.6.1	Treatment for Diabetes Mellitus	
2.2.7	New Strategies for Noninvasive Insulin Delivery	35
2.3	3T3-L1 Cell Line	39
2.4	Plants with antidiabetic potentials	41
2.5	<i>Stevia rebaudiana</i> Bertoni	43
2.5.1	Background and Origin	43
2.5.2	Stevioside as a Bioactive Compound	45
2.5.2.1	Medicinal Benefits and Antidiabetic Properties of Stevioside	46
<b>CHAPTER THREE: RESEARCH METHODS</b>		48
3.1	Materials	48
3.2	Methods	48
3.2.1	Cell Line	50
3.2.2	Maintenance, Subculture and Seeding of Cells	50
3.2.3	Cryopreservation	51
3.2.4	Cryothawing	51
3.2.5	Adipocyte Differentiation	51
3.2.6	Induction to Insulin Resistance	52
3.2.6.1	Tumour Necrosis Factor $\alpha$ (TNF- $\alpha$ )	52
3.2.6.2	S961, Insulin Receptor Antagonist	52
3.2.7	Sample Preparation	52