# STUDY ON MORPHOGENESIS OF Solanum lycopersicum AND Capsicum annuum IN COMBINATION MEDIA CONTAIN NAA AND PLANT BOOSTER BY USING SEEDS CULTURE

## NURUL NADIA BINTI YAHYA

BACHELOR OF SCIENCES (Hons.) BIOLOGY FACULTY OF APPLIED SCIENCES UNIVERSITI TEKNOLOGI MARA

#### **ACKNOWLEDGEMENT**

Alhamdulillah and gratitude to the Almighty for giving me the strength and patience to complete this final year project. I would like to express my thanks you to my supervisor, Miss NurSuria binti Md Setamam because without her guide my research study cannot be done properly. She always gives support and guide to complete of this final year project.

In addition I would like to take this opportunity to thanks those of friends who helped me and giving some guide to complete this project. Their views and tips are useful indeed. Last but not least, I am grateful for having such as supportive to all my family members and UiTM staffs especially staffs laboratory in order to accomplish my project. My Allah blesses each and every one of you.

### TABLE OF CONTENT

			PAGE	
ACK	NOWL	EDGEMENT	iii	
	TABLE OF CONTENT			
	LIST OF TABLES			
	OF FIG		vii	
LIST	OF AB	REVIATIONS	viii	
	TRACT		ix	
ABS	TRAK		x	
СНА	PTER 1	: INTRODUCTION		
1.1	Backg	round study	1	
1.2		em statement	3	
1.3	Signif	icance of the study	3	
1.4	_	tives of the study	4	
1.1	Object	arves of the study		
СНА	APTER 2	2: LITERATURE REVIEW		
2.1	Solani	um lycopersicum and Capsicum annum	5	
	2.1.1	Nutrient and Secondary Metabolite	9	
	2.1.2	Medicinal value	10	
2.2	Plant t	tissue culture	12	
2.3	Morph	nogenesis	13	
2.4	Seed o			
2.5	Callus	Callus		
2.6	Plant 1	Plant booster		
2.7	Napthaleneatic acid (NAA)		16	
		3: METHODOLOGY		
3.1	Mater		17	
		Raw materials	17	
		Material	17	
		Apparatus	17 17	
3.2		Method		
	3.2.1		17	
		Hormone stock solution preparation	18	
	3.2.3			
		Preparation media for all treatment	21	
	3.2.5	1 1	21	
	3.2.6	Sample preparation	24	

CHAP	TER 4: RESULT AND DISCUSSION		
4.1	Effect of seed germination in difference concentration		
	of NAA in media that contain plant booster.	26	
4.2	Effect of Auxin and plant booster in plant growth of		
	Capsicum annuum and Solanum lycopersicum	31	
4.3	Effect of callus present in media that contain Auxin and	34	
	plant booster		
CHAP	TER 5: CONCLUSION AND RECOMMENDATION	41	
	TER 3. CONCEDESTON THE RECOMMENDATION	-11	
CITE	CITED REFERENCES		
APPE	NDICES	46	
CURI	CULUM VITAE	48	

#### ABSTRACT

# STUDY ON MORPHOGENESIS OF Solanum lycopersicum AND Capsicum annuum IN COMBINATION MEDIA CONTAIN NAA AND PLANT BOOSTER BY USING SEEDS CULTURE.

Solanum lycopersicum has a much lower sugar content compare to other edible fruits and considered as vegetables for most culinary uses while Capsicum annuum has been used since ancient times as food flavoring and for human health. Chilies have numerous chemicals including steam-volatile oil, fatty oils, capsaicinods, vitamins, protein, fibre and mineral elements. In this research study the auxin hormone that been used is Napthaleneacetic acid (NAA) is a plant hormone that that stimulate elongation and involved in biological activity. While, MARDI plant booster that are used in this research study were consist organic supplements that has possibility also towards the growth and development of plants. In this research, effect of seed germination, plant growth and callus formation as an objective. There are total five treatment which are treatment A (control), B (1ml/L NAA), C (10ml/L plant booster), D (10ml/L plant booster and 1ml/L NAA) and E (10ml/L plant booster and 2ml/L NAA). Treatment B was shows a better result of Capsicum annuum as 3.6cm ±std and Solanum lycopersicum as 4.6cm ±std compare to another treatment. For Callus formation result treatment D was showed higher value of mean which is 1.4cm for Capsicum annuum and 1.6cm for Solanum lycopersicum. From the result, it was showed the NAA is good for plant growth compare plant booster because in NAA have their own nutrient that help plant to grow in short period. For recommendation can used NAA for plant growth in future.